

9444-189

1/29/2002

1/3

Pyrethrins Are Made From Chrysanthemum Flowers

CB-40-1 for Insect Control

- Provides Rapid Control of Residential, and Industrial Pests
- Excellent Flushing • 4 to 1 Synergist Ratio
- Quick Knockdown • Nonaqueous
- Highly Active Solvent System

KILLS and REPELS: Flies (Fruit Flies, Horn Flies, House Flies, Cluster Flies and Stable Flies) Mosquitoes, Gnats, Small Flying Moths, Wasps, Hornets, Mud Daubers, Cockroaches, Ants, Fleas, Crickets, Drug Store Beetles, Spiders, Silverfish, Sowbugs, Indianmeal Moths, Mediterranean Flour Moths, Angoumois Grain Moths, Confused Flour Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary, Weevils, Grain Mites, Booklice, Carpet Beetles, Centipedes, Chocolate Moths, Cigarette Beetles, Clover Mites and Bedbugs

FOR USE in Apartments, Campgrounds, Homes, Offices, Hotels, Motels, Schools, Hospitals, Nursing Homes, Resorts, Day-care Centers, Factories, Utilities, Warehouses, Stores, Supermarkets, Dining Areas and Pantries of Restaurants, Food Handling Establishments, Commercial or Institutional Kitchens, Food Processing Plants, Meat Packing Plants, Bottling Plants and Transportation Equipment (Automobiles, Buses, Trucks, Trailers, Boats, Ships, Trains, Aircraft¹)

1-Do not use in aircraft cabins.

CONTAINS NO CFCs OR OTHER OZONE DEPLETING SUBSTANCES. FEDERAL REGULATIONS PROHIBIT CFC PROPELLANTS IN AEROSOLS.

ACTIVE INGREDIENTS:

| | | |
|-------------------|--------------------------------------|---------------|
| | Pyrethrins | 0.50% |
| * | Piperonyl Butoxide, technical | 1.00% |
| | N-octyl bicycloheptene dicarboximide | 1.00% |
| INERT INGREDIENTS | | <u>97.50%</u> |
| | | 100.00% |

- * Equivalent to 0.80% (butylcarbityl) (6-propylpiperonyl) ether and 0.20% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel for Additional Precaution

EPA Reg. No. 9444-189

EPA Est. No. 9444-LA-1

NET WEIGHT 20 oz / 567 g

RECEIVED
JAN 29 2002

U.S. DEPARTMENT OF AGRICULTURE
NATIONAL SERVICE FOR
PEST MANAGEMENT
EPA 9444-189

2/3

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Wash thoroughly with soap and water after handling. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Harmful if inhaled. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not apply this product in or on electrical equipment due to the possibility of a shock hazard.

Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

Do not contaminate food or foodstuffs. Do not remain in treated areas. Vacate premises for at least 15 minutes after spraying. Ventilate the area after treatment is completed and before re-entering. In medical care facilities, remove all patients prior to treatment. Ventilate area for 2 hours before returning patients.

FLYING INSECTS: Flies, (Cluster Flies, Fruit Flies, Horn Flies, House Flies, Stable Flies,)

Mosquitoes, Mud Daubers, Small Flying Moths, Wasps, Hornets, Gnats—Calculate cubic feet of space to be treated. Close all doors and windows. Spray in a sweeping motion to all parts of the room. Apply for 1-3 seconds per 1,000 cubic feet of space. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Sweep up and destroy fallen insects. Repeat as necessary.

CRAWLING INSECTS: Ants, Booklice, Centipedes, Clover Mites, Cockroaches, Crickets, Silverfish, Sowbugs, Spiders, -General Volumetric Treatment: Open cabinets and doors in area to be treated. Shut off air conditioners, fans, and close doors and windows before spraying. Spray for 20 seconds per 1,000 cubic feet of space. Direct spray toward areas suspected of harboring insects. Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment as necessary.

FOR COCKROACHES: For best results, spray directly on cockroaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind stoves and refrigerators, food storage areas and wherever these pests are suspected of hiding. Also spray around door- and windowframes. Repeat as necessary.

Ants, Booklice, Centipedes, Clover Mites, Crickets, Fleas, Millipedes, Roaches, Silverfish, Sowbugs and Spiders: Inject into cracks and crevices in all hiding places behind baseboards, sinks, cabinets, meter boxes, door frames, windows. **GENERAL VOLUMETRIC TREATMENT:** Open cabinets and doors in areas to be treated. Turn off air conditioners and fans and close doors and windows before treating. Apply at the rate of 20 seconds per 1000 cu. ft. of space. Disperse toward area suspect of harboring the greatest insect infestations. Disperse in all locations contacting as many insects as possible. Keep area closed for 15 minutes. Open and ventilate the treated area before reoccupying. Repeat treatment as necessary.

CRACK & CREVICE APPLICATION: Food areas and/or food handling establishments, spray in cracks and crevices using the crack and crevice nozzle provided. A crack and crevice application can be done in food areas and/or food handling establishments while these facilities are in operation by taking extreme care not to contaminate surfaces.

VOLUMETRIC TREATMENT IN FOOD AREAS AND/OR FOOD HANDLING ESTABLISHMENTS

ESTABLISHMENTS: Food Areas of Food Handling Establishments: When applying a space treatment in food handling areas, the food handling operation should be shut down. Foods should be removed or covered during treatment. In food processing areas, all utensil, shelving, etc., where food will be handled should be covered or removed before treatment or thoroughly washed with an effective cleaning compound followed by a potable water rinse prior to use. Careful crack and crevice application for inspection purposes is permissible at all times.

VOLUMETRIC TREATMENT IN HOMES: In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Remove pets and birds and cover fish aquariums before spraying.

VOLUMETRIC TREATMENT FOR INACCESSIBLE VOIDS (Attics, False Ceilings, Wall and Equipment Voids): For crack and crevice application, calculate void volume in cubic feet. Treat 20 seconds per 1,000 cubic feet. Repeat as necessary.

FOR SPIDERS AND SILVERFISH: For control of these insects, frequent and repeated spraying of baseboards, windows, doorframes, bookcases and storage areas should be made. For best results, spray directly on insects.

FOR CRICKETS: Spray thoroughly and frequently behind stoves, refrigerators, under cabinets, in food storage areas and especially in closets, basements and other damp areas. For best results, spray directly on insects.

FOR FLEAS IN BUILDINGS: Close windows and doors. Hold aerosol 36 inches above floor and direct fog toward floor and lower walls at a rate of 10 seconds per 100 square feet making sure that all floor, sofa, and chair surfaces are contacted. Repeat treatment after 7 days.

FOR ANTS: Spray ant trails, nests and points of entry. For initial clean-up, apply for 10-15 seconds per 100 square feet of floor area. Then periodically spray as a space aerosol 3-5 seconds per 100 cubic feet of space.

INDIANMEAL MOTHS, CHOCOLATE MOTHS, MEDITERRANEAN FLOUR MOTHS, ANGOUMOIS GRAIN MOTHS: Close doors and windows, direct spray into all parts of room, especially around stored food product containers, pallets, and darkened areas. Do not spray in areas where food is exposed in anyway. Spray 7 to 10 seconds for each 1,000 cubic feet of space. Keep area closed for 15 minutes following treatment. Do not remain in treated areas. Ventilate after treatment. Repeat application weekly or as needed.

Exposed Adult and Larval stages of Bedbugs, Carpet Beetles, Cigarette Beetles, Confused Flour Beetles, Drugstore Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites: Spray into cracks and crevices, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.

CARPET BEETLES: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

FOR BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary.

HOME USE: To Kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible, and spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves and under appliances. Repeat as necessary.

Applications in Aircraft: This product can be applied in cargo areas but not in aircraft cabins.

FOOD PROCESSING PLANT USE: To Kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible. Spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves, under appliances, in food storage areas.

OUTDOOR GROUND APPLICATION: To Kill Flies, Gnats and Mosquitoes in open areas near buildings and campgrounds. Best results are obtained when wind speed is five (5) MPH or less. Apply at a rate of 1.5 to 2 seconds per thousand square feet. Apply in wide swaths across area to be treated. Allow treatment to penetrate dense foliage. Repeat treatment as necessary.

Authorized for use under USDA Inspection and Grading Programs.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area away from heat or open flame.

DISPOSAL: This container may be recycled in the few but growing number of communities where steel aerosol can recycling is available. Before offering for recycling, empty the can by using the product according to the label. **DO NOT PUNCTURE!** If recycling is not available, wrap the container and discard in the trash.

Manufactured by:
Waterbury Companies, Inc.
P.O. Box 1812
Waterbury, CT 06722
Made in USA