

NOV 22 2005

Ms. Cynthia L. Rohner  
Waterbury Companies, Inc.  
P.O. Box 640/129 Calhoun Street  
Independence, LA 70443

Subject: Label Amendment for Additional Marketing Claims  
Country Vet Farm & Dairy CV-80D Insect Control  
EPA Reg. Number: 9444-210  
Application Dated: October 22, 2004

Dear Ms. Rohner:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments listed below:

1. Delete "Farms" from the front panel. This is a vague use site, and could be implied to apply to any area on a farm, i.e. farm house, farm equipment, etc.
2. Modify your marketing claims to read Kills[& Repels] [Files &] Mosquitoes, [including the Culex species] [which may carry][the West Nile Virus].
3. Revise the directions for use as follows:

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product inconsistent with it labeling.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics, do not spray at close range.

**"For Use on Animals:**

**Do not spray on sick animals. Do not spray directly towards animals's eyes. Do not treat dogs or cats.**

**Diary Cattle:** Thoroughly spray entire animal from approximately 2 feet distance. Spot treat withers, shoulders and back where saliva accumulates from head tossing. Thoroughly wash teats before milking.

**Beef Cattle:** Thoroughly spray entire animal from approximately 2 feet distance. Repeat application daily.

**Horses and Ponies:** Lightly mist over the back of horses and ponies from a distance of 2 feet. Repeat application daily.

**Hogs and Swine:** Spray directly over the back of hogs for 1-2 seconds per hog. Repeat applications daily.

**For Use in Milking Parlors, Milk Rooms, Barnes, Horse Barns, Stables, Hog Houses, and Poultry Houses to kill or repel the listed insects:**

When using this product in an enclosed area, remove animals before spray. Cover all feed, and water troughs. Close all windows and doors. Do not remain in treated area. After 15 minutes, ventilate treated area. Repeat application daily.

**Milking Parlors and Milk Rooms:** In Milking Parlors and Milk Rooms, cover all milking utensils and milk to prevent contamination from spray and dead or falling insects. Apply as a space spray upwards, spraying for 1-3 seconds per 1,000 cubic feet of space.

**Beef Cattle Barn and Hog Barns:** Spray at a rate of 1-2 seconds per 1,000 cubic feet of space.

**Stanchion Barn Use:** Walk behind animals and direct spray over backs, allowing 1 second per animal.

**Poultry houses:** Use only in empty poultry houses (when birds are not present).

Apply as a space spray upwards at a rate of 1-2 seconds per 1,000 cubic feet."

**To Kill Mud Dauber and Wasps:** Apply mist directly on insects and their nests from approximately 2 feet distance. Contacted insects will fly away from the spray. Applications should be made in late evening, when insects are at rest. Spray into hiding and breeding places contacting as many insects as possible.

page No. 3  
EPA REG. NO. 9444-210

**To kill fleas in pet kennels and catteries: Do not use this product to treat Dogs or Cats. Remove animals before spraying. Change litter and pet bedding before treatment. Close all doors and windows.** Hold aerosol 36 inches above floor and direct spray towards floor and lower walls for 10 seconds per 100 square feet. Make sure that all floor areas are treated. **Keep treated area closed for 15 minutes, then open and ventilate and allow surfaces to thoroughly dry before reoccupying.** Retreat after 7 days.


**To kill flies, gnats, mosquitoes outdoors:** Spray be walking the length of the target area at 25 foot intervals while spraying downwind. This container treats up to one acre. Repeat treatment in 7 days."

4. Modify the "OUTDOOR GROUND APPLICATION" to read: Kills [& repels] [Flies &] Mosquitoes, [including the Culex species] [which may carry] [the West Nile Virus].

In these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed with the efficacy review.

Sincerely,



Richard J. Gebken  
Product Manager 10  
Insecticide Branch  
Registration Division (7505c)

4/6



[For Reviewer: Optional – Country Vet logo]

**CB**<sup>®</sup> [For Reviewer: Optional – CB<sup>®</sup> logo]

Pyrethrins Are Made From Chrysanthemum Flowers

# Country Vet Farm and Dairy CV-80D for Insect Control

Instant Knockdown

.50% Pyrethrins for Quick Kill

- Use as a premise spray or directly on animal

**Kills [& Repels] [Flies &] Mosquitoes, [including the Culex species] [which may carry] [that may carry] [that carry] [which carry] [and transmit] [the West Nile Virus] [including those that carry West Nile Virus]**

**KILLS and REPELS: Flies, Mosquitoes, Small Flying Moths, Wasps, Hornets, Gnats, Cockroaches, Ants, Fleas, Crickets, Spiders, Asian Beetles, Barn Flies, Deer Flies, Stable Flies, Horn Flies, Horse Flies, House Flies, Face Flies, Fruit Flies, Lice, Centipedes, Millipedes, Clover Mites, Cluster Flies and Mud Daubers**

**FOR USE [in] [and] [around] Beef Cattle Operations, Dairy Farms (including Milk House, Milk Parlor, Loafing Sheds, and Holding Lot), Hog Operations, Kennels, Barns, Stables, Farms, Animal Quarters, Milkrooms, and Poultry Houses**

**CONTAINS NO CFCs OR OTHER OZONE DEPLETING SUBSTANCES. FEDERAL REGULATIONS PROHIBIT CFC PROPELLANTS IN AEROSOLS.**

**ACTIVE INGREDIENTS:**

Pyrethrins .....	0.50%
* Piperonyl Butoxide, technical .....	4.00%
<b>INERT INGREDIENTS .....</b>	<b>95.50%</b>
	<b>100.00%</b>

\* Equivalent to 3.20% (butylcarbityl) (6-propylpiperonyl) ether and 0.80% related compounds.

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated  
NOV 22 2005**

**Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 9444-210**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**  
See Back Panel for Additional Precautionary Statements.

EPA Reg. No. 9444-210

EPA Est. No. 9444-LA-1

**NET WEIGHT 24 oz. / 680 g**

5/6

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS  
CAUTION**

Harmful if absorbed through skin or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

[For Reviewer: Boxed and "bulleted" format will be used if label space permits; otherwise a paragraph format will be used.]

**FIRST AID**

**If Swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice

**If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If In Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**HOT LINE NUMBER-** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information, call (985) 878-6751, Monday through Friday, 8:00 am to 4:30 pm CST. After 4:30pm call your poison control center. For product information call (203) 597-1812, Monday-Friday, 8:00 am to 5:00 pm EST.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Apply this product only as specified on this label.

**PHYSICAL OR CHEMICAL HAZARDS**

Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

*Note: When using this product in an enclosed area, such as a milkroom, barn or stable, remove animals before spraying.*

**DAIRY FARM USE: Milking Parlor and Milkroom**—Close all windows and doors. Direct spray upwards, spraying for 1-3 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray and dead or falling insects.

**ANIMAL USE** (Do not spray sick animals): Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

**BEEF CATTLE OPERATIONS:** In barns, close all windows and doors. Spray at the rate of 1-2 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat application as necessary.

**ANIMAL USE:** For animal application, thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

**STANCHION BARN USE:** Walk behind animals and direct spray over backs, allowing 1 seconds per animal. For most effective results, apply each morning.

**HORSE BARN OR STABLES:** In horse and pony barns, close all window and doors. Spray at the rate of 1-2 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

**ANIMAL USE:** For animal application, lightly mist over back of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

**HOG OPERATIONS:** In hog houses, close all doors and windows. Spray at the rate of 1-2 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

**ANIMAL USE:** For animal application, direct spray over back of hogs and spray for 1-2 seconds per hog. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

**POULTRY OPERATIONS:** Use only in empty poultry houses (when birds are not present). In poultry houses, close all windows and doors. Spray roosts, walls, nests, and cages thoroughly. Spray at the rate of 1-2 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Treat daily or as necessary.

**MUD DAUBER AND WASP NEST:** Spray mist directly on insects and their nests from approximately 2 ft. distance. Contacted insects will fly away from fog. Applications should be made in late evening when insects are at rest. Spray into hiding and breeding places contacting as many insects as possible.

**FLEAS: General Volumetric Treatment:** Hold aerosol 36" above floor and direct spray toward floor and lower walls at a rate of 10 seconds per 100 sq. ft. making sure that all floor areas are contacted. Change litter before application. Keep area closed for 15 minutes. Remove animals before spraying. Open and ventilate before reoccupying. Repeat treatment after 7 days or as necessary.

**KENNELS AND CATTERIES:** Remove animals before spraying. Ventilate after treatment is completed.

**OUTDOOR GROUND APPLICATION:** Flies, Gnats, and Mosquitoes, [including the Culex species] [which may carry] [that may carry] [that carry] [which carry] [and transmit] [the West Nile Virus] [including those that carry West Nile Virus], in open areas near buildings and in campgrounds. Best results are obtained when wind speed is five (5) MPH or less. Apply at a rate of 60-80 seconds per acre. Spray in wide swaths across area to be treated. Allow spray drift to penetrate dense foliage. Repeat treatment as necessary.

#### STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a cool, dry area away from heat or open flame.

**DISPOSAL:** Do not Puncture or Incinerate! **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1800-CLEANUP for disposal instructions.

[For Reviewer: The Steel graphics symbol is optional if label space allows it.]

For Additional Products and Information Visit Us At [www.Country-Vet.com](http://www.Country-Vet.com) [For Reviewer: Optional text]

Manufactured by  
**WATERBURY  
COMPANIES, INC.**  
P. O. Box 1812

Waterbury, CT 06722

985-878-6751 [For Reviewer: Optional text]

Made in USA

[www.CBProProducts.com](http://www.CBProProducts.com) [For Reviewer: Optional text]



[For Reviewer: Optional – CFCs logo]

APEPA210AmendWNV.oc4



[For Reviewer: Optional – Premium Pyrethrum logo]