



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

MAY 22 2006

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Ms. Cynthia Rohner  
Waterbury Company  
P.O. Box 640, 129 Calhoun Street  
Independence LA 70443

Dear Ms. Rohner,

Subject: Country Vet Farm and Dairy CV-80D for Insect Control  
EPA file symbol 9444-210  
Submission dated 1-31-06


The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments listed below.

The flies and mosquitoes claim must be split into two separate claims so it is clear that the claim about the culex species which may carry the West Nile Virus is not associated with flies.

"Livestock farms" must be clarified to read "livestock premises (or quarters) on farms".

One copy of the finished labeling must be submitted prior to releasing the product for shipment. If you have questions, please call Ann Sibold at 703 305-6502.

Sincerely,

  
for Richard Gebken  
Product Manager 10  
Insecticide Branch  
Registration Division (7505C)

2 8 4

**ACCEPTED  
WITH COMMENTS  
In EPA Letter Dated:**

**MAY 22 2006**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.  
9444-210**



[For Reviewer: Optional - Country Vet logo]



[For Reviewer: Optional - CB logo]

Pyrethrins Are Made From Chrysanthemum Flowers

# Country Vet Farm and Dairy CV-80D for Insect Control

Instant Knockdown

.50% Pyrethrins for Quick Kill

• Use as a premise spray or directly on animal

Kills [& Repels] [Flies &] Mosquitoes, including the Culex species which may carry the West Nile Virus

**KILLS and REPELS: Flies, Mosquitoes, Small Flying Moths,  
Wasps, Hornets, Gnats, Cockroaches, Ants, Fleas, Crickets,  
Spiders, Asian Beetles, Barn Flies, Deer Flies, Stable Flies, Horn Flies, Horse Flies,  
House Flies, Face Flies, Fruit Flies, Lice, Centipedes,  
Millipedes, Clover Mites, Cluster Flies and Mud Daubers**

**FOR USE [in] [and] [around] Beef Cattle Operations, Dairy Farms (including  
Milk House, Milk Parlor, Loafing Sheds, and Holding Lot),  
Hog Operations, Kennels, Barns, Stables, Livestock Farms, Animal  
Quarters, Milkrooms, and Poultry Houses**

**CONTAINS NO CFCs OR OTHER OZONE DEPLETING SUBSTANCES.  
FEDERAL REGULATIONS PROHIBIT CFC PROPELLANTS IN AEROSOLS.**

**ACTIVE INGREDIENTS:**

Pyrethrins .....	0.50%
* Piperonyl Butoxide, technical .....	4.00%
<b>INERT INGREDIENTS .....</b>	<b>95.50%</b>
	<b>100.00%</b>

\* Equivalent to 3.20% (butylcarbityl) (6-propylpiperonyl) ether and 0.80% related compounds.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Back Panel for Additional Precautionary Statements.

EPA Reg. No. 9444-210

EPA Est. No. 9444-LA-1

**NET WEIGHT 24 oz. / 680 g**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**CAUTION**

Harmful if absorbed through skin or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

[For Reviewer: Boxed and "bulleted" format will be used if label space permits; otherwise a paragraph format will be used.]

**FIRST AID**

**If Swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If In Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**HOT LINE NUMBER-** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information, call (985) 878-6751, Monday through Friday, 8:00 am to 4:30 pm CST. After 4:30pm call your poison control center. For product information call (203) 597-1812, Monday-Friday, 8:00 am to 5:00 pm EST.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Apply this product only as specified on this label.

**PHYSICAL OR CHEMICAL HAZARDS**

Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

*Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.*

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics, do not spray at close range.

**FOR USE ON ANIMALS:** Do not spray on sick animals. Do not spray directly towards animal's eyes. Do not treat dogs or cats.

**Dairy Cattle:** Thoroughly spray entire animal from approximately 2 feet distance. Spot treat withers, shoulders and back where saliva accumulates from head tossing. Thoroughly wash teats before milking.

**Beef Cattle:** Thoroughly spray entire animal from approximately 2 feet distance. Repeat application daily.

**Horses and Ponies:** Lightly mist over the back of horses and ponies from a distance of 2 feet. Repeat application daily.

**Hogs and Swine:** Spray directly over the back of hogs for 1-2 seconds per hog. Repeat applications daily.

**For Use in Milking Parlors, Milk Rooms, Barns, Horse Barns, Stables, Hog Houses, and Poultry Houses to kill or repel the listed insects.**

**When using this product in an enclosed area, remove animals before spraying. Cover all feed, and water troughs. Close all windows and doors. Do not remain in the treated area. After 15 minutes, ventilate treated areas. Repeat application daily.**

**Milking Parlors and Milk Rooms:** In Milking Parlors and Milk Rooms, cover all milking utensils and milk to prevent contamination from spray and dead or falling insects: Apply as a space spray upwards, spraying for 1-3 seconds per 1,000 cubic feet of space.

**Beef Cattle Barn and Hog Barns:** Spray at a rate of 1-2 seconds per 1,000 cubic feet of space.

**Stanchion Barn Use:** Walk behind animals and direct spray over backs, allowing 1 second per animal.

**Poultry Houses:** Use only in empty poultry houses (when birds are not present).

Apply as a space spray upwards at a rate of 1-2 seconds per 1,000 cubic feet.

**To Kill Mud Dauber and Wasps:** Apply mist directly on insects and their nests from approximately 2 feet distance.

Contacted insects will fly away from the spray. Applications should be made in late evening, when insects are at rest.

Spray into hiding and breeding places contacting as many insects as possible.

**To kill fleas in pet kennels and catteries:** Do not use this product to treat Dogs or Cats. Remove animals before spraying. Change litter and pet bedding before treatment. Close all doors and windows. Hold aerosol 36 inches above floor and direct spray towards floor and lower walls for 10 seconds per 100 square feet. Make sure that all floor areas are treated. Keep treated area closed for 15 minutes, then open and ventilate and allow surfaces to thoroughly dry before reoccupying. Retreat after 7 days.

to kill flies, gnats, and mosquitoes outdoors. Spray while walking the length of the target area at 20-foot intervals while spraying downwind. This container treats up to one acre. Repeat treatment in 7 days.

**OUTDOOR GROUND APPLICATION:** Kills and Repels Flies, Gnats, and Mosquitoes, including the Culex species which may carry the West Nile Virus, in open areas near buildings and in campgrounds. Best results are obtained when wind speed is five (5) MPH or less. Apply at a rate of 60-80 seconds per acre. Spray in wide swaths across area to be treated. Allow spray drift to penetrate dense foliage. Repeat treatment as necessary.

#### STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a cool, dry area away from heat or open flame.

**DISPOSAL:** Do not Puncture or Incinerate! **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1800-CLEANUP for disposal instructions.

[For Reviewer: The Steel graphics symbol is optional if label space allows it.]

For Additional Products and Information Visit Us At [www.Country-Vet.com](http://www.Country-Vet.com) [For Reviewer: Optional text]

Manufactured by  
**WATERBURY  
COMPANIES, INC.**

P. O. Box 1812

Waterbury, CT 06722

985-878-6751 [For Reviewer: Optional text]

Made in USA

[www.CBProProducts.com](http://www.CBProProducts.com) [For Reviewer: Optional text]



[For Reviewer: Optional – CFCs logo]



[For Reviewer: Optional – Premium Pyrethrum logo]