

U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Progra Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460

EPA Rog. Number: 9444-225

Date of Issuance:

MAY 2 4 2004

Term of Issuance: Conditional

Name of Pesticide Product:

CB Deltamethrin Cylinder Insecticide

NOTICE OF PESTICIDE:

x Registration _ Reregistration

Waterbury Companies, Inc. 129 Calhoun Street - P.O. Box 640 Independence, Louisiana 70443

Name and Address of Registrant (include ZIP Code):

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A), provided that you:

- Submit and/or cite all data required for registration/reregistration of your product 1 under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of you product under FIFRA section 4.
- Make the following labeling changes: 2.
 - Revise the EPA Registration Number to read "EPA Reg. No. 9444-225". a)
 - Delete all reference and directions for "foam" application. No data has been submitted to **b**) support this method of application.
 - Under Perimeter Treatment delete "D-Force HPX Residual with Deltamethrin" and c) replace with the product name.
- 3. The Confidential Statement of Formula (CSF) dated 6/16/03 submitted for this product is acceptable. Please note that this formulation was previously submitted as an alternate formulation for EPA Reg. No. 9444-217 (the product cited for this me-too registration). That

CSF, submitted via Notification per PR Notice 98-10 on 9-10-02 was inadvertently accepted and is now being rescinded. The flash point/flame extension reported on the confidential statement of formula is significantly different from the basic formulation for 9444-217.

4. Please submit two (2) copies of your final printed label for before your release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing amended labeling constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

If you have any questions regarding this action, please contact Olga Odiott of my team at (703) 308-9369.

George P. LaRocca, Product Manager (13), Insecticide Branch, Registration Division (7505C) Date:

MAY 2 4 2004

EPA Form 8570-6

Enclosure

CB Deltamethrin Cylinder Insecticide

Broad Spectrum Insect Control, KILLS: Ants, Bees, Boxelder Bugs, Carpet Beetles, Centipedes, Cluster Flies, Cockroaches, Crickets, Elmleaf Beetles, Merchant Grain Beetles, Saw-toothed Grain Beetles, Fire Ants, Firebrats, Fleas, Flour Beetles, Grain Beetles, Ground Beetles, Grain Mites, Gnats, Midges, Millipedes, Mole Crickets, Moths, Pillbugs, Scorpions, Silverfish, Sowbugs, Spiders, Springtails, Ticks, Wasps and Many Other Crawling Insects.

Also Controls and Kills: Wood Infesting Pests - Termites (Subterranean, Formosan & Drywood), Carpenter Ants and Carpenter Bees, Carpet Beetles and Wasps.

FOR USE IN Food Handling Establishments, Including Food Service, Manufacturing and Processing Establishments, Such as Restaurants, Cafeterias, Supermarkets, Bakeries, Breweries, Dairies, Meat, Poultry & Egg Processing & Packaging Plants, Canneries, Feed Handling Establishments Including Feed Manufacturing and Processing Establishments. Also For Use In Homes, Tents, Aircraft (in cargo areas only), Apartments, Atriums, Attics, Automobiles, Basements, Bathrooms, Boats, Bottling facilities, Buses, Cabins, Campers, Candy Plants, Carports, Cat Sleeping Quarters, Clothes Storage, Condominiums, Decks, Dens, Driveways, Garages, Gazebos, Granaries & Grain Mills, Hospitals, Hotels, Industrial Buildings, Residential, Institutional, Kitchens, Laboratories, Lanais, Living Rooms, Mausoleums, Offices, Parlors, Patios, Pet Sleeping Areas, Play Rooms, Porches, Railcars, Recreational Vehicles, Rooms, Schools, Ships & Vessels, Solariums, Storage Areas, Sun Rooms, Trailers, Trucks, Utility Rooms, Verandahs, Warehouses and Wineries.

·Protects for up to 8 Weeks

·Killing Action and Residual Control

·Keeps on Killing for Up to 8 Weeks

·For Indoor/Outdoor Use

·15% VOC Formulation

·Above-Ground Spot Treatment Termite Control

·Dry/Solvent Based Formulation

·For Residential, Industrial and Institutional Control of Major Nuisance Pests Listed on the label

·Controls Numerous Pests in and around Households and Other Structures

ACTIVE INGREDIENT:

Deltamethrin..... INERT INGREDIENTS...... 99.94% 100.00%

Contains Petroleum Distillates

EPA Reg. No. 9444-XXX

EPA Est. No. 9444-LA-1

KEEP OUT OF REACH OF CHILDREN

U.S. Patents Pending

NET WEIGHT 13 lbs./5.897 kg.

ACCEPTED with COMMENTS CAUTION In EPA Letter Dated
See Back Panel For Additional Precautionary Statements.

Under the Federal Insecticide. Fungicide, and Rodenticide Act. as amended, for the pesticide registered under EPA Reg. No.

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID	
If Inhaled	Move person to fresh air.
	• If person is not breathing, call 911 or an ambulance, then give artificial
	respiration, preferably by mouth-to-mouth, if possible.
	Call a poison control center or doctor for further treatment advice.
If on Skin or	Take off contaminated clothing.
Clothing	• Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
If In Eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	Remove contact lenses, if present, after the first 5 minutes, then continue
	rinsing eye.
	Call a poison control center or doctor for treatment advice.
If Swallowed	Call a poison control center or doctor for treatment advice.
	• Do not give any liquid to the person.
	• Do not induce vomiting unless told to do so by the poison control center or
	doctor.
	Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or	
going for treatment. For emergency information, call (985) 878-6751, Monday through Friday,	
8:00 am to 4:30 pm CST. After 4:30pm call your poison control center. For product information	
call (203) 597-1812, Monday-Friday, 8:00 am to 5:00 pm EST.	
NOTE TO PHYSICIAN	
Contains petroleum distillate - vomiting may cause aspiration pneumonia.	

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to any body of water.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product has residual action which keeps on killing for up to 8 weeks after you spray. Do not allow people or pets on treated surfaces until dry. Do not apply to classrooms when in use. Do not apply this pesticide in occupied patient rooms, or in any rooms occupied by the infirm, elderly or children for an extended period of time.

Remove pets, birds and cover fish aquariums before spraying.

Use appropriate equipment, such as crack & crevice extension tips for voids and wood injectors for wood injection treatments. Place injection tips into cracks, crevices, holes and small openings. Release approximately one-second of product. Several holes may be required in long-running voids.

INDOOR AND OUTDOOR TREATMENTS

WOOD INFESTING INSECTS- Termites (Subterranean, Formosan and Drywood)- For control of above-ground termites in localized areas, apply to voids and galleries in damaged wood, and in spaces between wooden structural members and between the sill plate and foundation where wood is vulnerable. Applications may be made to inaccessible areas by drilling, and then injecting with the supplied void injector into the damaged wood or wall voids. These applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplement for controlling active infestations of subterranean termites. Termite carton nests in building voids may be injected using a void injector. It is desirable to physically remove carton nest material from building voids when such nests are found. This type of application is intended to be a supplemental treatment for control of drywood and above-ground subterranean termites and other wood-infesting insects.

WOOD DESTROYING INSECTS-Localized treatment of wood-infesting insects: Carpet Beetles, Carpenter Ants, Carpenter Bees, Termites and Wasps: Locate suspected infestation and/or insect tunnels and cavities. Drill pilot holes into insect tunnels, galleries and inaccessible closed voids. Drill holes 8 to 10 inches apart to insure thorough coverage. Insect 5 to 10 seconds per hole. Do not contact electrical wiring, plumbing, etc. when drilling and never use metal wood injectors where electrical shock could occur. Use a standard plastic void injector if there is danger of electrical contact. For carpenter bees, seal bee entrances after application. This treatment is intended for localized termite infestations only.

For active and hibernating stages of Boxelder Bugs, Elmleaf Beetles and Cluster Flies: Insect cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices, behind siding, wall voids, around window and door frames. Application to attics, crawl spaces, unfinished basements, or manmade voids may be sprayed to control exposed worker and winged reproductive forms of termites. For control of swarming or reproductive stages of termites, spray to the swarm or area on which they collect.

The purpose of such applications of this product is to kill workers or winged reproductive forms which may be present at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but are merely a supplement. This product is not recommended as sole protection against termites. For active indoor termite infestations, get a professional inspection.

Carpenter Ants: For effective control in houses and other structures, apply as general surface, spot, crack and crevice or wall void application. Apply around doors and windows and other places where ants may enter the premises and where they crawl and hide. Spray into cracks and crevices or through small drilled holes into voids where these ants or their nests are present. For control of carpenter bees in structures, spray the liquid directly into gallery entrance holes. Following treatment, the entrance holes may be left open 24 hours to be certain that returning adult bees are killed. When there is no activity, the hole may be sealed with a suitable sealant.

INDOOR: Cockroaches (including adult and immature stages of both non-resistant and organophosphate and carbamate resistant strains), ants, booklice, boxelder bugs, centipedes, crickets, dermestids, firebrats, fleas palmetto bugs, silverfish, sowbugs, ticks and waterbugs: Apply thoroughly to all areas where these pests crawl and hide, especially in cracks and crevices, and hidden areas around sinks and storage areas, behind baseboards, around doors and windows, behind and under refrigerators, cabinets, sinks and stoves, the underside of shelves, drawers, bookcases, and similar areas. When pests are driven out, spray directly.

[COCKROACHES: Spray in cracks and crevices, around baseboards and other hiding places until wet. When driven out, spray bugs directly.]

Ants: Spray freely around ant trails, doors and windows and other places where ants enter the house. Ticks and Fleas on Surfaces: Remove pet bedding and destroy or clean thoroughly. Spray pet resting quarters. Adult fleas and larvae contacted by spray will be killed. Put fresh bedding down once spray has dried. Do not spray animals directly. For best results, pets should be treated with an appropriate

product before allowing them to return to the treated area. Spray rugs and carpets where infestations are bad. Test first for staining on an inconspicuous piece prior to use.

Spiders and Scorpions: Apply along and behind baseboards, to window and door frames, corners, pipes, storage localities, attics, crawl spaces and other areas over which these pests may crawl.

Centipedes, Ground Beetles, Pillbugs, Sowbugs and Ticks: Apply around doors and windows and other places where these pests may enter premises. Treat baseboards, storage areas and other locations where these pests are found.

[SILVERFISH, CRICKETS, SPIDERS: Apply to areas infested by these pests or spray pests directly.]

PANTRY PESTS – Grain Beetles (Rusty, Merchant and Saw-toothed), Flour Beetles (Red and Confused), Chocolate Moths, Cigarette Beetles, Clover Mites, Cluster Flies, Drugstore Beetles, Elmleaf Beetles, Rice Weevils, Lesser Grain Borers, Spider Beetles, Grain Mites and Tobacco Moths: Treat exposed stages. Remove all food-stuffs, utensils and shelf paper from area to be treated. Discard used shelf paper and any infested foodstuffs. Apply spray to shelves and in cracks and crevices behind and under cupboards and cabinets. Allow spray to dry prior to replacement of shelf paper, food and utensils. [For Foam Application]:PANTRY PESTS – Grain Beetles (Rusty, Merchant and Saw-toothed), Flour Beetles (Red and Confused), Chocolate Moths, Cigarette Beetles, Clover Mites, Cluster Flies, Drugstore Beetles, Elmleaf Beetles, Rice Weevils, Lesser Grain Borers, Spider Beetles and Tobacco Moths: Apply in cracks and crevices behind and under cupboards and cabinets.

Carpet Beetles (Black, Furniture and Varied): Spray on and under edges of floor coverings, under rugs and furniture and in closets or other localities where these insects are found. [For Foam Application]: Carpet Beetles (Black, Furniture and Varied): Apply in cracks and crevices in closets or other localities where these insects are found.

Webbing Clothes Moths: Use this product as a crack and crevice treatment in closets and other storage areas where these pests are found. This is an adjunct treatment and will not control moth larvae already on clothes. Do not apply to clothing.

Food/Feed Handling Establishments: For control of all labeled pests, applications are permitted in food/feed and non-food/non-feed areas of food/feed handling and processing establishments as a general surface, spot or crack and crevice treatment. I second spray per spot. This product is authorized for use in Federally Inspected Meat and Poultry Plants as a crack and crevice treatment. In the home, all food processing surfaces should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed.

Food/Feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food/feed. Serving areas where food/feed is exposed and the facility is in operation are also considered food/feed areas.

Non-Food/Non-Feed Areas: Examples of non-food/non-feed areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets and storage (after packaging, canning or bottling). All areas that insects inhabit or through which insects may enter should be treated.

General Surface Applications: Do not use this application method in the food/feed handling areas of the facility when the facility is in operation or foods/feeds are exposed. Do not apply directly to food/feed products. Cover or remove all food/feed processing and/or handling equipment during application. After application in food/feed processing plants, cafeterias and similar facilities, wash all equipment, benches, shelving and other surfaces which food/feed will contact. Clean food/feed handling or processing equipment and thoroughly rinse with clean, fresh water. After application in bakeries, clean all food contact surfaces thoroughly with appropriate equipment.

Spot or Crack & Crevice Applications: Except in federally inspected meat and poultry plants, spot or crack and crevice applications may be made while the facility is in operation provided exposed food is covered or removed from area being treated prior to application. Do not apply directly to food or

food-handling surfaces. [For Foam Application]: Spot or Crack & Crevice Applications: Except in federally inspected meat and poultry plants, spot or crack and crevice applications may be made while the facility is in operation. Do not apply directly to food or food-handling surfaces.

In the home, do not allow spray to contact food/feed surfaces. If it does, clean surfaces with soap and water.

OUTDOOR TREATMENTS

OUTDOORS: Remove all exposed food and cooking utensils. Cover all food handling surfaces or wash thoroughly after treatment and before use. Do not use on edible crops. Point nozzle away from face. Hold can on a slight downward angle, approximately 12 - 15 inches from surface to be sprayed. For best results, spray when air is calm. Spray with wind if breeze is blowing.

SUBTERRANEAN ANTS SUCH AS FIRE ANTS: To treat nests, use a soil injector. To maximize insecticide distribution, insert injector into the nest full length if possible. Inject the nest at 3-6 sites at a rate of 5-10 seconds per site. For very large nests (over 36" diameter), increase the number of injection sites. Space injection sites in a circular pattern on the nest surface with one site in the center. To quiet active ants on nest surface, sweep surface with spray from a distance of 12 inches. For best results, apply when temperature is 65-85°F, or in early morning or late evening hours. Treat new nests as they appear.

PESTS ON OUTSIDE SURFACES OF BUILDINGS: For control of building infestation to ants, clover mites, crickets and sowbugs (pillbugs). Spray foundation of building where insects are active and may find entrance up to a height of 2 to 3 feet. Apply as a residual spray to outside surfaces of buildings where insects tend to congregate. Spray areas include areas such as screens, window frames, eaves, porches, patios, garages and refuse dumps.

CRAWLING INSECTS: Ants, Cockroaches, (including Asian Cockroaches), Centipedes, Crickets, Mole Crickets and Silverfish. Spray infested surface of patio or picnic area, hitting as many insects as possible. Also spray legs of tables and chairs to repel ants.

PERIMETER TREATMENT: For crawling insects, apply D-Force HPX Residual with Deltamethrin to cracks and crevices where insects may enter premises or harbor. Areas to be treated include, but are not limited to, weep holds, under siding, wall voids, soffits, around attic vents, cracks and crevices openings around windows, doors and pipes. Treatment of infested fences and tree holes is permitted.

Ants: Apply to ant trails, around doors and windows and other places where ants maybe found. Lone Star Ticks, Dog Ticks, Crickets, Fleas and Earwigs: Apply thoroughly to infestation in bushes, grass or weeds.

Wasps, Mud Daubers, Hornets and Bees: Spray nests and bees directly.

FLYING INSECTS: Houseflies, Gnats, Mosquitoes and Small Flying Moths: Use only on outdoor surfaces. Spray outside surfaces of window and door frames and other areas where these pests may enter the home. Also spray localized resting areas, such as under eaves, porches, light fixtures and cords, railings, etc. where these insects may rest. Insects coming to rest on treated surfaces will be killed. [For Foam Application]: Construction practices, soil subsidence and other factors may create situations in which a continuous chemically treated barrier cannot be provided when using conventional liquid applications alone. When necessary, conventional treatments may be supplemented through the use of foam to provide a continuous treated barrier. Applications may be made behind veneers (concrete or wood), piers, chimney bases, into rubble foundations, into block voids, or structural voids, wall voids, under slabs, stoops, porches, or to the soil in crawl spaces and other similar voids

TERMITES AND CARPENTER ANTS LIVING INSIDE TREES, UTILITY POLES AND

FENCING: For control of above-ground termites inside trees, utility poles, fencing and decking materials and similar structural members, treat open holes or drill horizontal holes in a circular pattern where insects are suspected to find the interior infested cavity and spray 2 to 3 seconds of product per hole using void injector.

AERIAL TERMITE CARTON IN TREES: Termite carton nests in trees may be spray injected with product using the supplied void injector. Drill or punch hole into center of carton. Inject for 5 to 10

seconds. If carton is greater than 12 inches across, open another injection site. Inject for 5 to 10 seconds. Make one additional injection site for each 12 inches of carton. Multiple injection points to varying depths may be necessary. It is desirable to physically remove carton nest material from trees when such nests are found.

WOODBORERS IN FRUIT AND ORNAMENTAL TREES: Treat fruit and nut bearing trees when dormant. Locate borers entrance, remove accumulation of plant sap. Inject 1 to 2 seconds into gallery using standard injection equipment. Apply only to fruit trees that will not bear fruit within 12 months after application.

TREATMENT FOR INGROUND SERVICE BOXES: Ants, Centipedes, Cockroaches, Crickets, Fire Ants, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders and Springtails. Use soil injector or spray actuator to treat open boxes and exposed surfaces. Do not contact any electrical wiring or mechanical parts of meter.

STORAGE AND DISPOSAL

Storage: Do not store near heat or open flame. Store in a cool, dry area away from children.

Disposal: Do not Puncture or Incinerate! If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1800-CLEANUP for disposal instructions.

[For Reviewer: The Steel graphics symbol is optional if label space allows it.]

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Manufactured by
WATERBURY
COMPANIES, INC.
P. O. Box 1812
Waterbury, CT 06722
Made in USA
985-878-6751[For Reviewer: Optional text]
www.CBProProducts.com [For Reviewer: Optional text]

9444-EEL