

**For General  
Dairy Equipment Cleaning**

**Milk, Cheese, Butter Plant Sanitation.** One ounce to 4 gallons of water will give a 200 ppm solution of available chlorine. After cleaning and just prior to use, all equipment surfaces should be exposed to the germicide for not less than 2 minutes. Do not rinse equipment with water after sanitizing.

**BOTTLE CHLORINATION**

**Just prior to filling operation, sanitize bottles with a 200 ppm solution of available chlorine.**

**LOCAL AND STATE REGULATIONS**

**Where regulations are in effect consult authorities for additional information.**

**CONTAINER DISPOSAL**

**Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place. Deposit containers return to vendor.**

**Packaged for Hawkins Chemical Co., Inc.  
Minneapolis, Minn.  
By Vel-Tex Chemical Co., Inc.  
St. Paul, Minn.**

**TABLE OF PROPORTIONS**

**800 ppm —  
Use 1 oz. in 1 gal. water**

**200 ppm —  
Use 1 oz. in 4 gal. water**

**1 ppm —  
Use 1 oz. in 1000 gal. water**

**STORAGE**

**Degrades with age. Use a test kit and increase dosage as necessary to obtain required level of available chlorine.**

**FISH AND WILDLIFE CAUTIONS**

**This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.**