

[9594-13]

USE PRECAUTIONS

The use of ICC Concentrated Spray in food processing plants should be confined to time periods when the plant is not in operation. Foods should be removed or covered before treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Remove any birds, pets or fish aquarium before spraying.

When diluting ICC Concentrated Spray with water, prepare only the amount needed for immediate use. Holding a water dilution more than 24 hours is not recommended. Aerate occasionally. For oil dilutions, a deodorized petroleum oil is recommended.

Vacate premises after treatment and ventilate before reoccupying. Sweep up and remove dead insects.

Repeat treatment with ICC Concentrated Spray when reinfestation occurs.

DIRECTIONS

For maximum effectiveness, a combination of localized surface application and space treatment is recommended.


AS A SURFACE SPRAY: For initial clean up of severe insect infestations, dilute ICC Concentrated Spray at a rate of 1 part concentrate in 4 parts water or oil. This dilution may not be used in edible product areas of federally inspected meat and poultry plants in those plants and for normal infestation, dilute 1 part ICC Concentrated Spray in 9 parts water or oil.

To control **roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, meal worms, grain mites, and cadellies:** Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks, and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray milkcases. For ants, spray trails, nests and points of entry. If surface application only is desired, spray floors, walls and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control **carpet beetles,** spray base of carpet and under carpeting and rugs. Make localized application to baseboards. Spray directly into cracks, seams and other areas of shelving. Repeat treatment as needed.

To control **bedbugs,** spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in places. Treat baseboards, mouldings and floors. Repeat treatment as needed.

Batch # _____



CONCENTRATED SPRAY EMULSIFIABLE CONCENTRATE

EPA Reg. No. 9594-13

A FOOD INDUSTRY INSECT SPRAY

May be diluted with either water or oil

Approved for Use in All Departments of Official Establishments Operating Under the Meat, Poultry, Shell Egg Grading and Equine Products Inspection Programs

Provides Rapid Kill of Insects Named Without Objectionable Residues.

The Insecticidal Components (Piperonyl Butoxide and Pyrethrins) are Classified as Food Additives and When This Product Is Used as Directed, Acceptable Tolerances Levels will not be Exceeded.

ACTIVE INGREDIENTS

Pyrethrins	1%
Piperonyl Butoxide (Technical)	1%
Petroleum Distillate	1%
INERT INGREDIENTS	98%

ICC Concentrated Spray is a registered trademark of Intercontinental Chemical Corporation.

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

Net Contents Gallon(s)

MANUFACTURED FOR

Intercontinental Chemical Corporation

Manufacturing Plants: CINCINNATI, OHIO, SACRAMENTO, CALIF., ANAHEIM, CALIF.

DIRECTIONS FOR USE (CONT)

To control **fleas** and **brown dog ticks**, thoroughly spray the areas of both resting quarters, nearby racks and heavy curtains and behind baseboards, mouldings, window and door frames and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. For current treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

AS A SPACE SPRAY: Dilute at a rate of 1 part ICC Concentrated Spray to 19 parts water or oil. Apply with mechanical or compressed air equipment (non-thermo) adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems. For rapid control of **flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths:** direct spray at an upward angle distributing it uniformly through the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Ventilate and sweep up dead insects before reoccupying.

AS A FOG SPRAY: For rapid kill of exposed and inaccessible stages of insects named on this label, dilute 1 part ICC Concentrated Spray to 19 parts water or oil. Apply using compressed mechanical or compressed air equipment (non-thermo) following directions for space spray. Apply the above dilution at a rate of 1 to 2 ounces per 1000 cubic feet of space.

When used in meat and poultry processing plants, a dilution of more than 1 to 1,500 is acceptable and it must be dispensed from equipment adjusted to deliver a aerosol spray, no greater than 50 microns in diameter and 80 f.p.s. less than 35 m.p.h.

Always use proper ventilation.

This product is toxic to fish, birds, cats, dogs, and other animals. Do not use in areas where such animals are present. Do not use on birds.

Apply this product in accordance with the following directions:

For control of insects named on this label, use the following directions: For control of insects named on this label, use the following directions:

TERMINAL TREATMENT: For control of insects named on this label, use the following directions:

TERMINAL TREATMENT: For control of insects named on this label, use the following directions:

TERMINAL TREATMENT: For control of insects named on this label, use the following directions:

TERMINAL TREATMENT: For control of insects named on this label, use the following directions:

TERMINAL TREATMENT: For control of insects named on this label, use the following directions:

TERMINAL TREATMENT: For control of insects named on this label, use the following directions:

TERMINAL TREATMENT: For control of insects named on this label, use the following directions:

TERMINAL TREATMENT: For control of insects named on this label, use the following directions:

TERMINAL TREATMENT: For control of insects named on this label, use the following directions:

TERMINAL TREATMENT: For control of insects named on this label, use the following directions:

TERMINAL TREATMENT: For control of insects named on this label, use the following directions:

TERMINAL TREATMENT: For control of insects named on this label, use the following directions:

TERMINAL TREATMENT: For control of insects named on this label, use the following directions:

EPA EST. #9594-13