

New and Improved

Heavy Duty Weed Killer with 2, 4-D

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid **49.6%**

50.4°

CONTROLS THESE WEEDS

Redroot Fl. green
 Kochia
 Cocklebur
 Lambquarters
 E. of P. grasses
 F. of T. grasses

Can Be used On The Following Areas

Other recreation areas	Ditch banks
Air fields	Storage sheds
Water storage	Water towers



DEL Chemical Corporation

MENOMONEE FALLS, WISCONSIN

RENO, NEVADA

6. 11. 1954
1954
6. 11. 1954
- 1954
- 1954

GRA
the
for
and
+ 10
10
10
10
10
10

[illegible]

WHE
rche
24-
QAT

N

4-D

EDS

Gumweed
Burdock
Pissenerstock
Wild Radish
Redroot Pigweed
Kochia
Crackbur
Lambquarters
Barns or Peggards
Pitted Thistle

ng Areas

as Ditch banks
Pasture lands
Rain of height of 2 ft

15

ation

NEVADA

DIRECTIONS

SELECTIVE WEEDING IN CROPS

For use in corn, sorghum, and other crops. Apply 2 to 4 pints per acre in the early boot stage of the crop. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds.

GRASSES In corn, sorghum, and other crops. Apply 2 to 4 pints per acre in the early boot stage of the crop. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds.

In corn, sorghum, and other crops. Apply 2 to 4 pints per acre in the early boot stage of the crop. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds.

In corn, sorghum, and other crops. Apply 2 to 4 pints per acre in the early boot stage of the crop. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds.

CAUTION: Keep Out of Reach of Children

Avoid contact with skin, eyes, nose, mouth, and clothing. If contact occurs, wash immediately with plenty of water. If swallowed, drink plenty of water. Do not eat, drink, or smoke while using this product. Do not use near open flames or heat. Do not use in confined spaces. Do not use in areas where children or livestock are present. Do not use in areas where food is stored. Do not use in areas where food is prepared. Do not use in areas where food is served. Do not use in areas where food is consumed. Do not use in areas where food is stored. Do not use in areas where food is prepared. Do not use in areas where food is served. Do not use in areas where food is consumed.

Do NOT Store with Foodstuffs

WHEAT AND BARLEY: In spring, apply 2 to 4 pints per acre in the early boot stage of the crop. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds.

CORN

For use in corn, sorghum, and other crops. Apply 2 to 4 pints per acre in the early boot stage of the crop. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds.

WITH LIQUID NITROGEN SOLUTIONS

In corn, sorghum, and other crops. Apply 2 to 4 pints per acre in the early boot stage of the crop. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds.

SUGAR CANE

In corn, sorghum, and other crops. Apply 2 to 4 pints per acre in the early boot stage of the crop. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds.

SEED RELEASE

In corn, sorghum, and other crops. Apply 2 to 4 pints per acre in the early boot stage of the crop. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds.

NON SELECTIVE WEED CONTROL AND PREVENTION OF SEED FORMATION

In corn, sorghum, and other crops. Apply 2 to 4 pints per acre in the early boot stage of the crop. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds.

In corn, sorghum, and other crops. Apply 2 to 4 pints per acre in the early boot stage of the crop. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds.

In corn, sorghum, and other crops. Apply 2 to 4 pints per acre in the early boot stage of the crop. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds. Do not apply to crops with sensitive roots. Do not apply to crops with sensitive leaves. Do not apply to crops with sensitive stems. Do not apply to crops with sensitive seeds.

DEL CHEMICAL CORPORATION

MENOMONEE FALLS, WISCONSIN

RENO, NEVADA