


DIRECTIONS FOR USE

1. Mix 1 to 2 oz. of Okay Rat Kill in shallow container. Okay Rat Kill bait should be placed in bait stations near to walls, runways, or where rats are suspected to feed. Bait stations should be placed in corners, and in concealed locations where rats and mice are fed. Bait stations are desirable. Baiting should continue until all signs of feeding have stopped, usually 3 to 4 days. Bait stations should remain empty. Because of Okay Rat Kill in shallow container protects the bait from being contaminated and allows the bait stations to be moved from areas where feeding has stopped to areas where rodent activity is still noted. Bait stations should be placed in locations not accessible to children, pets and livestock, or in a corner of a room.



OKAY
The Best Yet Brand

RAT KILL

Contains Prolin* a better more effective Warfarin rodenticide. *Trademark of the Wisconsin Alumni Research Foundation.

ACTIVE INGREDIENTS:

N¹-(2-quinoxaline) sulfanilamide (sulfaguinoxaline) ----- 0.025%
2-(alpha-acetonyl-4-hydroxycoumarin (warfarin.) -- 0.025%

INERT INGREDIENTS ----- 99.950%

Total ----- 100.000%

Prolin (warfarin plus sulfaguinoxaline) is manufactured under license from the Wisconsin Alumni Research Foundation, U.S. Patent No. 2,427,876, other patent applications pending. The antibacterial agent, sulfaguinoxaline, increases the effectiveness of the bait and assures more thorough control of rats. Especially recommended for rat infestations which are difficult to control.


CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS, WHEN USING - WHEN CLEANING.

FOR ADDITIONAL DIRECTIONS, SEE SIDE PANEL.

LEUNG CHEUNG COMPANY
1407 Hickey St., Omaha, Nebraska

DIRECTIONS FOR USE FOR MICE

Because mice are nervous feeders and follow no particular feeding patterns, it is important that smaller portions of bait (in teaspoon amounts - 1/2 to 1 ounce) be placed at intervals of 3 to 12 feet. Place teaspoon amounts of Okay Rat Kill in shallow containers and set under stoves, refrigerators, and lower parts of the kitchen cabinets and sheltered areas along walls. Bait should be placed in locations not accessible to children, pets and livestock, or in a corner of a room.



CAUTIONS:

Warfarin is a poison. If a rat is seen eating bait in a glass of warm water and repeat until vomit fluid is clear. Call a veterinarian immediately!

Keep away from humans, domestic animals and pets. This material may reduce the clotting ability of the blood and cause bleeding. Usual doses, intravenous and oral administration of Vitamin K combine with blood coagulants are indicated in the case of bleeding caused by overdoses of 2-hydroxycoumarin.

ACCEPTED
JUL 19 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON, REGISTERED
ED UNDER NO. 1521-7
TO ATTACHED COMMENTS

7-19-67
9554-7