


FLOORS

Shut off pump for 24 hours. One pound per pool.

potency, pools. Disperses lightly throughout the pool. Distributes evenly.

NET WT.
2 5 LBS.

Pool Brite



ALGAE CURE

ACTIVE INGREDIENTS --- Trichloro-s-triazinetriene 100%
INERT INGREDIENTS None

WARNING KEEP OUT OF REACH OF CHILDREN.
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

U.S.D.A. R

WA

Keep out of reach of children. Strong oxidizing material. May stain clothing. In case of contact with skin, wash with cold water for at least 15 minutes. Attention. Avoid breathing vapors. May cause fire. Moisture may cause hazardous chlorine gas from open flame. Keep container closed when not in use. Discard or reuse.

POISON


External: Immediately flush with water.
Internal: Feed bread soaked in milk or oil or sooking oil.
CALL A PHYSICIAN

Manu

Orex

560 Ferry St.

Pool Brite



ALGAE CURE

ACTIVE INGREDIENTS — Trichloro-s-Triazinetrione..... 100%
INERT INGREDIENTS None

U.S.D.A. R

WA

Keep out of reach of children.
Strong oxidizing material.
Keep away from clothing.
In case of contact with skin, wash with cold water for at least 15 minutes.
Attention. Avoid breathing vapors.
May cause fire. Moisture and heat may cause hazardous chlorine gas.
Keep away from open flame. Keep container tightly closed when not in use.
Do not reuse or discard in household waste.

POISON

External: Immediately flush with water.

Internal: Feed bread soaked in milk or oil or sucking on lemon juice.

CALL A PHYSICIAN FOR TREATMENT.

Manu

Ortex

560 Ferry St.

U.S.D.A. Reg. No. 9556-5

WARNING

Keep out of reach of children. May be fatal if swallowed. Strong oxidizing material. Avoid contact with skin, eyes or clothing. In case of contact, immediately flush skin or eyes with cold water for at least 15 minutes. For eyes get medical attention. Avoid breathing dust. Contact with organic matter may cause fire. Moisture contamination may liberate irritating and hazardous chlorine-containing gases. Keep away from open flame. Keep container closed and in a cool, dry place when not in use. Rinse container thoroughly when discarding or if for re-use.

POISON — ANTIDOTE

External: Immediately flush with plenty of cold water.

Internal: Feed bread soaked in milk followed by olive oil or soaking coil.

CALL A PHYSICIAN.

Manufactured for

Ortex Products Inc.

560 Ferry St., Newark, N. J. 07105

KEEP YOUR POOL CLEAR & SANITARY

DIRECTIONS

1. Adjust pool water: pH to 7.2 to 7.6 as necessary with pH Increaser or Pool Acid.
2. Proper conditioning of pool water: with POOL CONDITIONER & STABILIZER is necessary to ensure maximum effectiveness of pool chlorine. To freshly filled or previously unconditioned and stabilized pool water add 1½ lbs. of CONDITIONER & STABILIZER for each 5000 gallons of pool water capacity.
3. To a pH adjusted and preconditioned pool as recommended in steps 1 & 2, add 2 oz. Granular Chlorine for every 4000 gals. of fresh water to super chlorinate and satisfy the chlorine demand of the fresh water. Test water after 1 to 4 hours with a chlorine test kit. If there is no chlorine residual recheck pH with test kit and readjust as in step 1. Repeat above chlorine dosage until a reading of 0.8 ppm or more residual chlorine is found with test kit. Thereafter periodically add 2 oz. Granular Chlorine for each 8000 gallons of water daily or as needed to maintain 0.8 to 1.5 ppm residual chlorine at all times. Add Granular Chlorine by broadcasting at the deep end of the pool by the main drain or through the skimmer. It will dissolve quickly, leaving no residue, contains no scale forming minerals. Test water at least daily for pH as well as chlorine residual as pH has a direct effect on chlorine action. 2 oz. of Granular Chlorine added to 8000 gallons of pool water will provide 1.1 ppm available chlorine.