

KELLEY'S RAI BAIT

812 W. Stubb • Ph. DU 3-1414 • Edinburg, Texas

• ANTICOAGULANT RAT AND MOUSE BAIT •

CAUTION KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS:

#2-(Alpha-acetonylfurfuryl) - 4-hydroxy-

coumarin _____ 0.025%

INERT INGREDIENTS _____ 99.975%

TOTAL _____ 100.000%

If swallowed by humans, domestic animals or pets, this material may reduce the clotting of the blood and cause bleeding. In such cases, intravenous and oral administration of Vitamin K combined with blood transfusions may be indicated, as in the case of hemorrhage caused by overdose of bishydroxycoumarin (Dicumarol). Protect baits from children, pets and livestock by bait boxes or other suitable means.

Net. Wt. _____ Lbs.



U S D A Reg. No. 9561-1

Directions

- a) place bait in locations where rats and mice feed, water or travel. Along walls in corners and in concealed locations where rats and mice can feed unobserved. Bait should be placed in locations not accessible to children and pets in tamperproof bait boxes.

FOR RAT CONTROL

Four ounces of bait per station should be the minimum amount used.
an uninterrupted supply of bait should be maintained for at least 10 days.

FOR MOUSE CONTROL

Tablespoon amounts 1/4 to 1/2 ounces of bait should be placed at 8 - 12 foot intervals.
an uninterrupted supply of bait should be maintained for at least 15 days.

