

1801

R.B. #2

KILLS RATS AND MICE

Kills Norway Rat, Roof Rat, and House Mice

Long term control may be attained through continuous baiting.

ACTIVE INGREDIENT:

Warfarin: [3-(alpha-Acetylbenzyl)-4-hydroxycoumarin].....

0.025%

INERT INGREDIENTS:.....

99.975%

NET WT.

Place Packs - ounces each

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel for Additional Precautionary Statements

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION: May cause eye irritation. Wash thoroughly with soap and water after handling. Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K, intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD: Keep out of lakes, streams or ponds.

This product is accepted as outlined by the U.S.D.A. for use in official establishments operating under the Federal meat, poultry, shell egg grading, and egg products inspection program.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restrictions: For control of Norway rats, Roof rats, and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamper-proof bait boxes or locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Selection of Treatment Areas: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

Application Directions: Rats: Apply one to five place packs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

Mice: Open pack and apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8-12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.


Rats and Mice: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL Storage: Store only in original container, in a dry place inaccessible to children and pets. Disposal: Do not reuse empty container. Securely wrap in newspaper and discard in trash.

Zoecon Corporation
A Sandoz Company
12200 Denton Drive, Dallas, Texas 75234

EPA Reg. No. 9562-7
0689-C:0128E

EPA Est. No. 39578-TX-1



BEST AVAILABLE COPY