

February 20, 1965

REGISTERED TRADEMARK
MARCH 1965
U.S. PATENT OFFICE
OFFICE OF TRADEMARKS AND PATENT APPEALS
WASHINGTON, D.C. 20530
ATTACHED FORM NO. 1

READ THE FRONT PANEL

TRADEMARK

NATIONWIDE
EXTERMINATING
USDA REG. NO. 659121
LIQUID DEATH
Water Soluble Warfarin
A concentrate of Sodium Salt and Warfarin
for control of Rats & Mice.
Active Ingredients:
Sodium Salt of Warfarin 7.74
Sodium Salt of 3-(alpha-
acetylbenzyl)-4-hydroxy-
coumarin
Inert Ingredients: (Silica) 90.26

CAUTION: Keep out of reach of Children,
domestic animals and pets. See side
panel for additional cautions.

If swallowed by humans, domestic animals or
pets this material may reduce the clotting
ability of the blood and cause bleeding. In
such case, intravenous and oral administration
of Vitamin K combined with blood transfusions
are indicated as in the case of hemorrhage
caused by overdose of dihydrocoumarin.

HOW TO USE

EXTERMINATING RATS & MICE

FOR RATS & MICE: Mix 4 ounces Liquid Death with 3 gallons of water,
drinking water in one quart mason jar or 10 used. No mixing or stirring
is required. Do not attempt to dissolve small white particles which remain
on the bottom of jar. Liquid Death should be set out in mason jars, whiskey
bottle or in shallow pans, with no less than 1/2 inch in each location.
Place where rats and mice are accustomed to feed. Do not place where freezing
may occur. Discard at once water in which hole or filthy is used. Protect
from children or pets and domestic animals.
It is recommended that dry bait be used in conjunction with liquid bait,
especially where there is an ample supply of fresh water.

When continuous source of infestation prevails from nearby dumps or fields,
permanent bait stations of dry bait should be used and baits replenished
as needed.

(16) (16) (continued)

FOR CONTROLLING MICE: Follow the same general directions as for rats except bait placements may be smaller and more placements should be made, space at 8-10 foot intervals. Mice are more difficult to control than rats and may require a longer baiting period.

BAITING SHOULD continue until no further sign of eating dry bait or consumption of Liquid Death solution is noted. For mice this will be a period of not less than 15 days; for rats not less than 12 days.

Placement of baits: Place along walls, in corners, and in concealed locations where rats and mice can feed undisturbed. Baits should be placed in locations not accessible to children, pets and domestic animals, or in tamperproof bait boxes. Continue baiting until all feeding has stopped. Immediately replace liquid baits that become cloudy.

Caution:

Place in locations inaccessible to children, pets and domestic animals.

ACCEPTED
3-21-68
UNDER FEDERAL INVENTORY
FUNDING AGENCY
FOR FEDERAL ACQUISITION
EO UNDER NO. 9591-1

W. W. C. COMPANY