

PRO KILL MILL SPRAY

A Ready-To-Use Insecticide for Control of Insect Pests of Grain and Food Products

For use in Flour Mills, Elevators, Granaries, Food Processing Plants and Warehouses, Bakeries, Beverage Plants, Restaurants and Supermarkets. Food additive tolerances have been established for pyrethrins and piperonyl butoxide. When this product is used as directed, allowable tolerance levels will not be exceeded.

ACTIVE INGREDIENTS:

- Pyrethrins 0.2%
- *Piperonyl Butoxide, Technical 2.0%
- Petroleum Distillate 97.8%
- 100.0%

*Equivalent to 1.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION
Harmful if swallowed. Avoid contamination of feed and foodstuffs. The use of this product in food processing plants should be confined to time periods when the plant is not in operation. Foods should be removed or covered before treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Clean out any waste grain, dust, dirt or debris and destroy before spraying. Remove pets, birds and fish aquariums before spraying.

ENVIRONMENTAL HAZARDS
This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL
Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide spray mixture or residue that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides. Triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill. Consult Federal, State or local disposal authorities for rules and alternative procedures.

SURFACE SPRAYING: This spray provides rapid initial kill of exposed or accessible stages of the following insect primary pests: weevils, rice weevil, infested flour beetle, rusted flour beetle, stored grain beetle, grain weevil, flour beetle, and stored grain beetle. Apply to coarse grain or spray using the following

**SURFACE SPRAYING (continued)
application rate:**

APPROXIMATE USE RATE	
Spray	Area to Be Covered
1 Pint	93 3/4 square feet
1 Quart	187 1/2 square feet
1 Gallon	750 square feet

Spray walls, floors and other surfaces of bins and storage and handling areas thoroughly being especially careful to treat cracks, crevices and similar hiding places. Treat unloading, handling and processing areas and inside conveying, processing and handling equipment. Spray thoroughly around the base of heavy machinery and equipment. Spray insects directly when possible.

For in-transit protection, clean and spray truck beds, boxcars and ships' holds before loading. Thoroughly treat walls and floors and spray each layer of cartons and containers lightly as stacked.

To control Indian meal moth and Angoumois grain moth in stored grain apply to the surface of binned grain at the following application rate

APPROXIMATE USE RATE	
Spray	Area to Be Covered
1 Pint	125 square feet
1 Quart	250 square feet
1 Gallon	1,000 square feet

Apply at regular intervals with the first appearance of moths and continue until early fall. In case of severe infestation, break up webbing with rake before spraying and make a second application in two weeks.

As an aid in the control of mosquitoes, gnats, wasps and skippers, treat walls, windows, etc., as follows:

SEE REVERSE SIDE

screens, door and window frames, beams, light cords and similar resting places.

SPACE SPRAYING: To control flies, mosquitoes, gnats and flying moths, close windows, doors and other ventilation. Using a good sprayer adjusted to deliver a fine mist, direct spray toward the upper portions of the enclosure. Distribute spray thoroughly the entire area using 1/2 ounce spray per 1000 cubic feet of space. To control active exposed stage of running and crawling insects previously named, direct mist toward and into places commonly inhabited by these insects. Using a rate of 2 ounces per 100 cubic feet of space. Keep area closed for at least 10 minutes after spraying. Vacate premises and ventilate before re-occupying.

FOGGING: This oil base spray is suitable for application using conventional, thermal, mechanical or compressed air fogging equipment. Use as directed under "Space Spraying" filling the entire area with mist at a rate of 2 to 4 ounces per 1000 cubic feet of space, to control exposed or accessible stages of any insect named on this label.

NATIONWIDE
P.O. BOX 3227
HAMILTON, OHIO 45013

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED do not induce vomiting, contains petroleum solvent. Call a physician at once.
IF INHALED remove victim to fresh air. Apply respiration if indicated.
IF ON SKIN remove contaminated clothing and wash affected areas with soap and water.
IF IN EYES flush eyes with plenty of water. Get medical attention.
See side panel for additional precautionary statements.

EPA Reg. No. 9591-53
EPA Est. No. 9591-04-1
NET CONTENTS _____ GALLONS

BEST DOCUMENT AVAILABLE