

1/2 15 11-1

# PRO KILL Multi-Purpose Insecticide

Professional Strength - Low Dosage Required

For use in Schools, Hospitals, Institutions, Supermarkets, Dairies, Hotels, Restaurants, Drive-ins, Motels, Drug Stores, Food Plants, Milk Rooms and Animal Quarters

Provides Rapid Control of Flies, Mosquitoes, Small Flying Moths (Millers), Gnats, Wasps and accessible stages of Roaches, Ants, Water Bugs, Silverfish, Flour Beetles, Weevils, Grain Mites and Spiders

KILLS AND REPELS Stable Flies, Horse Flies, Face Flies, Deer Flies, House Flies, Horn Flies, Mosquitoes and Gnats

Controls Garden Insects on Ornamental and Vegetable Plants

Acceptable for Use to Day of Harvest on Edible Plants

#### ACTIVE INGREDIENTS:

Pyrethrins .....	0.5%
*Piperonyl Butoxide, Technical .....	5.0%
Petroleum Hydrocarbons .....	4.3%
INERT INGREDIENTS: .....	90.2%
	100.0%

\*Equivalent to 4% (butylcarbityl) (6-propylpiperonyl) ether and 1% related compounds

Keep Out Of Reach Of Children  
**CAUTION**  
See Side Panels for Additional Cautions

NET Weight 14 oz.

**NATIONWIDE**  
CHEMICAL PRODUCTS INC  
P.O. BOX 3027  
HAMILTON, OHIO 45013

ACCEPTED  
JAN 9 1987  
Under the Federal Insecticide, Fungicide and Rodenticide Act.  
Manufactured for the pesticide industry.  
9591-95

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

##### CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in food processing plants, wash all equipment, benches, shelving, etc. (where exposed food will be handled) with potable water. All food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

##### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, ponds, and streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

##### PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks, and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Turn off pilot lights in area of spraying.

##### STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES: Flush with water for 15 minutes. See physician if irritation persists.
- IF ON SKIN: In case of contact, remove contaminated clothing and wash skin with soap and water.
- IF INHALED: Remove victim to fresh air. Apply artificial respiration if needed.
- IF SWALLOWED: Do not induce vomiting. Seek medical attention immediately.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

Flies, Mosquitoes, Gnats, Wasps and Small Flying Moths (Millers): Close all doors and windows and spray room at the rate of 1 second per 1000 cubic feet. Direct spray to all parts of the room especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat as necessary.

Roaches, Waterbug, Silverfish, Crickets, Spiders, Centipedes and Sow Bugs: Spray into hiding places, such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Exposed Stages of Indian Meal Moth and Angoumois Grain Moth: Close doors and windows. Direct spray into all parts of room, especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

Exposed Stages of Flour Beetles, Weavils and Grain Mites: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary. Infested stored food products in the home should be disposed of.

Carpet Beetles: Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, molding and floors. Allow treated fabric to dry thoroughly before using. Repeat treatment as necessary.

Fleas: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only as recommended pet spray on pets.

To Treat Animals: To protect beef and dairy cattle from attacks of Stable Flies, Horn Flies, Deer Flies, Houseflies, Horn Flies, Mosquitoes and Gnats, spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from face flies, spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.

In Animal Quarters and Milk Rooms: For rapid control of House Flies, Mosquitoes, Gnats, Wasps and Small Flying Moths, close windows and doors and spray at the rate of 1 second per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

For Use in Gardens: Spray with slow sweeping motion, not closer than 18 inches from plants. Avoid wetting plants. Spray plants only when air is still. For single plants, spray in short bursts. Repeat as necessary for control. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat application at 7 to 10 day intervals or after heavy rains.

## USE ON

Roses	Aphids, Rose Chafers, Japanese Beetles, Rose Slugs, Tent Caterpillars, Scales
Asters	Leafhoppers, Ternish Plant Bugs, Lace Bugs, Thrips (exposed), Mealy Bugs
Dahlias	Aphids, Red Spider Mites, Plant Bugs
African Violets	Broad Mites, Citrus Mealy Bugs
Azaleas	Azalea Lace Bugs, White Flies, Leaf Miners
Euonymus	Euonymus Scales, Aphids, Thrips (exposed)
Hydrangeas	Aphids, Red Spider Mites, Hydrangea Leaf-tiers
Junipers	Juniper Scales, Red Spider Mites, Bagworms
Lilacs	Leaf Miners, Leaf Rollers, Scales
Tomatoes	Flea Beetles, Hornworms, Stink Bugs, Colorado Potato Beetles, Green Peach Aphids
Peppers, Egg-Plants, Potatoes	Colorado Potato Beetles, Leafhoppers, Blister Beetles, Vine Beetles, Green Peach Aphids
Cucumbers, Squash, Melons	Striped & Spotted Cucumber Beetles
Beans	Mexican Bean Beetles, White Flies, 12-Spotted Beetles
Bunt Corn	Japanese Beetles

## STORAGE & DISPOSAL

**Storage:** Store in a cool dry area away from heat and open flame.

**Disposal:** Wrap container and dispose of in trash. Do not incinerate.