

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (77-567) WASHINGTON, D.C. 20460	EPA REGISTRATION NO. 9591-130	DATE OF ISSUANCE JUL 16 1985
	TERM OF ISSUANCE	
NAME OF PESTICIDE PRODUCT Pro Kill Concentrate 2.25% + 22.5%		

NOTICE OF PESTICIDE: REGISTRATION
 REREГИSTRATION
 (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

C Nationwide Chemical Products, Inc. T
 P.O. Box 3027
 Hamilton, OH 45013

L J

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by other products if conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Add the phrase "EPA Registration No. 9591-130" to your label before you release the product for shipment.
- 2. Submit five (5) copies of our final printed labeling before you release the product for shipment.
- 3. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of the above conditions.

A stamped copy of the label is enclosed for your records.

Timothy A. Gardner
 Product Manager (17)
 Insecticide-Rodenticide Branch
 Registration Division (75-767)

Enclosure

BEST AVAILABLE COPY

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL *Timothy A. Gardner* DATE 7-16-85

PRO KILL Concentrate 2.25% + 22.5%

Contains 0.18 lb/gal of Pyrethrins (21.6 g/L) and 1.82 lb/gal of Piperonyl Butoxide (218 g/L)
FOR CONTROL OF HOUSE FLIES, MOSQUITOES, GNATS (MIDGES), WASPS AND SMALL FLYING MOTHS (MILLER MOTHS), WATERBUGS, ANTS, SILVERFISH, CRICKETS, FLEAS, EARWIGS, CENTIPEDES, SLOWBUGS (PILLBUGS), BOXELDER BEETLES, DEER FLIES, STABLE FLIES, HORSE FLIES, SUCKING AND BITING LICE, TICKS, SHEEP KEDS, BED BUGS, AND

Authorized by USDA For Use in Edible Product Areas of Official Establishments Operating Under The Meat, Poultry, Shell Egg Grading, and Egg Products Inspection Program.

ACTIVE INGREDIENTS:

Pyrethrins	2.25% w/w
*Piperonyl Butoxide, Technical	22.50%
INERT INGREDIENTS:	75.25%

100.00% w/w

*Equivalent to 18.0% (butylcarbetyl)(6-propylpiperonyl) ether and 4.5% related compounds.
 **U.S. Patent No. 3,683,078.

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada amplemente.
(TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

**KEEP OUT OF REACH OF CHILDREN
 WARNING**

SEE SIDE PANEL FOR ADDITIONAL
 PRECAUTIONARY STATEMENTS AND
 STATEMENT OF PRACTICAL TREATMENT

BEST AVAILABLE COPY

NATIONWIDE CHEMICAL PRODUCTS, INC.
 P.O. Box 3027, Hamilton, Ohio 45013
 EPA Reg No 9591- EPA Est No 9591-OH-2

JUL 16 1985

NET CONTENTS:

Enter the product name, strength, and brand name as required for the product registered under EPA Reg. No. 9591-130

**PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS
 WARNING**

WARNING: Harmful if swallowed or absorbed through skin. May cause eye injury. Do not get in eyes. Avoid contact with skin or clothing and wash after handling. Remove contaminated clothing and shoes. Do not apply directly to food or feedstuffs. In commercial establishments and food processing equipment during application while food processing is underway. After space spraying in processing plants, wash all equipment, benches, shelving and floors thoroughly with an effective cleaning compound. At home, all food processing surfaces and utensils should be thoroughly washed before use. Remove pets, and cover fish tanks.

STATEMENT OF PRACTICAL TREATMENT:
IF SWALLOWED: Call a physician or Poison Control Center. This product contains petroleum solvent. Aspiration may be a hazard. **IF IN EYES:** Flush with plenty of water. Get medical attention. **IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.

ENVIRONMENTAL HAZARD: This product is toxic to fish. Do not apply directly to water. Do not clean equipment or dispose of wastes in water.

PHYSICAL AND CHEMICAL HAZARD: Flammable. Do not use or store near heat, or open flame. Do not use in confined spaces. Flash point minimum of 60°F (15.5°C).

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with the label.

STORAGE AND DISPOSAL:
 Do not contaminate water, food or feed by storage or disposal. **STORAGE:** This material may stratify upon standing, even after shaking. Before each use, do not expose to temperatures below 40°F below 40°F (4.5°C), allow to stand until product reaches room temperature before using. **DO NOT USE OPEN FLAME.**
PESTICIDE DISPOSAL: Wastes resulting from the use of this product should be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: For Containers 1 Gallon and Under: Wrap container and put in trash. For Containers Over 1 Gallon: Triple rinse (or equivalent) and reuse for non-pesticidal purposes. Do not reuse for food or feed. Reconditioning, or puncture and dispose of in a sanitary manner approved by State and Local authorities.

DIRECTIONS FOR USE (Continued on next page)

DIRECTIONS FOR USE (Continued)

EPA Reg. No. 9591

INDOORS: For use in hotels, motels, homes, preschools, apartment buildings, commercial, office, institutional and storage buildings, recreational buildings, packaging and processing plants, milk handling areas of milk plants, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries, boats, ship cabins, and hospitals. In hospitals, patients should be removed prior to treatment. Also for use in campers, mobile homes, buses, automobiles, trucks, boxcars and ship cargo holds. Also for use in barns, stables, kennels, poultry houses, and other animal quarters. When spraying in animal quarters, **DO NOT SPRAY FOOD OR FEED ITEMS.**

GENERAL INFORMATION

Mix this product with water as shown in the **SPRAY DILUTION CHART**. Use the 0.15 + 1.5 spray to quickly reduce heavy infestations of flying and crawling insects and the 0.1 + 1.0 spray to control light infestations of flying and crawling insects and for maintenance purposes. For animal treatment, use the 0.1 + 1.0 spray against flying insects, and the 0.01 + 0.1 dilution to control lice, fleas, ticks and sheep keds. Poultry should only be treated with the 0.01 + 0.1 dilution.

NOTE: THIS MATERIAL MAY STRATIFY UPON STANDING, EVEN AT ROOM TEMPERATURE. THOROUGHLY MIX BEFORE EACH USE. DO NOT EXPOSE TO TEMPERATURES BELOW 40°F (4.5°C). IF EXPOSED TO TEMPERATURES BELOW 40°F (4.5°C), ALLOW TO STAND UNTIL PRODUCT REACHES ROOM TEMPERATURE AND THOROUGHLY MIX BEFORE USING. DO NOT USE OPEN FLAME. DO NOT EXPOSE THE SPRAY MIXTURE TO TEMPERATURES BELOW 32°F (0°C).

SPRAY DILUTION CHART

Amount of Spray Mixture Desired		Amount of Product Required to Achieve Desired Dilution		
Gallons	Liters	0.15 + 1.5	0.1 + 1.0	0.01 + 0.1
1	3.78	0-2/28 oz (280 ml)	8 fl oz (177 ml)	2/2 fl oz (30 ml)
3	11.37	4/5 qt (9.76 L)	17-1/2 fl oz (513 ml)	3 fl oz (90 ml)
5	18.96	1-1/3 qt (1.26 L)	1 qt (9.06 L)	3 fl oz (90 ml)
10	37.92	2-2/3 qt (2.53 L)	1-4/5 qt (1.79 L)	8 fl oz (177 ml)

NOTE: 1 fluid ounce = 2 tablespoons = 8 teaspoons = 29.574 ml

DO NOT APPLY THIS PRODUCT IN CONDUITS, MOTOR HOUSINGS, JUNCTION AND SWITCH BOXES OR OTHER ELECTRICAL EQUIPMENT BECAUSE OF POSSIBLE SHOCK HAZARD. DO NOT USE ON SURFACES THAT CAN BE HARMED OR STAINED BY WATER.

NOTE TO PEST CONTROL OPERATORS: One gallon (3.78 L) of spray is sufficient to treat 1500 linear feet (457 meters) with a 6 inch (15.2 cm) spray band.

Space Spray: To control House Flies, Mosquitoes, Gnats (Midges), Wasps, and Small Flying Moths (Millers): Close all doors, windows and ventilators. Adjust spray equipment to deliver a fine mist and direct the nozzle upward and outward in all directions. Fill the room with mist at the rate of 0.5 to 1 ounce of spray for each 1,000 cu ft (28.3 cu m) of space. Keep the area closed for 15 minutes, then thoroughly ventilate before entering the treated area.

To Control Bed Bugs and Mites in Poultry Houses: Apply as a crack and crevice and spot treatment with a coarse spray to areas where the bed bugs and mites hide, such as cracks in walls and floor, crevices, roost poles and nests.

Contact Spray: To kill cockroaches (waterbugs), fleas, earwigs, centipedes, sowbugs (pillbugs), boxelder bugs, ants, silverfish and crickets: Adjust spray to produce large wetting type spray particles. Direct spray to contact insects in cracks and crevices in floors, baseboards, around pipes, behind cabinets and other harborage. Kill as many insects directly as possible. Apply the spray to the point of run-off. Repeat as necessary.

PESTS OUT-OF-DOORS: DO NOT TREAT OR ALLOW THE SPRAY TO CONTACT PLANTS (lawns, flowering plants, shrubs, trees, evergreens, etc.) BECAUSE OF POSSIBLE INJURY.

HOUSE FLIES, MOSQUITOES, GNATS, FLYING MOTHS (MILLERS), AND WASPS OUTDOORS: For outdoor use as an aid in reducing annoyance from these insects in municipalities, zoos, parks, playgrounds, stadiums, open-air theaters, recreational areas, dump areas, manure and compost piles. Treat the area with 1.5 gallons per acre (14 L/ha). Spray garbage cans, outhouses, barbecue pits, compost and manure piles, and other breeding areas until surfaces are thoroughly wet. Spray outside surfaces of tents, lean-tos, and shelters in camping areas, the outside of barns, dairies, kennels, creameries, food processing plants, and institutions until surfaces are thoroughly wet, paying particular attention to screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around outdoor light fixtures on porches, in garages, and other places where these insects alight or congregate. Repeat as necessary.

ANIMAL TREATMENT: To kill and repel House Flies, Horn Flies, Face Flies, Deer Flies, Stable Flies and Horse Flies on Dairy and Beef Cattle, Horses, Dogs, Hogs, Sheep and Goats: Apply the

0.1% + 1.0% dilution as a coarse wetting spray at the rate of 2 ounces per large animal or a Jolly Bath. Apply sufficiently to wet the hair but not to soak the hide. If longer kill or repellency is desired, use up to 1-2 quarts (0.95 to 1.69 L) applied to all parts of an animal the size of a mature cow. This latter treatment will protect for 3-8 days from stable flies, horn flies, face flies, and house flies, and offers up to 3 days protection from horse flies and deer flies. Avoid spraying in eyes. Repeat as necessary. Any animals made nervous from the sound or feel of the spray can be treated by applying the material with a sponge. Applicator should wear protective clothing and gloves for sponging.

For Control of Sucking and Biting Lice, Fleas, Ticks and Sheep Keds on Dairy and Beef Cattle, Horses, Dogs, Hogs, Sheep and Goats. Use the 0.01% + 0.1% dilution as a dip or a spray. If used as a spray, make sure the animal hair is thoroughly wet including the head and tail. Repeat the treatment in ten days to kill newly hatched insects.

For Biting Lice, Fleas, Ticks, Bedbugs, and Mites on Poultry: Use the 0.01% + 0.1% dilution to treat the birds by dipping them in the solution while holding the wings away from the body to insure thorough coverage. To avoid chilling of the poultry, dip on a warm day, or in a warm building. Applicator should wear protective clothing and gloves.

BEST AVAILABLE COPY

WARRANTY STATEMENT

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such, or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse or such, or in combination with other materials.

EPA 450/5-77-001