105 5 pm3 Umjera 1

## FRONT PANEL

LIQUID WEED & FEED 20-0-0

NET CONTENTS ONE QUART (NET WEIGHT: 2 LBS 5 0%

--KEEP FROM FREEZING-1 QUART COVERS UP TO 5,000 SQ.FT.

9591-149

KILLS: Bedstraw, Black Medic, Buckhorn, Burdock, Chicory; Chickweed,
Clover, Dandelion, Dock, Ground Ivy, Heal-All, Heabit, Knotweed,
Lambsquarter, Lespedeza, Mallow, Morning Glory, Oxalis, Peppergrass, Pigweed,
Plantains, Poison Ivy, Poison Oak, Purslane, Ragweed, Sheep Sorrel,
Shepherd's Purse, Speedwell, Spurge, Thistle, Wild Carrot, Wild Garlic, Wild
Lettuce, Wild Onion, Yarrow.

The fast, easy way to weed control and fertilize your lawn.

No Spreading - No Complicated Mixing Just attach to a garden hose & spray

## ACTIVE INGREDIENTS:

### This Product Contains:

\*0.44 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 4.711%
\*\*0.211 lb. 2-(2 methyl-4-chlorophenoxy) propionic acid per gallon or 2.261%
\*\*\*0.048 lb. Dicamba (3,6-dichloro-o-anisic acid) per gallon or 0.514%

# KEEP OUT OF REACH OF CHILDREN

-- CAUTION--

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

See back of panel for additional precautionary statements.

Panel 1

# STOP. READ THE ENTIRE LABEL FIRST. **OBSERVE ALL PRECAUTIONS AND** FOLLOW DIRECTIONS CAREFULLY

MEASURING GUIDE

### PRECAUTIONARY STATEMENTS

!Hazards to Humans (and Domestic Animals): (CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or iclothing. In case of contact, immediately flush eyes or skin with plenty of water. If irritation develops get medical lattention. Avoid contamination of food or feedstuffs.

ج ، ز۔

LENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not ! lapply when weather conditions favor drift from target area. Do ! Inot contaminate water by cleaning of equipment or Jisposal of !wastes. Do not contaminate domestic or irrigation waters. 

# 2000 SQFT

### DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner incorsistent with its labeling.

ISTORAGE: Keep from freezing. Store in original container in a llocked storage area inaccessible to children. Do not store near l lifood or feedstuffs. To prevent cross-contamination, do not store ! Inear other fertilizers, insecticides, fungicides, or seeds IPRODUCT DISPOSAL: Securely wrap original container with unused | iproduct in several layers of newspaper and discard in trash. :CONTAINER DISPOSAL: Do not reuse empty container. Rinse 3000 SQFT!thoroughly before discarding in trash

# OTHER USE PRECAUTIONS:

Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not apply to newly seeded lawn grasses until well established. Do not spray when the temperature exceeds 85°F. can be safely sown 3-4 weeks after application. Do not contaminate 4000 SQFTdomestic or irrigation waters. Do not use on carpetgrass, dichondra, St. Augustine, Centipede, closely-mowed bentgrass, nor

on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species.

5000 SQFT COVERAGE WHEN EMPTY



Panel 2.

### 20-0-0

### GUARANTEED ANALYSIS

Net Weight: 2 lbs 5 oz.

Manufactured by: Nationwide Chemical Products, Inc.

P. D. Box 13027 Hamilton, Ohio 45013 EPA Registration No. 9591-EPA Est. No. 9591-DH-2

### WHERE TO USE:

LIQUID WEED & FEED may be used on Bahia, Bermuda, Bluegrass, Fescue, Rye or Zoysia lawns, vacant lots, playgrounds, and other similar areas where broadleaf weeds are present. This product is especially effective on all hard-to-kill weeds such as Poison Oak, Sheep Sorrel and Spurge. In addition LIQUID WEED & FEED controls most other broadleaf weeds, including sensitive weeds such as Chickweed, Clover, Dandelion, Knotweed and Plantain.

### WHEN TO USE:

- 1. Spray at any time during the growing season when weeds are actively growing.
- On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks before putting new grass seed or stolons down in the bare spots.
- Poor weed control may result if spray is applied during drought or just before rain.
- 4. For best results, mow 1 to 2 days before treating.
- 5. Do not water lawn within 24 hours after treatment.

### HOW TO USE:

SHAKE WELL BEFORE USING.

\*Connect spray nozzle to Garden hose; remove safety cap from container; attach tube to applicator and insert tube section. Attach container to end of nozzle.

\*Turn on water; place finger over hole on right side of nozzle and proper ratio of LIQUID WEED & FEED will automatically be dispersed. Spray directly over area requiring application evenly and uniformly.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, seller makes no claims other than those stated on this label.