

1 of 3
PM 23.

9591-150

FRONT PANEL

LIQUID WEED & FEED
FOR SENSITIVE GRASSES
20-0-0

NET CONTENTS HALF GALLON
(NET WEIGHT: 4 LBS. 12 OZ)

--KEEP FROM FREEZING--
ONE HALF GALLON COVERS UP TO 5,000 SQ. FT.

ACCEPTED
JUL 6 1987
9591-150

KILLS: Bedstraw, Black Medic, Buckhorn, Bull Thistle, Burdock, Buttercup, Chicory, Chickweed, Clover, Cocklebur, Dandelion, Dock, Galinsoga, Ground Ivy, Hawkweed, Heal-All, Henbit, Knotweed, Lambsquarter, Lespedeza, Mallow, Morning Glory, Mustard, Oxalis, Peppergrass, Pigweed, Plantains, Poison Ivy, Poison Oak, Purslane, Ragweed, Sheep Sorrel, Shepherd's Purse, Speedwell, Spurge, Tansy Mustard, Thistle, Velvet Leaf, Wild Carrot, Wild Garlic, Wild Lettuce, Wild Onion, Yarrow, and many other broad leaf lawnweeds

The fast, easy way to weed control and fertilize your lawn.

No Spreading - No Complicated Mixing
Just attach to a garden hose & spray

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid.....	1.872%
**Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid.....	0.556%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid).....	0.229%
INERT INGREDIENTS.....	97.343%
TOTAL	100.000%

This Product Contains:

- *0.105 lb. 2-(2 methyl-4-chlorophenoxy) propionic acid per gallon or 1.54%
- **0.220 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 0.449%
- ***0.024 lb. Dicamba (3,6-dichloro-o-anisic acid) per gallon or 0.190%

KEEP OUT OF REACH OF CHILDREN
-- CAUTION--
STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

See back of panel for additional precautionary statements.

Panel 1

~~STOP.~~ READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY

MEASURING
GUIDE

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals):
CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes or skin with plenty of water. If irritation develops get medical attention. Avoid contamination of food or feedstuffs.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

2000 SQFT

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children. Do not store near food or feedstuffs. To prevent cross-contamination, do not store near other fertilizers, insecticides, fungicides, or seeds

PRODUCT DISPOSAL: Securely wrap original container with unused product in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash

3000 SQFT

OTHER USE PRECAUTIONS:

Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not apply to newly seeded lawn grasses until well established. Do not spray when the temperature exceeds 85°F. Seed can be safely sown 3-4 weeks after application. Do not contaminate domestic or irrigation waters. Do not use on carpetgrass, dichondra, St. Augustine, Centipede, closely-mowed bentgrass, nor on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of trees and other ornamental species.

4000 SQFT

5000 SQFT

COVERAGE WHEN EMPTY

Panel 2.

20-0-0

GUARANTEED ANALYSIS

Total Nitrogen (N), Not less than.....20.00%
20% Urea Nitrogen

Net Weight: 4 lbs 12 oz.

Manufactured by: Nationwide Chemical Products, Inc.

P. O. Box 13027
Hamilton, Ohio 45013
EPA Registration No. 9591-
EPA Est. No. 9591-OH-2

WHERE TO USE:

LIQUID WEED & FEED may be used on Bahia, Bermuda, Bluegrass, Fescue, Rye or Zoysia lawns, vacant lots, playgrounds, and other similar areas where broadleaf weeds are present. This product is especially effective on all hard-to-kill weeds such as Poison Oak, Sheep Sorrel and Spurge. In addition LIQUID WEED & FEED controls most other broadleaf weeds, including sensitive weeds such as Chickweed, Clover, Dandelion, Knotweed and Plantain.

WHEN TO USE:

1. Spray at any time during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks before putting new grass seed or stolons down in the bare spots.
3. Poor weed control may result if spray is applied during drought or just before rain.
4. For best results, mow 1 to 2 days before treating.
5. Do not water lawn within 24 hours after treatment.

HOW TO USE:

SHAKE WELL BEFORE USING.

*Connect spray nozzle to Garden hose; remove safety cap from container; attach tube to applicator and insert tube section. Attach container to end of nozzle.

*Turn on water; place finger over hole on right side of nozzle and proper ratio of LIQUID WEED & FEED will automatically be dispersed. Spray directly over area requiring application evenly and uniformly.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, seller makes no claims other than those stated on this label.