

FUL SAN

# QUIK-SAN

## Disinfectant - Sanitizer - Deodorizer

For Food Processing Plants, Dairy Farm and Home Uses,  
Hospital, Institutional, School and Industrial Use

ACTIVE INGREDIENTS:  
Alkyl (C<sub>12</sub> 50%; C<sub>12</sub> 40%; C<sub>16</sub> 10%)  
Dimethyl Benzyl Ammonium Chloride

INERT INGREDIENTS

ACCEPTED  
02 MAY 1980  
Total 100.0%

KEEP OUT OF REACH OF CHILDREN  
DANGER

EPA REG. No. 9598-13

EPA EST. No. 09598 CA-1

(SEE ADDITIONAL CAUTIONS ON SIDE PANEL)

Manufactured by

**FULTS CHEMICAL, INC.**

Tulare, California 93274



NET CONTENTS 1 GALLON (3.785 LITERS)

### Directions for Use - GENERAL CLASSIFICATION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### Disinfectant in Hospitals, Nursing Homes and Other Health Care Institutions.

For disinfecting floors, walls, door handles, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard surfaces.

Add 4 1/4 oz. Quik-San to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth.

At this use level Quik-San is effective against *Pseudomonas aeruginosa*.

#### Disinfectant in Institutions (i.e. Educational, Penal, Military, Industry, Schools and Homes)

For disinfecting floors, walls, door handles, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard surfaces.

Add 3 oz. of Quik-San to 4 gallons of water. Apply to previously cleaned hard surface with mop or cloth.

At 3 oz. per gallon Quik-San is effective against *Staphylococcus aureus*, and *Escherichia coli*.

#### Disinfection of Barber Tools

Barber tools such as combs, clippers, razors, and scissors can be disinfected by immersing in a 1/2-oz./gallon solution of Quik-San.

#### Disinfection of Poultry Equipment, Animal Quarters and Kennels.

Poultry brooders, watering fountains, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 4 1/4 oz. Quik-San to 4 gallons of water with a mop, cloth or brush. Small utensils should be immersed in this solution. Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate brooders, coops and other closed spaces. Do not house poultry, or other animals, or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

BEST DOCUMENT A.

32