

**WARNING**  
**Precautionary Statements**  
**ENVIRONMENTAL HAZARDS**

Toxic to fish. Do not discharge into lakes, streams, ponds, or public waters unless in accordance with an NPDES permit. For guidance contact your Regional Office of the EPA.

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in an approved landfill, or bury in a safe place. Metal containers over 30 gallons should be resealed before offering for reconditioning.

**GENERAL:** Consult Federal, State or Local disposal authorities for approved alternative procedures.

EPA Registration No. 9601-12  
 EPA Est. No. 9601-OH-1


23

# SLIMEX EE

*For the biological control of Algae, Bacteria, Fungi, and Slime in recirculating water cooling towers.*

Active Ingredients:

Disodium cyanodithioimidocarbonate.....	7.85%
Potassium N-methyldithiocarbamate.....	10.15%
Inert Ingredients.....	82.50%
<b>TOTAL</b> .....	<b>100.00%</b>



RECEIVED  
 JUN 10 1980  
 U.S. DEPARTMENT OF AGRICULTURE  
 PESTICIDE ACTION CENTER  
 WASHINGTON, D.C. 20250

**GENERAL**

is a violation of the manner inconsistent with the use of SLIMEX EE is used to control algae, bacteria, fungi, and slime in recirculating water cooling towers. The water should be cleaned to remove slime, and other debris. Use 3.3 to 6.6 fluid ounces per 1000 gallons of water. Repeat dosage until control is achieved. Subsequent slug dosing of SLIMEX EE per label instructions should be employed every 1 to 2 weeks. Frequency of additional dosing should be determined by microbiological procedures made in the summer.

**NET CONTENTS**

**LABORATORY CONTROL**

**CH**  
 539 Stimme Road

RECEIVED  
 JUN 10 1980  
 U.S. DEPARTMENT OF AGRICULTURE  
 PESTICIDE ACTION CENTER  
 WASHINGTON, D.C. 20250

# SLIMEX EE

For the biological control of Algae, Bacteria, Fungi, and Slime in recirculating water cooling towers.

<b>Active Ingredients:</b>	
Disodium cyanodithioimidocarbonate.....	7.35%
Potassium N-methyldithiocarbamate.....	10.15%
<b>Inert Ingredients:</b> .....	82.50%
<b>TOTAL</b> .....	100.00%

REGISTERED  
 JUN 20 1980  
 U.S. PATENT OFFICE  
 TRADEMARK  
 No. 2,661,112

## Directions for Use GENERAL CLASSIFICATION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SLIMEX EE is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime, and other deposits. An initial slug addition of 3.3 to 6.6 fluid ounces of SLIMEX EE per 1000 gallons of water is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 1.1 to 6.6 fluid ounces of SLIMEX EE per 1000 gallons of water should be employed every 1 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

NET CONTENTS

LAB CONTROL NO

Manufactured by

**CHARDON**

539 Stimmel Road Box 1004 Columbus, Ohio 43216

REGISTERED

24

REGISTERED