

VERTEX

CONCENTRATE

A CHLORINE BEARING SODIUM HYPOCHLORITE SOLUTION FOR SANITATION AND ALGAE CONTROL IN SWIMMING POOLS, PUBLIC WATER SUPPLIES AND WASTE WATER SYSTEMS.

ACTIVE INGREDIENT:

SODIUM HYPOCHLORITE 9.0%
INERT INGREDIENTS 91.0%

CAUTION:

Keep Out of Reach of Children

SEE SIDE PANEL FOR
ADDITIONAL PRECAUTIONS

CONTENTS 1 GALLON

MANUFACTURED BY

VERTEX CHEMICAL CORPORATION
Dupo, Illinois 62239

CAUTION

May be harmful if swallowed. Skin and eye contact may cause severe irritation. Do not mix with acids or ammonia as toxic vapors may be emitted.

ANTIDOTE – EXTERNAL: in case of contact with skin and eyes, flush with plenty of water. **INTERNAL:** drink mucilage, raw egg white, milk or rice gruel. Follow with tablespoon of mustard in glass of warm water. **CALL A PHYSICIAN.**

STORAGE – Store in a cool dark location. Contains 10.3% available chlorine by weight when packed. Degrades with age. This container should be used within one month of manufacture date. Use a test kit and increase dosage as necessary to obtain required level of available chlorine. When container is empty, rinse thoroughly with water before discarding.

FISH AND WILDLIFE CAUTION – This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water. Apply this product only as specified on this label.

TABLE OF PROPORTIONS

0.1 ppm - 0.1 oz. in 1000 gallons water
0.5 ppm - 0.5 oz. in 1000 gallons water
1.0 ppm - 1.0 oz. in 1000 gallons water
5.0 ppm - 5.0 oz. in 1000 gallons water
10 ppm - 10 oz. in 1000 gallons water

REPACKAGING – This product is sold for repackaging.

BATCH DATE

EPA REG. NO. 9616-2

ACCEPTED
FEB 15 1973
UNDER THE FEDERAL INSECTICIDE
FUNCTIONAL AND CHEMICAL ACT
REGISTERED UNDER NO. 9616-2
SUBJECT TO ATTACHED COMMENTS.

VERTEX

CSS-10

DIRECTIONS FOR USE

DAIRY FARMS – The strength of chlorine-bearing Sodium Hypochlorite is dissipated on contact with organic substances or soil. All equipment should be cleaned before sanitizing with VERTEX CSS-10. Refer to Table of Proportions and rinse utensils, bulk tanks, milking machines, pipeline milkers and udders with 200 ppm solution of CSS-10. Contact time should be at least 2 minutes. For udders, use a clean single service towel for each cow. Never dip used towel back into solution. Discard solution if visibly dirty. Prepare fresh solution for each milking. Dry with a clean towel or allow udder to air dry before attaching milker.

FOOD AND DAIRY PROCESSORS – After cleaning and before use, sanitize all nonporous surfaces with 200 ppm and all porous surfaces with 800 ppm solution of CSS-10. Rinse all surfaces with potable water supply following sanitation rinse. For mold control, a spray rinse of 500-10000 ppm is recommended.

RESTAURANTS AND TAVERNS – After washing with dishwashing detergent and rinsing with clean water, immerse utensils in 200 ppm solution of CSS-10 for at least 2 minutes. Allow utensils to air-dry.

STATE AND LOCAL REGULATIONS – Consult your dealer, state or local health authorities for additional information.

TABLE OF PROPORTIONS – AVAILABLE CHLORINE

- 200 ppm – 1 fluid oz. in 3¼ gallons water
- 800 ppm – 4 fluid oz. in 3¼ gallons water
- 5000 ppm – 30 fluid oz. in 4 gallons water
- 10000 ppm – 60 fluid oz. in 4 gallons water

A CHLORINE BEARING SODIUM HYPOCHLORITE SOLUTION FOR SANITIZATION IN THE DAIRY, FOOD PROCESSING AND FOOD SERVICE INDUSTRIES.

ACTIVE INGREDIENT:

SODIUM HYPOCHLORITE 9.0%

INERT INGREDIENTS 91.0%

CAUTION:


Keep Out of Reach of Children

SEE SIDE PANEL FOR
ADDITIONAL PRECAUTIONS

CONTENTS 1 GALLON

MANUFACTURED BY
VERTEX CHEMICAL CORPORATION

Dupo, Illinois 62239



May b
irritati
or am

FIRST
flush v
white,
glass o

STOR.
able c
contain
Use a
level of
with w

FISH
Treat
lakes,
specifi

REPA
BATC
EPA R
EPA E