

CLOW**ACCEPTED**

AUG 15 1980

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 9640-22

Water Management Division
Clow Corporation
408 Auburn Avenue
Pontiac, Michigan 48058**CLOW 3808****MICROBIOCIDICIDE**

EPA REG. NO. 9640-22

EPA EST. NO. 9640-MI-1

LOW FOAM WATER TREATMENT MICROBIOCIDICIDE**Active Ingredients:**

Diocetyl dimethyl ammonium chloride:	50%
Ethyl alcohol	10%

Inert Ingredients:

40%
100%

KEEP OUT OF REACH OF CHILDREN**DANGER****STATEMENT OF PRACTICAL TREATMENT:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****DANGER:** Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.**ENVIRONMENTAL HAZARDS:** This pesticide is toxic to Fish. Do not discharge treated effluent into lakes, streams, ponds, or public waters unless in accordance with an NPDES permit. For guidance contact your Regional Office of the EPA.**PHYSICAL HAZARDS:** Do not use, pour, spill, or store near heat or open flame.**Directions for Use****GENERAL CLASSIFICATION**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

To control algae and algae slimes, use **CLOW 3808** as directed. For best results, slug feed. The frequency of addition of microbiocide needed depends on many factors. To optimize your use of **CLOW 3808**, follow this procedure:

- Initially use 16 fluid ounces per 1000 gallons of water to be treated (60 ppm active quaternary).
Should the above dosage not give satisfactory results, use 21 fluid ounces per 1000 gallons of water.
Repeat the initial dose every seven days or increase the frequency if needed.
- When the above treatment level is successful, use 6 to 9 fluid ounces per 1000 gallons of water to maximize efficiency. Repeat weekly as needed.
Should slime develop again, return to initial dosage.

Cooling tower waters that are inherently low in algae growth may be adequately controlled by the lower range of these dosages, slug feed every seven days.

Dilute the appropriate amount of **CLOW 3808** in 1 to 2 gallons of water then add to the sump of the tower. Note: This product weighs 8.0 lbs. per gallon (at 20° C.).

Should tower be heavily fouled, a precleaning is required.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. **PESTICIDE****DISPOSAL:** Pesticide that cannot be used, or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.**CONTAINER DISPOSAL:** Dispose of in an incinerator or landfill approved for pesticide containers, bury in a safe place or return to drum reconditioner.

0480

FOR INDUSTRIAL USE ONLY

Lbs. Net Wt. 39