

ACTIVE INGREDIENTS:

Sodium Cacodylate 22.2%
Dimethylarsinic Acid
(Cacodylic Acid) 3.8%

INERT INGREDIENTS:

74.0%
Total Arsenic (as
elemental) all in water
soluble form 12.5%

Product contains the equivalent
of 2.48 Lbs. cacodylic acid per
gallon.

Net Contents:

EPA Reg. No. 9003-16

ACCEPTED

GENERAL INFORMATION: DILIC Herbicide is useful for general post-emergent weed control. It is non-selective in its action. Its phytotoxic properties are quickly inactivated on contact with soil. It contains a surfactant. *It is unnecessary to add any other surfactant to the spray solution.* Best results are obtained on young actively growing weeds. It produces top-kill only so repeat applications are required for season long weed control of perennials.

MIXING INSTRUCTIONS: DILIC Herbicide is completely water soluble. Any spray equipment that gives good coverage may be used. Fill the spray tank about half full with water and add the required amount of herbicide with agitation. Finish filling the tank with water and apply. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period.

DIRECTIONS FOR USE

NON-CROP: DILIC Herbicide is useful for general weed control on drainage ditchbanks and rights-of-way along sidewalks, driveways and fences, around buildings and ornamentals, and on similar non-crop areas. Mix 1 to 2 gallons of DILIC Herbicide in 100 gallons of water. Spray unwanted vegetation to just short of run-off. If regrowth occurs reapply as re-

quired. Do not allow spray to come in contact with foliage, green bark, grafted unions, or scuffed, damaged or broken bark when spraying around trees and ornamentals.

AGRICULTURAL PLANTINGS — CITRUS (Except Florida): DILIC Herbicide is useful as a directed application in non-bearing citrus orchards such as orange, grapefruit, tangerine, lemon and lime orchards. It should be applied at the rate of 1-1/2 to 2 gallons per acre.

Mix 1 to 2 gallons of DILIC Herbicide in 50 to 100 gallons of water. Apply as a directed spray in inter-spaces and around base of trees. Spray unwanted vegetation to just short of run-off. If regrowth occurs, reapply as required, however, do not exceed 3 applications per year.

Do not allow spray solution to contact leaves, stems or bark. Use a shield if necessary, for nursery plantings or young trees. Do not apply around trees from which fruit will be harvested within one year of treatment.

Sodium Cacodylate Liquid
Plus Surfactant For General
Post-Emergent Weed Control

DILIC
HERBICIDE



READ ENTIRE LABEL
BEFORE USING THIS PRODUCT

CAUTION**Keep Out of the Reach of Children.**

Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.

Do not contaminate waters used for domestic consumption, or by animals, wildlife and aquatic life, or for irrigation purposes. Do not graze treated areas to livestock.

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

LAWN RENOVATION: Mow lawn to about 1 inch high before treatment. Mix 10 fluid ounces of DILIC Herbicide in 4 gallons of water and apply to 1,000 sq. ft. Spray foliage thoroughly, preferably on a warm sunny day. Foliage will usually turn brown in about five days. Bermudagrass and other deep-rooted perennials, or partially sprayed foliage, may require a second treatment. Reapply five days later if green areas remain.

When foliage is all brown, matted areas should be raked to remove dead vegetation. The lawn may then be re-established according to local practice. *Keep children and pets off treated areas until after first rain or sprinkling following treatment. Do not track from treated to untreated areas.*

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent, crush if possible, and discard in a safe place.

NOTICE: Use of this material in a manner, or at any time other than in accordance with the directions may produce plant injury or other undesirable results. Crystal Manufacturing Corporation makes no express or implied warranties, other than specified on this label.

CRYSTAL MANUFACTURING CORPORATION



1525 N. Post Oak Road • Houston, Texas 77055