

Lite Weight Crabgrass Preventer
20 lbs. Net Weight
Covers up to 2500 Sq. Ft.

Contains Dacthal

Total Nitrogen (N)	10.00%
Available Phosphoric Acid (P ₂ O ₅)	6.00%
Soluble Potash (K ₂ O)	4.00%

Active Ingredients:

* Dimethyl ester of tetrachloroterophthalic acid	2.88%
** Technical Chlordane	0.79%
Total Active Ingredients	3.67%
Inert Ingredients (Plant Food)	96.33%
TOTAL	100.00%

- * Use as a pre-emergence herbicide patented under U.S. Pat. 2,923,634
- ** Equivalent to 0.47% Octachloro - 4.7-methano - tetrahydroindane and 0.32% of related compounds.


CAUTION: Keep out of the reach of children.
See back panel for additional CAUTIONS.

U.S.D.A. Registration Number


Mfg. By Occidental Chemical Co., P.O. Box 636,
Springfield, Ohio 45501

NOTE: (caution will be capitals - 18 point - balance 12 point type)

SIDE I



10-6-4
Lite Weight - Plant Food
Crabgrass Preventer - Insect Control
Covers 2500 Square Feet



1. First Action - FEEDS - contains three primary plant foods for healthier, greener lawns.
2. Second Action - CRABGRASS PREVENTER - contains Dacthal which is a highly effective pre-emergence herbicide.
3. Third Action - INSECT CONTROL - contains chlordane which controls many insects.

Weeds Prevented:

When applied as recommended, Triple Action Crabgrass Preventer will give excellent control of crabgrass and the following weeds in turfs: Carpet-weed, Common chickweed, Fall panicum, Florida pussley, Green foxtail, Hairy crabgrass, Lambsquarters, Lovegrass, Purslane, Smooth crabgrass, Texas millet, Witchgrass, Yellow foxtail and Goosegrass.

Insects Controlled:

Controls the following insects - ants, white grubs, Japanese beetle grub, sod webworm, June beetle grub, May beetle grub, white fringe beetle and wireworms. Also, box elder bugs, chiggers, ticks, millipedes, imported fire ants.

When to Apply:

Triple Action Crabgrass Preventer kills crabgrass and other annual weeds as they germinate, but not after they have rooted. To be effective, Triple Action Crabgrass Preventer must be applied before weed seed germination in early spring. This usually coincides with the time when Forsythia blossoms start to fall. A single spring application should suffice for most geographic locations. Make application on a calm day to avoid wind drift.

How to Apply:

Before making application, lawn should be raked to remove extraneous material and insure uniform application. In order to obtain expected results, Triple Action Crabgrass Preventer should be evenly applied with no skips or misses. It is suggested that the area be covered twice; the second time at right angles to the first to insure thorough coverage*. Do not rake the lawn or disturb soil surface after application. Water in lightly if no rain falls within a week after application.

Re-Seeding:

When some spring seeding is necessary, there should be a delay of approximately 60 days after the application of Triple Action Crabgrass Preventer. This delayed seeding improves the survival of desirable grass seedlings. If re-seeding is done before 60 days after application the amount of seed should be increased 50% to insure thrifty stands. An early spring application may be made to newly seeded lawns after they have exhibited greening of newly sprouted lawn grasses.

SIDE II (Continued):

Caution: Keep out of the reach of children.
Avoid contact with eyes.
Keep children and animals off treated areas until this material
is watered into the soil and the grass is thoroughly dry.
Harmful if swallowed.
Avoid contact with the skin; in case of contact with skin, wash
with soap and water.
Avoid breathing dust.
Do not take internally.

Avoid contamination of feed and foodstuff. Do not pour, store or use near
flame. Thoroughly clean equipment after use. Destroy containers when empty,
by burning. Wash after using.

This product is toxic to fish and wildlife. Keep out of lakes, streams and
ponds. Birds feeding on treated areas may be killed.

Do not use on Dichondra or Toronto and Cohansey Bents or on newly seeded lawns.

Spreader settings may vary with age, model and condition of spreader. Make
a trial check on a measured area of 1000 square feet with 8 pounds of this
product. Avoid overlap.

SUGGESTED SPREADER SETTINGS

Scott.....8 *(6½) Jackson.....7½ *(6) Vaughn.....7½ *(6)
Cyclone.....6½ *(5) N.K.....5½ *(4)
*(Setting for application in two directions)

Apply preferable in late winter or early spring when the grass is dry. Do
not apply to areas that will be used for gardening or where run-off will be onto
a garden area. Do not use on dichondra lawns or on newly seeded lawns.

For content guarantee see front panel. The guarantee of Manufacturer or
seller is only that the material contained herein conforms to the Chemical
described on the label and is reasonably fit for the use herein described
when used in accordance with the direction for use. Manufacturer and seller
make no other expressed or implied guarantee, including any other express or
implied guarantee of fitness or of merchantability.

Mfg. by Occidental Chemical Co., P.O. Box 636, Springfield, Ohio 45501

