

RIVERSIDE INDUSTRIES

MARKS, MISSISSIPPI

MARKS PHONE
FA 6-2411

ACCEPTED

RIVERSIDE
PEANUT INSECTICIDE FUNGICIDE

(*Sevin-Sulphur-Copper)

1979 62

ACTIVE INGREDIENTS:

*Carbaryl (1-Naphthyl-N-methylcarbamate)	10.0%
Copper (In Basic Copper Sulfate) Expressed as Metallic	3.4%
Sulphur (Expressed as "S") (Sulphur 93% passed 325 mesh)	65.0%

INERT INGREDIENTS:

21.6%

TOTAL

100.0%

*Sevin, Trade Mark of UCC for 1-Naphthyl-N-methylcarbamate.

Effective for the control of leaf spot (*Cerospora Arachidicola*)
Velvet Bean Caterpillar or the Peanut Worm.

WARNING: KEEP OUT OF REACH OF CHILDREN. See other Cautions on Right Panel.

WHEN TO BEGIN: When First Spots appear on Basal Leaves.

NUMBER OF APPLICATIONS: Usually 4. Applied at about 10 day intervals,
the first dusting should be about the time of last cultivation.

AMOUNT OF MATERIAL: From 20 to 30 pounds of dust per acre for each
dusting depending on size of plant.

CAUTION

HARMFUL IF INHALED OR SWALLOWED.

Do not breathe dust. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands before eating. Take a shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOODSTUFFS. Avoid contamination of feed, feeding troughs, and watering receptacles.