Please read Instruction on	reverse before comp	leting form.	Fo	m Appro	oved. O	MB N	lo. 2070	0-0060. A	Approval expires 11-30-9
**EPA	Was	invironmental esticide Progr shington, DC ation for f	rams (H750 : 20460)5C)		X		stration ndment r	
	1		Section	1			-	-	<u> </u>
1. Company/Product Numb				. EPA Pr	oduct N	Manag	er	3. [Proposed Classification
9688-89				. LaRoc			•	-	•
4. Company/Product (Name	e)		P	M#					X None Restricte
Chemsico Diazinon 2G				3					
5. Name and Address of Al Chemsico, Division of Un P.O. Box 15842 St. Louis, MO 63114	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b) (I), my product is similar or identical in composition and labeling to: EPA Reg. No.								
Check if t	this is a new address	;	P	roduct N	lame _				
			Section		_				
Amendment - Explain below Resubmission in response to Agency letter dated				Final printed labels in response NOTIFICATION Agency letter dated "Me Too" Application					
X Notification - Explain b				"Me Too" Application Other - Explain below MAY 1 8 1998					
Explanation: Use additional Notification per P.R. Noti			ingredients	stateme	ent to de	elete	chemica	al name.	
			Section	111					
1. Material This Product Wi	ill Be Packaged In:	•							
Child-Resistant Packaging	Unit Packaging	_	Water Sol	luble Pac	kaging		2. Type	e of Conta —	ainer
Yes* No	Yes No		N.	No Plas Glas			Metal Hastic Glass		
*Certification must be submitted.	If "Yes", Unit Package wgt.	No. per container	If "Yes", Package v	wgt.	No. p conta r			Paper Other (S	Specify)
3.Location of Net Contents	Information	4. Size(s) of	f Retail Cor	ntainer	<u> </u>	5. Lo	cation c	of Label Di	irections
Label	Container						On La		companying product
6. Manner In Which Label is	s Affixed to Product		Lithograph Paper glue Stenciled			Othe	r(}
		•	Section I						
1. Contact Point (Complete i	items directly below f	for identificati	ion of indivi	idual to b	be conta	acted			• •
Name		· <u>-</u>	Title					Telephone Code)	e No. (Include Area
Kathie J Tryson			Manager, Registrations 800-242-11				1166		
I certify that the statements I acknowledge that any kno under applicable law.			attachments nt may be p						
2. Signature	- つ,	•	3. Title					_ ,,	
Bartin	Manager, Registrations					`			
4. Typed Name		· ,	5. Date						

May 6, 1998

Kathie J. Tryson



Diazinon 2G Lawn Insect Control

For outdoor use on the home lawn only
This product must not be used on golf courses and sod farms.
Ready to use • Contains 2% Diazinon Insecticide


Kills Home Invading Pests—Ants, Brown dog ticks, Fleas, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs.

Effective Ant Control • Kills ants and other insects in lawns & gardens • Kills lawn insects—As listed on back

Treats up to 200 square feet

KEEP OUT OF REACH OF CHILDREN

See additional precautionary statements on back



Chemsico Diazinon 2G kills:

Ants
Armyworms
Bermudagrass mites
Billbugs
Brown dog ticks
Chiggers
Chinch bugs
Clover mites
Crickets

Cutworms
Digger wasps
Earwigs
Fire ants**
Fleas
Lawn moths
Leafhoppers
Millipedes
Sod webworms

Sowbugs (Pillbugs) Springtails White grubs of: • Japanese beetle

European chafer
 Southern chafer

"Aids in control

STOP, READ ENTIRE LABEL BEFORE USE

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

LAWN APPLICATION

Spectracide controls lawn pests on both grass and dichondra lawns.

Special Precautions To Reduce Risk To Birds

Birds, especially waterfowl (ducks or geese), feeding or drinking on treated areas may be killed. To reduce risks to birds:

- 1. Do not treat areas larger than 15,000 sq. ft.
- Because of the migratory habits of certain waterlowl, do not apply in Nassau County, New York between November 1 and May 20.
- Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterlowl.
- Avoid overlapping treated areas.
- Follow all watering instructions in the directions for use; however, stop watering before puddling occurs.
- Do not apply this product to sites occupied by waterfowl or to sites within 75 ft. of any body of water that may attract waterfowl, such as takes or ponds.
- If it is necessary to repeat application, wait a minimum of 3 weeks between applications. Make no more than 4 applications per year.

Apply at the rate of 1 lb. per 200 sq. ft. (an area 20' by 10'), or 8 oz. per 100 sq. ft. (an area 10' by 10').

NOTIFICATION
MAY 1 8 1998

Ants, Armyworms, Bermudagrass mites, Brown dog ticks, Chiggets, Chinch bugs, Clover mites, Crickets, Cutworms, Earwigs, Fleas, Lawn moths (sod webworms), Leafhoppers, Millipedes, Sowbugs (pillbugs) and Springtails (Collembola): Apply when grass is dry. Water lawn lightly after application. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year.

Billbugs, and White grubs (Japanese beetle, European chafer, Southern chafer, Ataenlus spretulus [dung beetle]): Water grass with at least ¼-inch of water after application; however, stop watering before puddling occurs. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in Spring or early Summer when damage is first noticed. Water grass with at least ¼-inch of water after application; however, stop watering before puddling occurs. Treat again in late Summer or early Fall, if a second generation occurs.

Hyperodes weevils: Water grass with at least ¼-inch of water after application; however, stop watering before puddling occurs. Treat problem areas in mid-April and again in mid-May.

OPTIONAL SPREADER DIRECTIONS FOR OVERALL LAWN APPLICATION

Use 5 lbs. per 1,000 sq. (t. (an area 20' by 50'). Apply with a fertilizer spreader. See spreader settings below. For application with other types of equipment, also apply at a rate of 5 lbs. per 1,000 sq. ft (an area 20' by 50'), 1 lb. per 200 sq. ft. (an area 20' by 10'), or 8 oz. per 100 sq. ft. (an area 10' by 10').

Ants, Armyworms, Bermudagrass mites, Brown dog ticks, Chiggers, Chinch bugs, Clover mites, Crickets, Cutworms, Earwigs, Fleas, Lawn moths (sod webworms), Leafhoppers, Millipedes, Sowbugs (pillbugs) and Springtails (Collem-bola): Apply when grass is dry. Water lawn lightly after application. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year.

Billbugs and White grubs (Japanese beetle, European chafer, Southern chafer, Ataenius spretulus [dung beetle]): Water grass with at least %-inch of water after application; however, stop watering before puddling occurs. For billbug control, apply when activity is first observed or when chowoid or brown grass Indicates damage. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in Spring or early Summer



when damage is first noticed. Water grass with at least ¼-inch of water after application; however, stop watering before puddling occurs. Treat again in late Summer or early Fall, if a second generation occurs.

Hyperodes weevils: Water grass with at least ¼-inch of water after application; however, stop watering before puddling occurs. Treat problem areas in mid-April and again in mid-May.

SPREADER SETTINGS FOR LAWN APPLICATIONS					
Spreader Make & Model	5 lbs./1,000 sq. ft.				
Central Imperial Mark No. 1662-G	7 1/2				
Central Suburbia Lawn Spreader No. 1622-0	7 ½				
Cyclone Model B1 Rotary	6				
Greenfield Model 105	8				
J.C. Penney Model 6005	8 1/2				
J. C. Penney Rotary Spreader	8				
Scott's Precision Flow Spreader PF-1	6 1/2				
Sears Craftsman Model 671.19198	7 1/2				

NOTE: These settings are approximate, due to variability and wear of spreader. Check settings and calibrations on a small area before making total application.

SPOT TREATMENT

Ants: Apply 1 tablespoon over and around each individual ant hill.

Fire ants: To aid in the control of fire ants in lawns and other outdoor home recreation areas, apply 1% cup over, and 2 ft. around, each fire ant mound. Immediately after application, water in thoroughly. Apply water gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. A minimum of 1 gallon of water should be sprinkled over each mound area. Reapply as necessary. NOTE: For best results, apply in cool weather, 65–80° F, or in early morning or late evening hours. Treat new mounds as they appear. Digger wasps: Apply 1 tables boon 1 and around each individual ground nest opening. Apply 2 tables boon 1 and around each individual ground nest opening. Apply 2 tables boon 1 around each individual ground nest opening. Apply 2 tables boon 3 are not active.

BAND TREATMENT AROUND HOUSE FOUNDATION

The ploying pests are commonly found around or near house foundations and may enter the house?

Ants	Earwigs	Sowbugs		
Brown dog ticks	Fleas	(Pilibugs)		
Clover mites	Millipedes	Springtails		
Crickets	,	(Collembola		

Apply a 2-foot band of granules next to the house foundation. Distribute granules uniformly, using a rate of 8 oz. for each 100 sq. ft. of area (2' by 50') treated. If necessary, repeat application at 3-week intervals.

STORAGE AND DISPOSAL

Storage: Store in original container in a cool, dry, well-ventilated area, inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, seeds, food or feed. Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash. Container Disposal: Do not reuse empty container. Discard container in trash.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin, or clothing. Wash thoroughly after handling and before smoking or eating. May cause contact sensitization following repeated contact with skin of susceptible individuals. If sensitization reactions result, contact a physician. Do not store near food or feed products. Food utensils such as tablespoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow people or pets on treated areas until granules have been watered in and the grass has detect.

Statement of Practical Treatment

If Swallowed: Call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing. If Inhaled: Remove victim to fresh air and apply artificial respiration, if indicated. If on Skin: Wash promptly with soap and water. Rinse thoroughty. If In Eyes: Rinse eyes with plenty of water and call a physician immediately. Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl (ducks or geese), feeding or drinklog on treated areas may be killed. Because of the migratory habits of cortain Atlantic Coast waterfowl,

do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds. Avoid overlapping granules. Follow all watering instructions in the directions for use; however, stop watering before puddling occurs. Do not apply this product to sites occupied by waterfowl, or within 75 ft. of any body of water which may attract waterfowl, such as ponds or lakes. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to fish and other aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Questions or comments? Call 800-332-5553

Made in the USA

by Chemsico, Division of United Industries Corp., P. O. Box 15842, St. Louis, MO 63114-0842

UPC SYMBOL 0 00000 0

EPA Reg. No.: 9688-89 EPA Est. Nos.: 10370-TX-1 F , 9198-OH-1 M

, 61282-WI-1 H., 44616-MO-1 G, 961-PA-1 L , 9198-OH-2 B, 769-GA-1 S Superscript used is first letter of lot number

12 20

Form R2 © 1998

