Inchworms, Leafminers, Leafrollers, Mealybugs, Spider mites,

hemsico

AEROSOL MP

For outdoor residential use only.

[New! Fast Acting • Dual disease and insect control (action) . Disease plus insect control (killer) . Prevents and cures diseases on roses, flowers and ornamental shrubs • Protects roses, flowers and ornamental shrubs . Perfect for roses, flowers and ornamental shrubs • Can be used on delicate ornamentals • Ideal for use on delicate ornamentals . Controls powdery mildew, black spot, rust, blights and leaf spot . Prevents and cures black spot, rust, powdery mildew, blights and other listed diseases • Protects roses, flowers and ornamental shrubs from ants, aphids, chinch bugs, spider mites and many (other) [more] similar insects • Kills harmful insects on contact • Quickly kills insect[s] [pests] on contact • Kills aphids, whiteflies, spider mites, beetles and other harmful insects · Protects (for) up to 2 weeks · Can't be washed (off) away by rain · Water-Based)

Active Ingredients:	
Myclobutanil	0.012%
Permethrin	0.020%
Other Ingredients:	99.968%

KEEP OUT OF REACH OF CHILDREN CAUTION See back for additional precautionary statements

NET WT. 15 OZ.[/426tg]: _

OPTIONAL MARKETING LANGUAGE

IOPTION A

Chemsico Aerosol MP protects roses, flowers, ornamental shrubs from damaging diseases and insects. Its exclusive formula can actually prevent diseases and is ideal for use on delicate ornamentals.

OPTION B

Chemsico Aerosol MP prevents and stops a wide variety of diseases and insects on roses, flowers and shrubs. It can't be washed off by rain or sprinklers once dried and it works against diseases and insects for up to 2 weeks. Use this product throughout your yard to keep plants thriving.

Prevents and cures . . . Black Spot. Powderv Mildew. Rust, Blight and Leaf Spot

Also kills . . . aphids, spider mites, leafhoppers . . . and many [other] [more] similar insects!]

STOP, READ ENTIRE LABEL BEFORE USE. DIRECTIONS FOR USE

It is a violation of Fuderal law to use this product in a manner inconsistent with its labeling.

- Shake well before using.
- 2. Begin early in the season. Start treating when insects first appear or conditions favoring disease start to develop.
- 3. Hold can upright and at least 18 inches from foliage while
- Can be used on: Ageratum, aster, azalea, begonia, carnation, chrysanthemum, iris, lilac, marigold, mock-orange, pansy, petunia, privet, rhododendron, rose, salvia, snapdragon, spirea, sunflower, zinnia.
- 5. Spray with a slow sweeping motion, covering both upper and lower leaf surfaces and the stem.
- For best results, repeat every 10 14 days.

INSECTS CONTROLLED

Aphids, Bagworms, Cladas, Fall cankerworms, Fall webworms,

Under the Poderal Inensitie blo. Lunglishte, and Hodennettde Act, as unumded, for the permitte registered under EFA Rog. No. 1688-123

Thrips (exposed), Whiteflies

DISEASES CONTROLLED

PLANTS	DISEASES
Rose (1)	Black spot, Powdery mildew, Rust
Azaiea ⁽²⁾ , Rhododendron ⁽²⁾	Petal blight (ovulinia spp.) Powdery mildew
Carnation, Mock-orange, Pansy, Petunia, Salvia, Snapdragon, Sunflower	Powdery mildew, Rust
Ageratum, Aster, Chrysanthemum, Columbine, Cornllower, Iris, Marigold	Rust
Begonia, Lilac, Spirea	Powdery mildew
Privet, Zinnia	Cercospora leaf spot, Powdery mildew

- (1) Treat on a 7 to 10 day schedule. If black spot is not a problem, a 10 to 14 day schedule is recommended.
- (2) Begin spraying when flowers start to show color.

USE TIPS

- ✓ When using this product on plants not listed, test on a small
- ✓ Do not spray if temperature is above 90 OF as damage to some delicate plants can occur.
- Spray early in the morning or late in the evening, when air is
- ✓ If it rains before product has dried, re-apply.
- Regular light sprayings are more effective than a single heavy treatment.

SENSITIVE PLANT CARE

Use with care on ferns, palms and fuchsia. As with all plants, especially carnations, chrysanthemums, and roses, there exists a wide variety of species, and therefore, a wide variety of susceptibility to injury. To test for plant sensitivity, spray a few plants as described above, wait 24 hours, and check for pos-

[Bracketed Text] = Optional Language

page 1 of 2

sible injury.

OPTIONAL FORMAT

USE ON

Ageratum, aster, azalea, begonia, carnation, chrysanthemum, iris, lilac, marigold, mock-orange, pansy, petunia, privet, rhododendron, rose, salvia, snapdragon, spirea, sunflower, zinnia.

INSECTS CONTROLLED

Aphids, Bagworms, Cicadas, Fall cankerworms, Fall webworms, Inchworms, Leafminers, Leafrollers, Mealybugs, Spider mites, Thrips (exposed), Whiteflies

DISEASES CONTROLLED

PLANTS	DISEASES
Rose (1)	Black spot, Powdery mildew, Rust
Azalea (2), Rhododendron (2)	Petal blight (ovulinia spp.) Powdery mildew
Carnation, Mock-orange, Pansy, Petunia, Salvia, Snapdragon, Sunflower	Powdery mildew, Rust
Ageratum, Aster, Chrysanthemum, Columbine, Cornflower, Iris, Marigold	Aust
Begonia, Lilac, Spirea	Powdery mildew
Privet, Zinnia	Cercospora leaf spot, Powdery mildew

- Treat on a 7 to 10 day schedule. If black spot is not a problem, a 10 to 14 day schedule is recommended.
- (2) Begin spraying when flowers start to show color.

HOW TO TREAT

- Shake well before using.
- 2. Begin early in the season. Start heating when invects (1st

appear or conditions favoring disease start to develop.

- Hold can upright and at least 18 inches from foliage while spraying.
- Spray with a slow sweeping motion, covering both upper and lower leaf surfaces and the stern.
- 5. For best results, repeat every 10 14 days.

USE TIPS

- When using this product on plants not listed, test on a small area first.
- ✓ Do not spray if temperature is above 90 ^O F as damage to some delicate plants can occur.
- Spray early in the morning or late in the evening, when air is still.
- ✓ If it rains before product has dried, re-apply.
- Regular light sprayings are more effective than a single heavy treatment.

SENSITIVE PLANT CARE

Use with care on ferns, palms and tuchsia. As with all plants, especially carnations, chrysanthemums, and roses, there exists a wide variety of species, and therefore, a wide variety of susceptibility to injury. To test for plant sensitivity, spray a few plants as described above, wait 24 hours, and check for possible injury.

PRECAUTIONARY STATEMENTS AND STORAGE & DISPOSAL FOR EITHER FORMAT

STORAGE AND DISPOSAL

Store in a cool area away from heat or open flame. Discard container in trash. Do not incinerate or puncture.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domostic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with scop and water after handling.

First Ald

If in Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water.

Physical or Chemical Hazards

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130 °F may cause bursting.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Questions or comments? Call 800-332-5553

Contains NO CFC's or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.

Made in the USA by Chemsico, Division of United Industries Corporation, P.O. Box 15842, St. Louis, MO 63114-0842

EPA Reg. No. 9688-122 EPA Est. No. 9688-MO-1 © 1998