



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

January 12, 2026

Michael Sarli
Sr. Director Regulatory
Chemsico, A Div. of United Industries Corp.
P.O. Box 142642
St. Louis, MO 63114

Subject: Label Amendment - Registration Review Mitigation for lambda-Cyhalothrin
Product Name: Chemsico Insecticide Concentrate 5L
EPA Registration Number: 9688-183
Case Number: 480103
Application Dates: February 17, 2022

Dear Michael Sarli:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the lambda-Cyhalothrin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for

shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Tracy Jackson by phone at 202-566-2268, or via email at jackson.tracy@epa.gov.

Sincerely,



Julie Javier, Team Leader
Risk Mitigation and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

CHEMSICO

Insecticide Concentrate 5L

For Outdoor Use Around the Home Only.

Claims for a General Use Product

[Protects Listed Areas:] [Protects] Vegetables, Fruit and Nut Trees, Roses, Flowers, Trees, Shrubs • Ants*, Aphids, Chinch Bugs, Cutworms, Flea Beetles, Grubs, Mites, Sod Webworms & many others • Kills listed outdoor insects• Kills Garden Insects...including grubs, chinch bugs, European crane flies, fleas, fire ants, mosquitoes, sod webworms, aphids, Japanese beetles, whiteflies, ants*, mites, deer ticks and other listed insects • Kills grubs • Keeps Killing Nuisance Pests Outside the Home for Up to 8 Weeks [2 months] • Keeps Killing Pests on Outdoor Surfaces for Up to 8 Weeks [2 months] • Up to 8 week Control of Nuisance Pests • Low Odor [Formula] • Kills Fast • Fast Kill Formula • Connect and Spray (Optional for hose-end sprayer) • Spray Concentrate • Concentrated Insect Spray • Kills 180+ Pests [on Contact]•

**Guaranteed Results or Your Money Back • Makes up to 128 gallons • Covers up to 16,000 square feet • Long Lasting Formula • Fast Acting Formula • For Broadcast and Barrier Use • ‡Non-staining • Works [Kills] in 24 Hours • Kills Grubs in 24 Hours • Broad Spectrum [Insect Killer] [Insecticide] [Concentrate] • Fast Acting [Long Lasting] Formula • Kills over 180 pests on contact • For Garden Insects • Kills Japanese Beetles Fast • Long Lasting Protection From Insect Damage • Kills Insects • Kills Above & Below Ground • Mix with water to use with tank sprayer, dial-style hose-end sprayer, concentrate plus water hose-end sprayer • Kills Carpenter Ants & other listed ant species • Kills Ants* on Contact • Use on shrubs & other listed areas • *Argentine, Southern, Field, Allegheny mound, Florida carpenter, Cornfield, Honey, Pavement, Nuisance, Black carpenter]

Fire Ant Claims

[Kills Fire Ants in 24 Hours • Kills Fire Ant Mounds in 24 hours • Kills the Queen • Kills the Fire Ant Mound • Kills the Queen and Mound • Controls Fire Ants for up to Four Weeks]

Active Ingredient

Lambda-Cyhalothrin 0.5%

Other Ingredients 99.5%

KEEP OUT OF REACH OF CHILDREN

CAUTION. See end of booklet for additional precautionary statements [Option for booklet label: See back booklet for additional precautionary statements.]

Net [Contents] 1 qt/ 32 fl oz / 946 mL

ACCEPTED

Jan 12, 2026

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under
EPA Reg. No. 9688-183

OPTIONAL PRODUCT FACT BOX

PRODUCT FACT BOX

What it Does:

Kills Grubs, Ants*, Fleas, Chinch Bugs, Mosquitoes, Sod Webworms, Aphids, Beetles, Whiteflies, Mites, Deer Ticks, Japanese Beetles, European Crane Flies, Fire Ants and Other Listed Insects

Where to Use: (icon outdoors)

Questions & Comments: (phone icon) (computer icon)
Call 1-800-332-5553 or Visit our Website at www.xxxx.com

*Argentine, Southern, Field, Allegheny mound, Florida carpenter, Cornfield, Honey, Pavement, Nuisance, Black carpenter.

STOP. READ ENTIRE LABEL BEFORE

USE. DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



Do not allow children or pets into the treated area until dry.

DO NOT apply directly to residential lawns and turf in residential settings (e.g., homes, parks, schools, athletic fields or any other area frequented by the general public).

All outdoor spray applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1. Application to pervious surfaces such as soil, and vegetation;
2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or groundcover that abuts the perimeter of the treated structure);
3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch;
5. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;
6. Applications to vertical surfaces directly above pervious surfaces, such as soil, mulch or vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).

Do not apply the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.

Do not apply or irrigate to the point of runoff.

Do not apply when the wind speed is greater than 15 mph.

For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).

Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

Treat surfaces to ensure thorough coverage but avoid runoff.

To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

In New York State, this product may not be applied within 100 ft of a coastal marsh, or stream that drains directly into a coastal marsh.

HOW TO MIX AND APPLY

STEP 1

Determine size of spraying job. For vegetables, measure the area to be sprayed. [For trees, shrubs and flowers, apply as a thorough cover spray.] Make applications when insects first appear. Applications should be repeated only as directed to maintain control.

STEP 2

Use a clean sprayer. Carefully measure and mix the amount of product and water as indicated in the [rate chart][directions]. It is always a good idea to wear [disposable] gloves when handling pesticides. **Kitchen utensils such as measuring cups and measuring spoons should not be used for food**

purposes after use with pesticides.

STEP 3

Spray as directed [in rate chart]. Thorough coverage is important.

STEP 4

Flush sprayer with clean water after each use. **See [chart][directions] for list of use sites, plants and insects controlled.** Use the rate given in the [chart] [directions] for the type of insect you want to control on the site or plant you are spraying.

MEASUREMENT CONVERSIONS

1 Tablespoon = 3 teaspoons, 1 fl oz = 6 teaspoons or 2 Tablespoons,
6 fl oz = $\frac{3}{4}$ cup

1 $\frac{1}{2}$ Tablespoons in 3 gallons of water = $\frac{1}{2}$ teaspoons in 1 gallon of water

2 Tablespoons in 3 gallons of water = 2 teaspoons in 1 gallon of water

†NOTE: This product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding in particular), test in an inconspicuous area and recheck in a few hours. Do not use if any staining is observed.

[NUISANCE] PESTS OUTSIDE THE HOME

[HOME BARRIER TREATMENT] Mix 0.5 fl oz (1 Tbsp) in 1 gallon of water and apply as a continuous band of insecticide around building foundation and around windows, doors, eaves, vents and other areas to greatly reduce the potential for entry by crawling pests. For easier application, remove debris and leaf litter from next to the foundation, cut back vegetation and branches that touch the foundation, and move or rake back rocks, deep mulch, or other potential pest harborage next to the foundation. Apply a 3 ft band around the structure and upwards along the foundation to 3 feet and around windows, doors, and roof overhangs. Apply as a coarse spray to thoroughly and uniformly wet the foundation and/or band area so that the insecticide will reach the soil or thatch level where pests may be active. One gallon will treat 250 sq ft of surface area. Repeat application every 8 weeks [2 months].

OUTDOOR SURFACES USE: For control of ants*, centipedes, cockroaches, crickets, fleas, millipedes, mosquitoes, palmetto bugs, scorpions, sowbugs, pillbugs, spiders, deer ticks and waterbugs. Mix 0.5 fl oz (1 Tbsp) in 1 gallon of water and apply as a residual treatment to ornamental plants next to foundations of buildings and to surfaces of buildings, porches, screens, window frames, eaves, patios, garages and other similar areas where these pests are active. Repeat application every 8 weeks [2 months].

*Argentine, Southern, Field, Allegheny mound, Florida carpenter, Cornfield, Honey, Pavement, Nuisance, Black carpenter.

FIRE ANT MOUNDS

Mix 1 fl oz (2 Tbsp) in 1 gallon of water and apply to each fire ant mound. Thoroughly wet the mound and surrounding area to a 4 ft diameter. For best results apply in cool weather, 65° to 80°F, or in the early morning or late evening. Kills the entire mound, including the queen, in 24 hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment.

ORNAMENTAL TREES, SHRUBS AND FLOWERS

[For use on ornamental trees, shrubs and flowers such as but not limited to: Ageratum, Ash, Aster, Arborvitae, Arizona Cypress, Azaleas, Begonia, Birch, Boxwood, Box Elder, Camellias, Cherry (ornamental), Non-bearing Citrus, Chrysanthemums, Carnations, Coleus, Common Ninebark, Conifers, Douglas Fir, Elm, English Ivy, Euonymus, Exacum, Weeping Fig, Fir, Gladioli, Gold Bells, Hawthorn, Holly, Honeysuckle, Hypoestes, Ivy, Juniper, Lilac, Locust, Maple, Marigold, Mimosa, Mock Orange, Nannyberry, Oak, Orchid, Palm, Pansy, Pea Shrub, Petunia, Philodendron, Pine, Ornamental Plum, Poinsettia, Poplar, Tulip Poplar, Rhododendron, Roses, Snapdragon, Snowberry, Spruce, Taxus, Willow, Zinnia.]

To control [kill] ants*, armyworms, azalea caterpillar, aphids, bagworms, black vine weevil (adult), box elder bugs, budworms, California oakworm, cankerworms, cockroaches, crickets, cutworms, Eastern tent caterpillar, elm leaf beetles, European sawfly, fall webworm, flea beetles, forest tent caterpillar, gypsy moth larvae, Japanese beetles (adults), June beetles (adults), lace bugs, leaf-feeding caterpillars, leafhoppers, leafminers (adults), leaf rollers, leaf skeletonizers, midges, oleander moth larvae, pillbug, pine sawfly, pine shoot beetles, pinetip moths, plant bugs, root weevil, sawfly, scale insects (crawlers), spiders, spittlebugs, striped beetles, striped oakworm, thrips, tip moth, tussock moth larvae: Mix 0.25 fl oz (1/2 Tbsp) in 1 gallon of water.

To control [kill] broadmites, brown softscale, California redscale (crawler), clover mites, mealybugs, pineneedle scale (crawler), spider mites and whiteflies: Mix 0.5 fl oz (1 Tbsp) in 1 gallon of water.

*Argentine, Southern, Field, Allegheny mound, Florida carpenter, Cornfield, Honey, Pavement, Nuisance, Black carpenter.

USE TIPS

- For best results, apply early before insect populations are high.
- For scale control, treat entire plant, including stem, trunks and twigs.
- For bagworm control, apply when bagworm larvae begin to hatch.
- For best results, apply to both sides of plant leaves and all plantsurfaces where insects are present.

VEGETABLES

Mix 0.5 fl oz (1 Tbsp) in 1 gallon of water and apply to 300 sq ft. Apply when insects appear or when damage occurs. Spray to wet all infested plant surfaces. Do not apply more than 5 times per growing season to corn. For all other crops do not apply more than 9 times per growing season.

[Bracketed Text] = Optional language

For use on Broccoli, Brussels Sprouts, Cabbage, Cavallo Broccoli, Cauliflower, Chinese Broccoli, Chinese Cabbage, Chinese Mustard, and Kohlrabi to control [kill]: **Alfalfa Looper, Cabbage Looper, Imported Cabbageworm, Southern Cabbageworm, Cutworm, Cabbage Webworm, Diamondback moth, Armyworm, Beet Armyworm, Fall Armyworm, Yellow-striped Armyworm, Corn Earworm, Flea Beetle, Japanese Beetle (Adult), Grasshopper, Leafhopper, Plant Bug, including Lygus Bug, Stink Bug, Meadow Spittlebug, Aphid, Whitefly, Thrips, Spider Mite.**

For use on Corn (sweet and pop) to control [kill] **Cutworm, Western Bean Cutworm, Corn Earworm, Green Cloverworm, Meadow Spittlebug, Tobacco Budworm, European Corn Borer, Southwestern Corn Borer, Stalk Borer, Hop Vine Borer, Armyworm, Beet armyworm, Western Corn Rootworm Beetle (adult), Northern Corn Rootworm Beetle (adult), Southern Corn Rootworm Beetle (adult), Mexican Corn Rootworm Beetle (adult), Bean Leaf Beetle, Cereal Leaf Beetle, Japanese Beetle (adult), Sap Beetle (adult), Stink Bug, Grasshopper, Corn Leaf Aphid, English Grain Aphid, Chinch Bug, Green Bug.**

Sweet corn only: Southern Armyworm, Western Bean Cutworm, Tarnished Plant Bug, Aster Leafhopper, Aphid, Spider Mite, Corn silkfly (adult).

For use on Lettuce (head and leaf) to control [kill]: **Alfalfa Looper, Cabbage Looper, Imported Cabbageworm, Cutworm, Saltmarsh Caterpillar, Green Cloverworm, Diamondback moth, Armyworm, Beet Armyworm, Fall Armyworm, Southern Armyworm, Corn Earworm, Tobacco Budworm, European Corn Borer, Flea Beetle, Japanese Beetle (Adult), Vegetable Weevil (adult), Grasshopper, Leafhopper, Plant Bug, Stink Bug, Meadow Spittlebug, Aphid, Whitefly, Spider Mite.**

For use on Onion (bulb) and Garlic to control [kill]: **Cutworm, Seedcorn Maggot (adult), Onion Maggot (adult), Leafminer, (adult), Armyworm, Onion Thrips, Tobacco Thrips, Western Flower Thrips, Flower Thrips, Aphid, Plant Bug, Stink Bug.**

For Use on Tomato, Tomatillo, Peppers (bell and nonbell), Eggplant, Ground Cherry and Pepino to control [kill]: **Cabbage Looper, Cutworm spp., Hornworm, Tomato Fruitworm, Tobacco Budworm, Tomato Pinworm, Beet Armyworm, Southern Armyworm, Yellow-striped Armyworm, Fall Armyworm, European Corn Borer, Leafminer, Colorado Potato Beetle, Flea Beetle, Grasshopper, Leafhopper, Aphid, Whitefly, Meadow Spittlebug, Stink Bug, Plant Bug, Stalk Borer, Blister Beetle, Japanese Beetle (adult), Vegetable Weevil, Cucumber Beetle (adult).**

For use on Asparagus beans, Snap Beans, Wax Beans, Black-eyed Peas, Garden Peas, Green Peas and Sugar Snap Peas to control [kill]: **Cutworm spp., Green Cloverworm, Imported Cabbageworm, Saltmarsh Caterpillar, Velvetleaf Caterpillar, Mexican Bean Beetle, Corn Earworm, Painted Lady Butterfly (Larva), European Corn Borer, Looper spp., Western Bean Cutworm, Armyworm, Yellow-Striped Armyworm, Western Yellow-Striped Armyworm, Bean Leaf skeletonizer, Webworm spp., Leaffier spp., Alfalfa Caterpillar, Stalk Borer, Cucumber Beetle (adult), Corn Rootworm Beetle (adult), Flea Beetle (adult), Blister**

Beetle, Bean Leaf Beetle, Japanese Beetle (adult), Leafhopper spp., Fleas Hopper spp., Three-Cornered Alfalfa Hopper, Meadow Spittle Bug, Stink Bug spp., Plant Bug spp., Grasshopper, Thrips spp., Aphid spp.

NUMBER OF DAYS TO WAIT BETWEEN THE LAST APPLICATION AND HARVEST	
VEGETABLES	PHI
Broccoli, Brussels sprouts, Cabbage, Cavallo Broccoli, Cauliflower, Chinese Broccoli, Chinese Cabbage, Chinese Mustard, Kohlrabi, Lettuce (Head & Leaf)	1
Tomato, Tomatillo, Peppers (bell and nonbell), Eggplant, Ground Cherry and Pepino	5
Onions (bulb), Garlic	14
Corn (sweet and pop)	21
Asparagus beans, Snap Beans, Wax Beans, Black-eyed Peas, Garden Peas, Green Peas and Sugar Snap Peas	7
Lettuce (Leaf and Head)	1
PHI = Pre-Harvest Interval	

FRUIT AND NUT TREES

Mix 0.5 fl oz (1 Tbsp) in 1 gallon of water and apply to 300 sq ft. Apply when insects appear or when damage occurs. Spray to wet all infested plant surfaces.

For Fruit Trees do not apply more than 9 times per year, and only apply 5 times after blooms have formed. For Nut Trees do not apply more than 6 times per year, and only apply 5 times after blooms have formed.

For use on Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear and Quince to control [kill]: **Leafhopper spp., Codling Moth, Tufted Apple Budworm, Oriental Fruit Moth, Lesser Appleworm, Green Fruitworm, Tent Caterpillar spp., Tentiform Leaf Miner spp., Apple Maggot (adult), Pear Sawfly, Plum Curculio, Japanese Beetle, Plant Bug, Stink Bug, Leafhopper, Periodical Cicada, Apple Aphid, Rosy Apple Aphid, Pear Psylla, San Jose Scale (on fruit only).**

For use on Apricot, Sweet Cherry, Tart Cherry, Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot and Prune to control [kill]: **Leafroller spp., Peach Twig Borer, Oriental Fruit Moth, Peachtree Borer spp., Green Fruitworm, Tent Caterpillar spp., American Plum Borer, Cherry Fruit Fly spp., (adult), Plum Curculio, Rose Chafer, Japanese Beetle, Plant Bug, Stink Bug, Leafhopper spp., Periodical Cicada, Black Cherry Aphid.**

For use on Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut, Walnut (Black and English) and Pecan to control [kill]: **Leafroller spp., Navel Orangeworm, Codling moth, Filbertworm, Peach Twig Borer, Walnut Husk Fly spp. (adult), Ants, Plant Bug, Stink Bug, Chinch Bug, Leaffooted Bug, Walnut Aphid, Hickory Shuckworm, Pecan Casebearer spp., Pecan Weevil, Pecan Aphid spp., Pecan Spittlebug, Pecan Phylloxera spp.**

[Bracketed Text] = Optional language

NUMBER OF DAYS TO WAIT BETWEEN THE LAST APPLICATION AND HARVEST	
Fruit and Nut Trees	PHI
Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear, Quince	21
Apricot, Sweet Cherry, Tart Cherry, Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot and Prune	14
Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut, Walnut (Black and English) and Pecan	14
PHI = Pre-Harvest Interval	

STORAGE AND DISPOSAL

Storage: Store in a cool dry area away from heat or open flame.

If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling, if available. **If partly filled:** Do not pour or dispose down-the-drain or sewer. Call your local solid waste agency for disposal instructions.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID

If Swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person. **If in Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor for treatment advice.

Environmental Hazards

This product is extremely toxic to fish and other aquatic organisms. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainagesystems.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protect pollinating insects by following label directions intended to minimize drift and reduce pesticide risk to these organisms.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

[Questions or comments? Call 1-800-332-5553 or visit our website.]

[**If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.]

Made in the USA for
Chemsico, A Division of United Industries Corporation
PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No.: 9688-183
EPA Est. No.: 9688-MO-1

© 2025

OPTIONAL READY SPRAY NOZZLE DIRECTIONS

STOP. READ ENTIRE LABEL BEFORE USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



Do not allow children or pets into the treated area until dry.

DO NOT apply directly to residential lawns and turf in residential settings (e.g., homes, parks, schools, athletic fields or any other area frequented by the general public).

All outdoor spray applications must be limited to spot or crack-and- crevice treatments only, except for the following permitted uses:

1. Application to pervious surfaces such as soil, and vegetation;
2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or groundcover that abuts the perimeter of the treated structure);
3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch;
5. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;
6. Applications to vertical surfaces directly above pervious surfaces, such as soil, mulch or vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surfacewaters.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).

Do not apply the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.

Do not apply or irrigate to the point of runoff.

Do not apply when the wind speed is greater than 15 mph.

For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).

Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

Treat surfaces to ensure thorough coverage but avoid runoff.

To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

In New York State, this product may not be applied within 100 ft of a coastal marsh, or stream that drains directly into a coastal marsh.

HOW TO APPLY

For trees, shrubs and flowers, apply as a thorough cover spray. Begin treating when insects first appear. Applications should be repeated only as directed to maintain control. Apply as directed [in the rate chart]. Remember, thorough coverage is important. Be sure to follow the important use limitations for the specific vegetable you are treating, including the amount of spray that can be applied, the interval between treatments and the days to wait before harvest.

USING THE READY SPRAY NOZZLE (Optional subheading)

1. *Shake container well before using.*
2. Connect a garden hose to the Ready Spray nozzle. Make sure the dial on the nozzle is in the "OFF" position with the [green] safety tab in the valve notch.
3. Turn on water at faucet. Extend hose to the farthest area to be treated and work back toward the faucet so you don't come in contact with the treated area.
4. To BEGIN spraying, point nozzle toward treatment site and a.) bend the [green] safety tab back (located at right of yellow dial) with your thumb, b.) hold while turning yellow dial clockwise until it stops. Water will automatically mix with the product.
5. Spray until wet to control insects. Walk at a steady pace while spraying using an even sweeping motion, slightly overlapping treated areas.

[Bracketed Text] = Optional language

6. To STOP spraying, QUICKLY turn the dial counterclockwise until it stops and the safety tab engages into the notch on the valve. Turn water off at faucet. To relieve pressure before removing nozzle from hose, bend the safety tab back and turn dial "ON" until water stops spraying.
7. To STORE unused product, make sure the dial is in the "OFF" position with [green] safety tab in the valve notch. Place in cool area away from heat, sunlight or open flame.

‡NOTE: This product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding in particular), test in an inconspicuous area and recheck in a few hours. Do not use any if any staining is observed.

ORNAMENTAL TREES, SHRUBS AND FLOWERS

[For use on ornamental trees, shrubs and flowers such as but not limited to: Ageratum, Aster, Arborvitae, Arizona Cypress, Azaleas, Begonia, Birch, Boxwood, Box Elder, Camellias, Carnations, Cherry (ornamental), Chrysanthemums, Carnations, Coleus, Common Ninebark, Conifers, Douglas Fir, Elm, English Ivy, Euonymus, Exacum, Weeping Fig, Fir, Gladioli, Gold Bells, Hawthorn, Holly, Honeysuckle, Hypoestes, Ivy, Juniper, Lilac, Locust, Maple, Marigold, Mimosa, Mock Orange, Nannyberry, Oak, Orchid, Palm, Pansy, Pea Shrub, Petunia, Philodendron, Pine, Ornamental Plum, Poinsettia, Poplar, Tulip Poplar, Rhododendron, Roses, Snapdragon, Snowberry, Spruce, Taxus, Weeping Fig, Willow, Zinnia.]

To control [kill] ants*, armyworms, azalea caterpillar, aphids, bagworms, black vine weevil (adult), boxelder bugs, budworms, California oakworm, cankerworms, cockroaches, crickets, cutworms, Eastern tent caterpillar, elm leaf beetles, European sawfly, fall webworm, flea beetles, forest tent caterpillar, gypsy moth larvae, Japanese beetles (adults), June beetles (adults), lace bugs, leaf-feeding caterpillars, leafhoppers, leafminers (adults), leafrollers, leaf skeletonizers, midges, oleander moth larvae, pillbug, pine sawfly, pine shoot beetles, pinetip moths, plant bugs, root weevil, sawfly, scale insects (crawlers), spiders, spittlebugs, striped beetles, striped oakworm, thrips, tip moth, tussock moth larvae, broadmites, brown softscale, California redscale (crawler), clover mites, mealybugs, pine needle scale (crawler), spider mites and whiteflies.

*Argentine, Southern, Field, Allegheny mound, Florida carpenter, Cornfield, Honey, Pavement, Nuisance, Black carpenter.

Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to spray underside of leaves and penetrate dense foliage.

USE TIPS

- For best results, apply early before insect populations are high.
- For scale control, treat entire plant, including stem, trunks and twigs.
- For bagworm control, apply when bagworm larvae begin to hatch.
- For best results, apply to both sides of plant leaves and all plantsurfaces

where insects are found.

VEGETABLES

Apply when insects appear or when damage occurs. Spray to wet all infested plant surfaces. Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to spray underside of leaves and penetrate dense foliage. One qt will treat 19,200 sq ft of vegetable plants.

Do not apply more than 5 times per growing season to corn. For all other crops do not apply more than 9 times per growing season.

For use on Broccoli, Brussels Sprouts, Cabbage, Cavallo Broccoli, Cauliflower, Chinese Broccoli, Chinese Cabbage, Chinese Mustard, and Kohlrabi to control [kill]: **Alfalfa Looper, Cabbage Looper, Imported Cabbageworm, Southern Cabbageworm, Cutworm, Cabbage Webworm, Diamondback moth, Armyworm, Beet Armyworm, Fall Armyworm, Yellow-striped Armyworm, Corn Earworm, Flea Beetle, Japanese Beetle (Adult), Grasshopper, Leafhopper, Plant Bug, including Lygus Bug, Stink Bug, Meadow Spittlebug, Aphid, Whitefly, Thrips, Spider Mite.**

For use on Corn (sweet and pop) to control [kill] **Cutworm, Western Bean Cutworm, Corn Earworm, Green Cloverworm, Meadow Spittlebug, Tobacco Budworm, European Corn Borer, Southwestern Corn Borer, Stalk Borer, Hop Vine Borer, Armyworm, Beet armyworm, Western Corn Rootworm Beetle (adult), Northern Corn Rootworm Beetle (adult), Southern Corn Rootworm Beetle (adult), Mexican Corn Rootworm Beetle (adult), Bean Leaf Beetle, Cereal Leaf Beetle, Japanese Beetle (adult), Sap Beetle (adult), Stink Bug, Grasshopper, Corn Leaf Aphid, English Grain Aphid, Chinch Bug, Green Bug.**

Sweet corn only: Southern Armyworm, Western Bean Cutworm, Tarnished Plant Bug, Aster Leafhopper, Aphid, Spider Mite, Corn silkfly (adult).

For use on Lettuce (head and leaf) to control [kill]: **Alfalfa Looper, Cabbage Looper, Imported Cabbageworm, Cutworm, Saltmarsh Caterpillar, Green Cloverworm, Diamondback moth, Armyworm, Beet Armyworm, Fall Armyworm, Southern Armyworm, Corn Earworm, Tobacco Budworm, European Corn Borer, Flea Beetle, Japanese Beetle (Adult), Vegetable Weevil (adult), Grasshopper, Leafhopper, Plant Bug, Stink Bug, Meadow Spittlebug, Aphid, Whitefly, Spider Mite.**

For use on Onion (bulb) and Garlic to control [kill]: **Cutworm, Seedcorn Maggot (adult), Onion Maggot (adult), Leafminer, (adult), Armyworm, Onion Thrips, Tobacco Thrips, Western Flower Thrips, Flower Thrips, Aphid, Plant Bug, Stink Bug.**

For Use on Tomato, Tomatillo, Peppers (bell and nonbell), Eggplant, Ground Cherry and Pepino to control [kill]: **Cabbage Looper, Cutworm spp., Hornworm, Tomato Fruitworm, Tobacco Budworm, Tomato Pinworm, Beet Armyworm, Southern Armyworm, Yellow-striped Armyworm, Fall Armyworm, European Corn Borer, Leafminer, Colorado Potato Beetle, Flea Beetle, Grasshopper, Leafhopper, Aphid, Whitefly, Meadow Spittlebug, Stink Bug, Plant Bug, Stalk**

Borer, Blister Beetle, Japanese Beetle (adult), Vegetable Weevil, Cucumber Beetle (adult).

For use on Asparagus beans, Snap Beans, Wax Beans Black-eyed Peas, Garden Peas, Green Peas and Sugar Snap Peas to control [kill]: **Cutworm spp., Green Cloverworm, Imported Cabbageworm, Saltmarsh Caterpillar, Velvetleaf Caterpillar, Mexican Bean Beetle, Corn Earworm, Painted Lady Butterfly (Larva), European Corn Borer, Looper spp., Western Bean Cutworm, Armyworm, Yellow-Striped Armyworm, Western Yellow-Striped Armyworm, Bean Leaf skeletonizer, Webworm spp., Leaftier spp., Alfalfa Caterpillar, Stalk Borer, Cucumber Beetle (adult), Corn Rootworm Beetle (adult), Flea Beetle (adult), Blister Beetle, Bean Leaf Beetle, Japanese Beetle (adult), Leafhopper spp., Fleas Hopper spp., Three-Cornered Alfalfa Hopper, Meadow Spittle Bug, Stink Bug spp., Plant Bug spp., Grasshopper, Thrips spp., Aphid spp.**

For use on Lettuce (Head and Leaf) to control [kill]: **Alfalfa Looper, Cabbage Looper, Imported Cabbageworm, Cutworm spp., Saltmarsh Caterpillar, Green Cloverworm, Armyworm, Beet Armyworm, Fall Armyworm, Corn Earworm, European Corn Borer, Flea Beetle spp., Japanese Beetle (adult), Vegetable Beetle (adult), Grasshopper, Leafhopper, Plant Bug, Stink Bug, Meadow Spittlebug.**

NUMBER OF DAYS TO WAIT BETWEEN THE LAST APPLICATION AND HARVEST	
VEGETABLES	PHI
Broccoli, Brussels sprouts, Cabbage, Cavallo Broccoli, Cauliflower, Chinese Broccoli, Chinese Cabbage, Chinese Mustard, Kohlrabi, Lettuce (Head & Leaf)	1
Tomato, Tomatillo, Peppers (bell and nonbell), Eggplant, Ground Cherry and Pepino	5
Onions (bulb), Garlic	14
Corn (sweet and pop)	21
Asparagus beans, Snap Beans, Wax Beans Black-eyed Peas, Garden Peas, Green Peas and Sugar Snap Peas	7
Lettuce (Leaf and Head)	1
PHI = Pre-Harvest Interval	

FRUIT AND NUT TREES

Apply when insects appear or when damage occurs. Spray to wet all infested plant surfaces. Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to spray underside of leaves and penetrate dense foliage. One qt will treat 19,200 sq ft of fruit and nut trees.

For Fruit Trees do not apply more than 9 times per year, and only apply 5 times after blooms have formed. For Nut Trees do not apply more than 6 times per year, and only apply 5 times after blooms have formed.

For use on Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear and Quince to control [kill]: Leafhopper spp., Codling Moth, Tufted Apple Budworm, Oriental Fruit Moth, Lesser Appleworm, Green Fruitworm, Tent Caterpillar spp., Tentiform Leaf Miner spp., Apple Maggot (adult), Pear Sawfly, Plum Curculio, Japanese Beetle, Plant Bug, Stink Bug, Leafhopper, Periodical Cicada, Apple Aphid, Rosy Apple Aphid, Pear Psylla, San Jose Scale (on fruit only).

For use on Apricot, Sweet Cherry, Tart Cherry, Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot and Prune to control [kill]: Leafroller spp., Peach Twig Borer, Oriental Fruit Moth, Peachtree Borer spp., Green Fruitworm, Tent Caterpillar spp., American Plum Borer, Cherry Fruit Fly spp., (adult), Plum Curculio, Rose Chafer, Japanese Beetle, Plant Bug, Stink Bug, Leafhopper spp., Periodical Cicada, Black Cherry Aphid.

For use on Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut, Walnut (Black and English) and Pecan to control [kill]: Leafroller spp., Navel Orangeworm, Codling moth, Filbertworm, Peach Twig Borer, Walnut Husk Fly spp. (adult), Ants, Plant Bug, Stink Bug, Chinch Bug, Leaffooted Bug, Walnut Aphid, Hickory Shuckworm, Pecan Casebearer spp., Pecan Weevil, Pecan Aphid spp., Pecan Spittlebug, Pecan Phylloxera spp.

NUMBER OF DAYS TO WAIT BETWEEN THE LAST APPLICATION AND HARVEST	
Fruit and Nut Trees	PHI
Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear, Quince	21
Apricot, Sweet Cherry, Tart Cherry, Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot and Prune	14
Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut, Walnut (Black and English) and Pecan	14
PHI = Pre-Harvest Interval	

[Bracketed Text] = Optional language

PRECAUTIONARY STATEMENTS**Hazards to Humans and Domestic Animals**

CAUTION. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID

If Swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person. **If in Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor for treatment advice.

Environmental Hazards

This product is extremely toxic to fish and other aquatic organisms. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Protect pollinating insects by following label directions intended to minimize drift and reduce pesticide risk to these organisms.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

[Questions or comments? Call 1-800-332-5553 or visit our website.]

[**If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.]

Chemsico, A Division of United Industries Corporation
PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No.: 9688-183

EPA Est. No.: 9688-MO-1

© 2025

OPTIONAL DIRECTIONS FOR OUTDOOR SURFACE USE ONLY

OPTIONAL PRODUCT FACTS BOX

PRODUCT FACT BOX

What it Does: Kills Grubs, Ants*, Fleas, Chinch Bugs, Mosquitoes, Sod Webworms, Aphids, Beetles, Whiteflies, Mites, Deer Ticks, Japanese Beetles, European Crane Flies, Fire Ants and Other Listed Insects

Where to Use: Outdoor Surfaces

How Much to Use: 0.5 fl oz per gallon of water

Coverage (32 oz): Up to 16,000 sq ft

Questions & Comments: (phone icon) (computer icon)

Call 1-800-332-5553 or Visit our Website at www.xxxx.com

*Argentine, Southern, Field, Allegheny mound, Florida carpenter, Cornfield, Honey, Pavement, Nuisance, Black carpenter.

STOP. READ ENTIRE LABEL BEFORE USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



Do not allow children or pets into the treated area until dry.

DO NOT apply directly to residential lawns and turf in residential settings (e.g., homes, parks, schools, athletic fields or any other area frequented by the general public).

All outdoor spray applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1. Application to pervious surfaces such as soil, and vegetation;
2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or groundcover that abuts the perimeter of the treated structure);
3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch;
5. Applications to vertical surfaces (such as the side of a man-made structure)

[Bracketed Text] = Optional language

directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;

6. Applications to vertical surfaces directly above pervious surfaces, such as soil, mulch or vegetation only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).

Do not apply the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.

Do not apply or irrigate to the point of runoff.

Do not apply when the wind speed is greater than 15 mph.

For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).

Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

Treat surfaces to ensure thorough coverage but avoid runoff.

To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

In New York State, this product may not be applied within 100 ft of a coastal marsh, or stream that drains directly into a coastal marsh.

HOW TO MIX AND APPLY

STEP 1

Determine size of spraying job. For vegetables, measure the area to be

sprayed. [For trees, shrubs and flowers, apply as a thorough cover spray.] Make applications when insects first appear. Applications should be repeated only as directed to maintain control.

STEP 2

Use a clean sprayer. Carefully measure and mix the amount of product and water as indicated in the [rate chart][directions]. It is always a good idea to wear [disposable] gloves when handling pesticides. **Kitchen utensils such as**

measuring cups and measuring spoons should not be used for food purposes after use with pesticides.

STEP 3

Spray as directed [in rate chart]. Thorough coverage is important.

STEP 4

Flush sprayer with clean water after each use.

[Optional Sprayer Information]**[For Tank Sprayers]**

Dilution Rate: 0.5 fl. oz. per gallon

For Dial-Style Hose-End Sprayers

Dilution Rate: 0.5 fl. oz. per gallon

Step 1 Fill sprayer jar with enough concentrate to cover area to be treated.

Step 2 Attach sprayer top and set dial to appropriate dilution rate.

Step 3 Attach Sprayer to hose and apply.

Step 4 Begin by spraying area farthest from faucet, walking backwards as you apply to avoid contact with the treated area.

For Concentrate Plus Water Hose-End Sprayers

Dilution Rate: 0.5 fl. oz. per gallon

Step 1 Pour concentrate into sprayer jar to the required fluid ounce level.

Step 2 Add the appropriate amount of water to the gallons level.

Step 3 Replace sprayer top and shake well before applying.

Step 4 Begin by spraying area farthest from faucet, walking backwards as you apply to avoid contact with the treated area.]

See [chart][directions] for list of use sites, plants and insects controlled. Use the rate given in the [chart] [directions] for the type of insect you want to control on the site or plant you are spraying.

MEASUREMENT CONVERSIONS

1 Tablespoon = 3 teaspoons, 1 fl oz = 6 teaspoons or 2 Tablespoons,

6 fl oz = $\frac{3}{4}$ cup

1 $\frac{1}{2}$ Tablespoons in 3 gallons of water = 1 $\frac{1}{2}$ teaspoons in 1 gallon of water

2 Tablespoons in 3 gallons of water = 2 teaspoons in 1 gallon of water

†NOTE: This product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home

siding (vinyl siding in particular), test in an inconspicuous area and recheck in a few hours. Do not use if any staining is observed.

[NUISANCE] PESTS OUTSIDE THE HOME

OUTDOOR SURFACES USE: For control of ants*, centipedes, cockroaches, crickets, fleas, millipedes, mosquitoes, palmetto bugs, scorpions, sowbugs, pillbugs, spiders, deer ticks and waterbugs. Mix 0.5 fl oz (1 Tbsp) in 1 gallon of water and apply as a residual treatment to ornamental plants next to foundations of buildings and to surfaces of buildings, porches, screens, window frames, eaves, patios, garages and other similar areas where these pests are active. Repeat application every 8 weeks [2 months].

*Argentine, Southern, Field, Allegheny mound, Florida carpenter, Cornfield, Honey, Pavement, Nuisance, Black carpenter.

STORAGE AND DISPOSAL

Storage: Store in a cool dry area away from heat or open flame.

If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling, if available. **If partly filled:** Do not pour or dispose down-the-drain or sewer. Call your local solid waste agency for disposal instructions.

PRECAUTIONARY STATEMENTS**Hazards to Humans and Domestic Animals**

CAUTION. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID

If Swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person. **If in Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor for treatment advice.

Environmental Hazards

This product is extremely toxic to fish and other aquatic organisms. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will

help avoid run off to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Protect pollinating insects by following label directions intended to minimize drift and reduce pesticide risk to these organisms.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

[Questions or comments? Call 1-800-332-5553 or visit our website.]

[**If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.]

Chemsico, A Division of United Industries Corporation
PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No.: 9688-183

EPA Est. No.: 9688-MO-1

© 2025

OPTIONAL DIRECTIONS FOR A FIRE ANT MOUND CONTROL READY TO SPRAY PRODUCT

OPTIONAL PRODUCT FACTS BOX

PRODUCT FACT BOX

What it Does: Kills fire ant queen and colony in 24 hours.

Where to Use: Outdoors, on fire and mounds

How Much to Use: 30 second application

Coverage (32 oz): 32 fire ant mounds

Questions & Comments: (phone icon) (computer icon)

Call 1-800-332-5553 or Visit our Website at www.xxxx.com

Optional Marketing Claims

[Treats 32 mounds • Fast Acting [Fire Ant Killer] Solution • Quick Action Formula [Against Fire Ants] • Kills Fire Ants By Contact • Easy to Use [Fire Ant Control] Formula • No Hassle Fire Ant Killer [Control] Solution • Penetrates Different Soil Types • Penetrates the Mound • Does not Harm Vegetation [or Surrounding Grass]
• Deep Penetrating Formula • Kills the Queen and [Mound][Colony] in 24 Hours • Kills the Queen and Colony • One Application Kills the [Fire Ant] Mound • One Application Kills the Queen and [Colony][Mound]

STOP. READ ENTIRE LABEL BEFORE USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



Do not allow children or pets into the treated area until dry.

DO NOT apply directly to residential lawns and turf in residential settings (e.g., homes, parks, schools, athletic fields or any other area frequented by the general public).

All outdoor spray applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1. Application to pervious surfaces such as soil, and vegetation;
2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or groundcover that abuts the perimeter of the treated structure);
3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch;

5. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;
6. Applications to vertical surfaces directly above pervious surfaces, such as soil mulch, or vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).

Do not apply the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.

Do not apply or irrigate to the point of runoff.

Do not apply when the wind speed is greater than 15 mph.

For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).

Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

Treat surfaces to ensure thorough coverage but avoid runoff.

To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

In New York State, this product may not be applied within 100 ft of a coastal marsh, or stream that drains directly into a coastal marsh.

HOW TO APPLY

1. Make sure control [lever][dial][valve] is in the **OFF** position. Shake well and attach to hose.
2. Turn water on at faucet, aim nozzle over mound [push down tab] and turn the control [lever][dial][valve] to **[ON][Spray]** to begin application.
3. Gently soak mound, moving the applicator to fully wet the mound for 30 seconds (approximately one gallon of spray). Turn control [lever][dial][valve] to **OFF**.
4. Repeat for each mound to be treated.
5. When finished turn control [lever][dial][valve] to **OFF**. Turn faucet **OFF**. Turn control [lever][dial][valve] **ON** and **OFF** again to relieve pressure in hose before removing bottle from hose.

STORAGE AND DISPOSAL

Storage: Store in a cool dry area away from heat or open flame.

If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling, if available. **If partly filled:** Do not pour or dispose down-the-drain or sewer. Call your local solid waste agency for disposal instructions.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID

If Swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person. **If in Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor for treatment advice.

Environmental Hazards

This product is extremely toxic to fish and other aquatic organisms. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Protect pollinating insects by following label directions intended to minimize drift and reduce pesticide risk to these organisms.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

[Questions or comments? Call 1-800-332-5553 or visit our website.]

[**If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.]

Chemsico, A Division of United Industries Corporation
PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No.: 9688-183 EPA
Est. No.: 9688-MO-1
© 2025