

## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Ms. Vee Lee Product Registration CHEMSICO Div. of United Industries Corp. PO Box 142642 St. Louis, MO 63114

MAY 8 2009

SUBJECT:

Application for Pesticide Notification (PRN 98-10) Request General Label Change (Marketing Claims)

EPA Reg. No. 9688-237

Application Dated April 16', 2009

#### Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 04/16/09 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

Please read instructions on reverse before completing form. Form Approved. OMB No. 2070-0060. Approval expires 11-30-93 United States Environmental Protection Agency **OPP Identifier Number** Office of Pesticide Programs (H7505C) Registration **\$EPA** Amendment Washington, DC 20460 Other Application for Pesticide: Section I 1. Company/Product Number 2.Product Manager 3. Proposed Classification 9688-237 George LaRocca 4. Company/Product (Name) PM# Restricted x None **Chemsico Insecticide Concentrate 25G** 13 5. Name and Address of Applicant (include ZIP Code) 6 Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or iden a life to be in the labelling Chemsico, Div. Of United Industries P.O. Box 142642 MAY 08 2009 St. Louis, MO 63114 EPA Reg. No. Check if this is a new address **Product Name** Section II Amendment - Explain below Final printed label in response to Agency letter date "ME TOO" Application. Resubmission in response to Agency letter dated Notification - Explain below. Other - explain below **Explanation:** Use additional page(s) if necessary. (For section I and Section II.) Notification per PR Notice 98-10 to add the following optional marketing claim(s): For Lawns & Landscapes Section III 1. Material in Which This Product Will Be Packaged: Child-Resistant Packaging Unit Packaging Water-Soluble Packaging 2. Type of Container Yes ' Yes Yes Metal No No No Plastic Glass If "YES." If "YES," \* Certification must be No. per Paper No. per submitted container Other Unit package wgt Package wgt container 3. Location of Net Contents Information 4. Size(s) of Retail Container Location of Label Directions, On Laber 2 Label Container On Labeling accompanying product 0.0 Other 6. Manner in Which Label Is Affixed To Product Lithographed Paper glued 10000 Stenciled ( ( ( ( ( ( Section IV 1. Contact Point (Complete items directly below for identification of the individual to be contacted, if necessary, to process this application) Name Telephone No. (include Area Code) **Regulatory Affairs Specialist** 314-683-2438 Vee Lee Certification Date Application I certify that the statements I have made on this form and all attachments thereto are true, accurate, and complete. Received

EPA Form 8570-1 (Rev.12-90)

2. Signature

4. Typed Name

Vee Lee

both under applicable law.

Previous editions are obsolete

I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or

3. Title

5. Date

April 16, 2009

White - EPA File Copy (original)

**Regulatory Affairs Specialist** 

Yellow - Applicant copy

(Stamped)

# Chemsico

Division of United Industries P.O. Box 142642 St. Louis, MO 63114-0642

April 16, 2009

Document Processing Desk (NOTIF)
Office of Pesticide Programs
United States Environmental Protection Agency
Room S4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Attention:

Mr. George LaRocca

Product # 13

Subject:

Notification per PR Notice 98-10

Chemsico Insecticide Concentrate 25G

EPA Reg. No. 9688-237

Dear Mr. Hebert:

Please find the enclosed notification per PR Notice 98-10 to add the optional marketing claim, For Lawns & Landscapes. This is a variation of the already-approved optional marketing claims and language appearing on the label.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

The following documents are enclosed in support of this submission:

- 1. Application for Pesticide (EPA Form 8570-1)
- 2. One copy of the proposed label with changes highlighted
- 3. Five copies of the proposed label

Please call me at 1-800-242-1166, Ext. 2438 you have questions or need further information. Thank you were much for your attention to our application.

Sincerely,

Vee Lee

Regulatory Specialist

# Chemsico Insecticide Concentrate 25G

For Outdoor Use Around the Home Only

#### Claims for a General Use Product

[Protects Listed Areas:] [Protects] Lawns, Vegetables, Fruit and Nut Trees, Roses, Flowers, Trees, Shrubs • Wide Range of Control: [Controls] Ants, Aphids, Chinch Bugs, Cutworms, Flea Beetles, Grubs, Mites, Sod Webworms & many others . Controls tough outdoor insects • Kills Lawn and Garden Insects...including grubs, chinch bugs. European crane flies, fleas, fire ants, mosquitoes, sod webworms, aphids, Japanese beetles, whiteflies, ants, mites, ticks [Deer, Lone star, Brown dog, American dog] and other listed insects • Kills grubs • Keeps Killing Nuisance Pests Outside the Home for Up to 8 Weeks [2 months] • Keeps Killing Pests on Outdoor Surfaces for Up to 8 Weeks [2 months] • Up to 8 week [2 month] Control of Nuisance Pests • Low Odor [Formula] • Kills Fast • Fast Kill Formula • Connect and Spray (Optional for hose-end sprayer) • Spray Concentrate • Concentrated Insect Spray • Kills 180+ Pests [on Contact] • \*Guaranteed Results or Your Money Back • Makes up to 128 gallons • Covers up to 16,000 square feet of lawn ● Water-based ● Long Lasting Formula ● Fast Acting Formula • Soil and Turf Insecticide • For Soil and Turf Insect Control • For Broadcast and Barrier Use • ‡Non-staining • Works [Kills] in 24 Hours • Kills Grubs in 24 Hours • Broad Spectrum [Insect Killer] [Insecticide] [Concentrate] • Fast Acting [Long Lasting] Formula • Won't [Harm] [Kill] Earthworms • Kills over 130 pests on contact • For Lawn [Garden] Insects • Kills Japanese Beetles Fast • Long Lasting Protection From Insect Damage • Kills and Repels Insects • Kills 180+ Insects [on Contact] • Kills Above & Below Ground] • For Lawns & Landscapes

#### Fire Ant Claims

[Kills Fire Ants in 24 Hours ● Kills Fire Ant Mounds in 24 hours ● Kills the Queen ● Kills the Fire Ant Mound ● Kills the Queen and Mound ● Controls Fire Ants for up to Four Weeks]

Active Ingredient	2	:		_	=	υž	
Gamma-Cyhalothrin							. 0.25%
Other Ingredients							

#### KEEP OUT OF REACH OF CHILDREN

CAUTION See end of booklet for additional precautionary statements, [Option for booklet label: See back booklet for additional precautionary statements].

NET CONTENTS 1 QUART/ 32 fl oz / 946 mL

#### OPTIONAL PRODUCT FACT BOX

#### PRODUCT FACT BOX

#### What it Does:

Kills Grubs, Ants, Fleas, Chinch Bugs, Mosquitoes, Sod Webworms, Aphids, Beetles, Whiteflies, Mites, Deer Ticks, Japanese Beetles, European Crane Flies. Fire Ants and Other Listed Insects

Where to Use: (icon outdoors)

**Questions & Comments:** (phone icon) (computer icon) Call 1-800-332-5553 or Visit our Website at <a href="https://www.xxxx.com">www.xxxx.com</a>

#### STOP. READ ENTIRE LABEL BEFORE USE.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



Do not allow children or pets into the treated area until dry.

In New York State, this product may not be applied to lawns within 100 ft of a coastal marsh, or stream that drains directly into a coastal marsh.

### HOW TOM APPLY

STEP 1 Determine are of spraying job. For lawns and vegetables, measure the area to be sprayed. For trees, shrubs and flowers, apply as a thorough cover spray. Make applications when insects first appear. Applications should be repeated only asstirected to maintain control.

STEP 3

Spray as directed [in rate chart]. Thorough coverage is important.



STEP 4

Flush sprayer with clean water after each use.

See [chart][directions] for list of use sites, plants and insects controlled. Use the rate given in the [chart][directions] for the type of insect you want to control on the site or plant you are spraying.

#### **MEASUREMENT CONVERSIONS**

1 Tablespoon = 3 teaspoons, 1 fl. oz. = 6 teaspoons or 2 tablespoons,

6 fl. oz. =  $\frac{3}{4}$  cup

1 ½ Tablespoons in 3 gallons of water = 1½ teaspoons in 1 gallon of water

2 Tablespoons in 3 gallons of water = 2 teaspoons in 1 gallon of water

‡NOTE: This product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding in particular), test in an inconspicuous area and recheck in a few hours. Do not use if any staining is observed.

#### **LAWNS**

UNDERGROUND INSECTS/THATCH INFESTING INSECTS including Blue grass billbug (adult), Black turfgrass Ataenius (adult), chiggers, chinch bugs, white grubs (Japanese beetle, European chafer, Southern chafer), dung beetle and hyperodes weevils (adult), European crane flies, and mole crickets (nymphs and young adults) Mix 0.5 fl.oz. (1 Tablespoon) in 1 gallon of water to treat 125 sq.ft. or 4 fl.oz. in 8 gallons water to treat 1000 sq.ft. For sub-surface insect control (mole crickets, grubs) water treated area with additional ¼ to ½ inch of water for optimum results. [1 quart of Chemsico Insecticide Concentrate 25G treats 8,000 sq. ft.] Do not apply more than seven (7) times per year per location.

SURFACE INSECTS including ants\*, armyworms, cockroaches, crickets, cutworms, earwigs, fleas, lawn moths (sod webworms), millipedes, mosquitoes, palmetto bugs, sowbugs, mites, spittlebugs, and waterbugs. Thoroughly wet down grass a few hours before applying. Mix 0.5 fl.oz. (1 Tablespoon) in 1 gallon of water to cover [treat] 250 sq. ft. or 2 fl.oz. (4 Tablespoons) in 4 gallons of water to cover [treat] 1000 sq.ft. of lawn. Delay additional watering or mowing for 24 hours after application for optimum control of surface insects. For heavy infestations or to kill Lone sianticks, Brown dog ticks, American dog ticks and deer ticks (which may transmit syme Disease) use rate can be increased to 3 fl.oz./1000 sq.ft. [1 quarte of Chemsico Insecticide Concentrate 25G treats 16,000 sq. ft.] Do not apply more than 14 (fourteen) times per year per location.

WHEN TO TREAT: Treat the following pests during the minimal indicated or when first noticed

JAN	FEB	MAR	APR	MAY	JUN	\ <u>````</u> L	AUG	SEP°	, OCT	NOV₀	DEC
				Billbugs		100	Gr. Gr	ubs	\$* 36° €		
			, Dun	g beetle gru	ıbs 🛴						

	Hyperodes wee	vils		
Treat ants	and other insects v	vhen they first a	ppear or you	notice lawn damage

<sup>\*</sup>Argentine, Southern, Field, Allegheny mound, Florida carpenter, Cornfield, Honey, Pavement, Nuisance.

#### [NUSIANCE] PESTS OUTSIDE THE HOME

[HOME BARRIER TREATMENT] Mix 0.5 fl.oz. (1 Tablespoon) in 1 gallon of water and apply as a continuous band of insecticide around building foundation and around windows, doors, eaves, vents and other areas to greatly reduce the potential for entry by crawling pests. For easier application, remove debris and leaf litter from next to the foundation, cut back vegetation and branches that touch the foundation, and move or rake back rocks, deep mulch, or other potential pest harborage next to the foundation. Apply a 3 ft. band around the structure and upwards along the foundation to 3 feet and around windows, doors, and roof overhangs. Apply as a coarse spray to thoroughly and uniformly wet the foundation and/or band area so that the insecticide will reach the soil or thatch level where pests may be active. 1 gallon will treat 250 sq. ft. of surface area. Repeat application every 8 weeks [2 months].

**OUTDOOR SURFACES USE:** For control of ants\*, centipedes, cockroaches, crickets, fleas, millipedes, mosquitoes, palmetto bugs, scorpions, sowbugs, pillbugs, spiders, deer ticks and waterbugs. Mix 0.5 fl.oz. (1 Tablespoon) in one gallon of water and apply as a residual treatment to ornamental plants next to foundations of buildings and to surfaces of buildings, porches, screens, window frames, eaves, patios, garages and other similar areas where these pests are active. Repeat application every 8 weeks [2 months].

\*Argentine, Southern, Field, Allegheny mound, Florida carpenter, Comfield, Honey, Pavement, Nuisance.

#### **FIRE ANT MOUNDS**

Mix 1.0 fl. oz. (2 Tbsp) in 1 gallon of water and apply to each fire ant mound. Throughly wet the mound and surrounding area to a 4 ft. diameter. For best results apply in cool weather, 65° to 80° F, or in the early morning or late evening. Kills the entire mound, including the queen, in 24 hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment.

#### ORNAMENTAL TREES, SHRUBS AND FLOWERS

[For use on ornamental trees, shrubs and flowers such as but not limited to: Ageratum, Ash, Aster, Arborvitae, Arizona Cypress, Azaleas, Begonia, Birch, Boxwood, Box Elder, Camellias, Cherry (ornamental), Non-bearing Citrus, Chrysanthemums, Carnations, Coleus, Common Ninebark, Conifers, Douglas Fir, Elm, English Ivy, Euonymus, Exacum, Weeping Fig, Fir, Gladioli, Gold

Bells, Hawthorn, Holly, Honeysuckle, Hypoestes, Ivy, Juniper, Lilac, Locust, Maple, Marigold, Mimosa, Mock Orange, Nannyberry, Oak, Orchid, Palm, Pansy, Pea Shrub, Petunia, Philodendron, Pine, Ornamental Plum, Poinsettia, Poplar, Tulip Poplar, Rhododendron, Roses, Snapdragon, Snowberry, Spruce, Taxus, Willow, Zinnia.]

To control [kill] ants\*, armyworms, azalea caterpillar, aphids, bagworms, black vine weevil (adult), box elder bugs, budworms, California oakworm, cankerworms, cockroaches, crickets, cutworms, Eastern tent caterpillar, elm leaf beetles, European sawfly, fall webworm, flea beetles, forest tent caterpillar, gypsy moth larvae, Japanese beetles (adults), June beetles (adults), lace bugs, leaf-feeding caterpillars, leafhoppers, leafminers (adults), leaf rollers, leaf skeletonizers, midges, oleander moth larvae, pillbug, pine sawfly, pine shoot beetles, pinetip moths, plant bugs, root weevil, sawfly, scale insects (crawlers), spiders, spittlebugs, striped beetles, striped oakworm, thrips, tip moth, tussock moth larvae: Mix 0.25 fl. oz. (1/2 Tablespoon) in 1 gallon of water.

To control [kill] broadmites, brown softscale, California redscale (crawler), clover mites, mealybugs, pineneedle scale (crawler), spider mites and whiteflies: Mix 0.5 fl.oz. (1 Tablespoon) in 1 gallon of water.

\*Argentine, Southern, Field, Allegheny mound, Florida carpenter, Comfield, Honey, Pavement, Nuisance.

#### **USE TIPS**

- For best results, apply early before insect populations are high.
- For scale control, treat entire plant, including stem, trunks and twigs.
- For bagworm control, apply when bagworm larvae begin to hatch.
- For best results, apply to both sides of plant leaves and all plant surfaces where insects are present.

#### **VEGETABLES**

Mix 0.5 fl.oz. (1 Tablespoon) in 1 gallon of water and apply to 300 sq.ft. Apply when insects appear or when damage occurs. Spray to wet all infested plant surfaces.

Do not apply more than 5 times per growing season to corn? For all other crops do not apply more than 9 times per growing Season.

For use on Broccoli, Brussels Sprouts, Cabbage, Cavalo Broccoli, Cauliflower, Chinese Broccoli, Chinese Cabbage, Chinese Mustard, and Konirabi to control [kill]: Alfalfa Looper, Cabbage Looper, Imported Catbageworm; Southern Cabbageworm, Cutworm, Cabbage Webworm, Diamondback moth, Armyworm, Beet Armyworm, Fall Armyworm, Yellow-striped Armyworm, Corn Earworm, Flea Beetle, Japanese Beetle (Adult), Grasshopper, Leafhopper,

Plant Bug, including Lygus Bug, Stink Bug, Meadow Spittlebug, Aphid, Whitefly, Thrips, Spider Mite.

For use on Corn (sweet and pop) to control [kill] Cutworm, Western Bean Cutworm, Corn Earworm, Green Cloverworm, Meadow Spittlebug, Tobacco Budworm, European Corn Borer, Southwestern Corn Borer, Stalk Borer, Hop Vine Borer, Armyworm, Beet armyworm, Western Corn Rootworm Beetle (adult), Northern Corn Rootworm Beetle (adult), Southern Corn Rootworm Beetle (adult), Mexican Corn Rootworm Beetle (adult), Bean Leaf Beetle, Cereal Leaf Beetle, Japanese Beetle (adult), Sap Beetle (adult), Stink Bug, Grasshopper, Corn Leaf Aphid, English Grain Aphid, Chinch Bug, Green Bug.

Sweet corn only: Southern Armyworm, Western Bean Cutworm, Tarnished Plant Bug, Aster Leafhopper, Aphid, Spider Mite, Corn silkfly (adult). For use on Lettuce (head and leaf) to control [kill]: Alfalfa Looper, Cabbage Looper, Imported Cabbageworm, Cutworm, Saltmarsh Caterpillar, Green Cloverworm, Diamondback moth, Armyworm, Beet Armyworm, Fall Armyworm, Southern Armyworm, Corn Earworm, Tobacco Budworm, European Corn Borer, Flea Beetle, Japanese Beetle (Adult), Vegetable Weevil (adult), Grasshopper, Leafhopper, Plant Bug, Stink Bug, Meadow Spittlebug, Aphid, Whitefly, Spider Mite.

For use on Onion (bulb) and Garlic to control [kill]: Cutworm, Seedcorn Maggot (adult), Onion Maggot (adult), Leafminer, (adult), Armyworm, Onion Thrips, Tobacco Thrips, Western Flower Thrips, Flower Thrips, Aphid, Plant Bug, Stink Bug.

For Use on Tomato, Tomatillo, Peppers (bell and nonbell), Eggplant, Ground Cherry and Pepino to control [kill]: Cabbage Looper, Cutworm spp., Hornworm, Tomato Fruitworm, Tobacco Budworm, Tomato Pinworm, Beet Armyworm, Southern Armyworm, Yellow-striped Armyworm, Fall Armyworm, European Corn Borer, Leafminer, Colorado Potato Beetle, Flea Beetle, Grasshopper, Leafhopper, Aphid, Whitefly, Meadow Spittlebug, Stink Bug, Plant Bug, Stalk Borer, Blister Beetle, Japanese Beetle (adult), Vegetable Weevil, Cucumber Beetle (adult).

For use on Asparagus beans, Snap Beans, Wax Beans Black-eyed Peas, Garden Peas, Green Peas and Sugar Snap Peas to control [kill]: Cutworm spp., Green Cloverworm, Imported Cabbageworm, Saltmarsh Caterpillar, Velvetleaf Caterpillar, Mexican Bean Beetle, Corn Earworm, Painted Lady Butterfly (Larva), European Corn Borer, Looper spp., Western Bean Cutworm, Armyworm, Yellow-Striped Armyworm, Western Yellow-Striped Armyworm, Bean Leafskeletonizer, Webworm spp., Leaftier spp., Alfalfa Caterpillar, Stalk Borer, Cucumber Beetle (adult), Corn Rootworm Beetle (adult), Flea Beetle (adult), Blister Beetle, Bean Leaf Beetle, Japanese Beetle (adult), Leafhopper spp., Fleas Hopper spp., Three-Cornered Alfalfa Hopper, Meadow Spittle Bug, Stink Bug spp., Plant Bug spp., Grasshopper, Thrips spp., Aphid spp.

For use on Lettuce (Head and Leaf) to control [kill]: Alfalfa Looper, Cabbage Looper, Imported Cabbageworm, Cutworm spp., Saltmarsh Caterpillar, Green Cloverworm, Armyworm, Beet Armyworm, Fall Armyworm, Corn Earworm, European Corn Borer, Flea Beetle spp., Japanese Beetle (adult), Vegetable Beetle (adult), Grasshopper, Leafhopper, Plant Bug, Stink Bug, Meadow Spittlebug,

NUMBER OF DAYS TO WAIT BETWEEN THE LAST APPLICATION AND HARVEST					
VEGETABLES	PHI				
Broccoli, Brussels sprouts, Cabbage, Cavalo Broccoli, Cauliflower, Chinese Broccoli, Chinese Cabbage, Chinese Mustard, Kohlrabi, Lettuce (Head & Leaf)	1				
Tomato, Tomatillo, Peppers (bell and nonbell), Eggplant, Ground Cherry and Pepino	5				
Onions (bulb), Garlic	14				
Corn (sweet and pop)	21				
Asparagus beans, Snap Beans, Wax Beans Black-eyed Peas, Garden Peas, Green Peas and Sugar Snap Peas	7				
Lettuce (Leaf and Head)	1				
PHI = Pre-Harvest Interval					

#### **FRUIT AND NUT TREES**

Mix 0.5 fl.oz. (1 Tablespoon) in 1 gallon of water and apply to 300 sq.ft. Apply when insects appear or when damage occurs. Spray to wet all infested plant surfaces.

For Fruit Trees do not apply more than 9 times per year, and only apply 5 times after blooms have formed. For Nut Trees do not apply more than 6 times per year, and only apply 5 times after blooms have formed.

For use on Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear and Quince to control [kill]: Leafhopper spp., Codling Moth, Tufted Apple Budworm, Oriental Fruit Moth, Lesser Appleworm, Green Fruitworm, Tent Caterpillar spp., Tentiform Leaf Miner spp., Apple Maggot (adult), Pag Sawfly, Flum Curculio, Japanese Beetle, Plant Bug, Stink Bug, Leafhoppes, Remodical Cicada, Apple Aphid, Rosy Apple Aphid, Pear Psylla, San Jose Scale (on fruit only).

For use on Apricot, Sweet Cherry, Tart Cherry, Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot and Prune to control [kill]: Leafroller spp., Peach Twig Borer, Oriental Fruit Moth, Peachtres Borer spp., Green Fruitworm, Tent Caterpillar spp., American Plum Borer, Cherry Fruit Fly spp., (adult), Plum Curculio, Rose Chafer, Japanese Beetle, Plant Bug, Stink Bug, Leafhopper spp., Periodical Cicada, Black Cherry Aphid.

For use on Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut, Walnut (Black and English) and Pecan to control [kill]: Leafroller spp., Navel Orangeworm, Codling moth, Filbertworm, Peach Twig Borer, Walnut Husk Fly spp. (adult), Ants, Plant Bug, Stink Bug, Chinch Bug, Leaffooted Bug, Walnut Aphid, Hickory Shuckworm, Pecan Casebearer spp., Pecan Weevil, Pecan Aphid spp., Pecan Spittlebug, Pecan Phylloxera spp.

NUMBER OF DAYS TO WAIT BETWEEN THE LAST APPLICATION A HARVEST	ND
Fruit and Nut Trees	PHI
Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear, Quince	21
Apricot, Sweet Cherry, Tart Cherry, Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot and Prune	14
Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut, Walnut (Black and English) and Pecan	14
PHI = Pre-Harvest Interval	<del></del>

#### STORAGE AND DISPOSAL

**Storage**: Store in a cool dry area away from heat or open flame. **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### PRECAUTIONARY STATEMENTS

**Hazards to Humans and Domestic Animals** 

**CAUTION:** Harmful if swallowed. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

#### **FIRST AID**

**if Swallowed:** Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a Poison Control Center or doctor for treatment advice.



#### **Environmental Hazards**

This product is extremely toxic to fish. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

[Questions or comments? Call 1-800-332-5553]

[\*If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.]

Made in the USA for Chemsico, Division of United Industries Corporation PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No.: 9688-237

EPA Est. No.: 9688-MO-1

© 2006

# OPTIONAL READY SPRAY NOZZLE DIRECTIONS

STOP. READ ENTIRE LABEL BEFORE USE.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow children or pets into the treated area until dry.

In New York State, this product may not be applied to lawns within 100 ft of a coastal marsh, or stream that drains directly into a coastal marsh.

#### **HOW TO APPLY**

Determine the size of the spraying job. For lawns measure the area to be sprayed. For trees, shrubs and flowers, apply as a thorough cover spray. Begin treating when insects first appear. Applications should be repeated only as directed to maintain control. Apply as directed [in the rate chart]. Remember, thorough coverage is important. Be sure to follow the important use limitations for the specific vegetable you are treating, including the amount of spray that can be applied, the interval between treatments and the days to wait before harvest.

USING THE READY SPRAY NOZZLE (Optional subheading)

- 1. Shake container well before using.
- 2. Connect a garden hose to the Ready Spray nozzle. Make sure the dial on the nozzle is in the "OFF" position with the [green] safety tab in the valve notch.
- 3. Turn on water at faucet. Extend hose to the farthest area to be treated and work back toward the faucet so you don't come in contact with the treated area.
- 4. To BEGIN spraying, point nozzle toward treatment site and a.) bend the [green] safety tab back (located at right of yellow dial) with your thumb, b.) hold while turning yellow dial clockwise until it stops. Water will automatically mix with the product.
- 5. Spray until wet to control insects. Walk at a steady pace while spraying using an even sweeping motion, slightly overlapping treated areas.
- 6. To STOP spraying, QUICKLY turn the dial counter clockwise until it stops and the safety tab engages into the notch on the valve. Turn water off at faucet. To relieve pressure before removing nozzle from hose, bend the safety tab back and turn dial "ON" until water stops spraying.
- 7. To STORE unused product, make sure the dial is in the "OFF" position with [green] safety tab in the valve notch. Place in cool area away from heat, sunlight or open flame.

‡NOTE: This product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding in particular), test in an inconspicuous area and recheck in a few hours. Do not use any if any staining is observed.

#### **LAWNS**

THATCH INFESTING INSECTS/UNDERGROUND INSECTS including Blue grass billbug (adult), Black turfgrass Ataenius (adult), chiggers, chinch bugs, white grubs (Japanese beetle, European chafer, Southern chafer), dung beetle and hyperodes weevils (adult) and mole crickets (nymphs and young adults). Use 1 quart of Chemsico Insecticide Concentrate 25G to treat [cover] 8,000 sq. ft. of lawn. For sub-surface insect control (mole crickets, grubs) water treated area with additional ½ to ½ inch of water for optimum results. Do not apply more than seven (7) times per year per location.

SURFACE INSECTS including ants\*, armyworms, cockroaches, crickets, cutworms, earwigs, fleas, lawn moths (sod webworms), millipedes, mosquitoes, palmetto bugs, sowbugs, mites, spittlebugs, Lone star ticks, Brown dog ticks, American dog ticks, deer ticks (which may transmit Lyme Disease) and waterbugs. Use 1 quart of Chemsico Insecticide Concentrate 25G to treat [cover] 16,000 sq. ft. of lawn. Thoroughly wet down grass a few hours before applying. Delay additional watering or mowing for 24 hours after application for optimum control of surface insects. Do not apply more than fourteen (14) times per year per location.

WHEN TO TREAT: Treat the following pests during the months indicated or when first noticed

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
				Billbugs		(20)	Gı	ubs			
		Section 1	Dùn	g beetle gri	ıbs	Y 2					
			Hyperc	des weevils.							
Trea	t ants*	and of	hêr ins	ects wher	î thev	first a	ppear (	or vou	notice i	awn da	mage

<sup>\*</sup>Argentine, Southern, Field, Allegheny mound, Florida carpenter, Cornfield, Honey, Pavement, Nuisance.

#### ORNAMENTAL TREES, SHRUBS AND FLOWERS

[For use on ornamental trees, shrubs and flowers such as but not limited to: Ageratum, Aster, Arborvitae, Arizona Cypress, Azaleas, Begonia, Birch, Boxwood, Box Elder, Camellias, Carnations, Cherry (ornamental), Chrysanthemums, Carnations, Coleus, Common Ninebark, Conifers, Douglas Fir, Elm, English Ivy, Euonymus, Exacum, Weeping Fig, Fir, Gladioli, Gold Bells, Hawthorn, Holly, Honeysuckle, Hypoestes, Ivy, Juniper, Lilac, Locust, Maple, Marigold, Mimosa, Mock Orange, Nannyberry, Oak, Orchid, Palm, Pansy, Pea Shrub, Petunia, Philodendron, Pine, Ornamental Plum, Poinsettia, Poplar, Tulip Poplar, Rhododendron, Roses, Snapdragon, Snowberry, Spruce, Taxus, Weeping Fig. Willow, Zinnia,

To control [kill] ants\*, armyworms, azalea caterpillar, aphids, bagworms, black vine weevil (adult), boxelder bugs, budworms. California oakworm, cankerworms, cockroaches, crickets, cutworms, Eastern tent caterpillar, elm leaf beetles, European sawfly, fall webworm, eflea beetles, forest tent caterpillar, gypsy moth larvae, Japanese beetles (adults); June beetles (adults), lace bugs, leaf-feeding caterpillars, leafhoppers, leafminers (adults), leafrollers, leaf skeletonizers, midges, oleander moth larvae, pillbug, pine sawfly, pine shoot beetles, pinetip moths, plant bugs, root wedvil, pawfly, scale insects (crawlers), spiders, spittlebugs, striped beatlass otriped oakworm, thrips, tip moth, tussock moth larvae proadmites brown coffecale. California redscale (crawler), clover mites, mealybugs, pineneedle scale (crawler), spider mites and whiteflies.

\*Argentine, Southern, Field, Allegheny mound, Florida carpenter, Cornfield, Honey, Pavement, Nuisance,

Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to spray underside of leaves and penetrate dense foliage.

#### **USE TIPS**

- For best results, apply early before insect populations are high.
- For scale control, treat entire plant, including stem, trunks and twigs.
- For bagworm control, apply when bagworm larvae begin to hatch.
- For best results, apply to both sides of plant leaves and all plant surfaces where insects are found.

#### **VEGETABLES**

Apply when insects appear or when damage occurs. Spray to wet all infested plant surfaces. Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to spray underside of leaves and penetrate dense foliage. 1 gt. will treat 19,200 sq. ft. of vegetable plants.

Do not apply more than 5 times per growing season to corn. For all other crops do not apply more than 9 times per growing season.

For use on Broccoli, Brussels Sprouts, Cabbage, Cavalo Broccoli, Cauliflower, Chinese Broccoli, Chinese Cabbage, Chinese Mustard, and Kohlrabi to control [kill]: Alfalfa Looper, Cabbage Looper, Imported Cabbageworm, Southern Cabbageworm, Cutworm, Cabbage Webworm, Diamondback moth. Armyworm, Beet Armyworm, Fall Armyworm, Yellow-striped Armyworm, Corn Earworm, Flea Beetle, Japanese Beetle (Adult), Grasshopper, Leafhopper, Plant Bug, including Lygus Bug, Stink Bug, Meadow Spittlebug, Aphid, Whitefly, Thrips, Spider Mite.

For use on Corn (sweet and pop) to control [kill] Cutworm, Western Bean Cutworm, Corn Earworm, Green Cloverworm, Meadow Spittlebug, Tobacco Budworm, European Corn Borer, Southwestern Corn Borer, Stalk Borer, Hop Vine Borer, Armyworm, Beet armyworm, Western Corn Rootworm Beetle (adult), Northern Corn Rootworm Beetle (adult), Southern Corn Rootworm Beetle (adult), Mexican Corn Rootworm Beetle (adult), Bean Leaf Beetle, Cereal Leaf Beetle, Japanese Beetle (adult), Sap Beetle (adult), Stink Bug, Grasshopper, Corn Leaf Aphid, English Grain Aphid, Chinch Bug, Green Bug.

Sweet corn only: Southern Armyworm, Western Bean Cutworm, Tarnished Plant Bug, Aster Leafhopper, Aphid, Spider Mite, Corn silkfly (adult).

For use on Lettuce (head and leaf) to control [kill]: Alfalfa Looper, Cabbage Looper, Imported Cabbageworm, Cutworm, Saltmarsh Caterpillar, Green Cloverworm, Diamondback moth, Armyworm, Beet Armyworm, Fall Armyworm, Southern Armyworm, Corn Earworm, Tobacco Budworm,

European Corn Borer, Flea Beetle, Japanese Beetle (Adult), Vegetable Weevil (adult), Grasshopper, Leafhopper, Plant Bug, Stink Bug, Meadow Spittlebug, Aphid, Whitefly, Spider Mite.

For use on Onion (bulb) and Garlic to control [kill]: Cutworm, Seedcorn Maggot (adult), Onion Maggot (adult), Leafminer, (adult), Armyworm, Onion Thrips. Tobacco Thrips, Western Flower Thrips, Flower Thrips, Aphid, Plant Bug, Stink Bug.

For Use on Tomato, Tomatillo, Peppers (bell and nonbell), Eggplant, Ground Cherry and Pepino to control [kill]: Cabbage Looper, Cutworm spp., Hornworm, Tomato Fruitworm, Tobacco Budworm, Tomato Pinworm, Beet Armyworm, Southern Armyworm, Yellow-striped Armyworm, Fall Armyworm, European Corn Borer, Leafminer, Colorado Potato Beetle, Flea Beetle, Grasshopper, Leafhopper, Aphid, Whitefly, Meadow Spittlebug, Stink Bug, Plant Bug, Stalk Borer, Blister Beetle, Japanese Beetle (adult), Vegetable Weevil, Cucumber Beetle (adult).

For use on Asparagus beans, Snap Beans, Wax Beans Black-eved Peas, Garden Peas, Green Peas and Sugar Snap Peas to control [kill]: Cutworm spp., Green Cloverworm, Imported Cabbageworm, Saltmarsh Caterpillar, Velvetleaf Caterpillar, Mexican Bean Beetle, Corn Earworm, Painted Lady Butterfly (Larva), European Corn Borer, Looper spp., Western Bean Cutworm. Armyworm, Yellow-Striped Armyworm, Western Yellow-Striped Armyworm, Bean Leafskeletonizer, Webworm spp., Leaftier spp., Alfalfa Caterpillar, Stalk Borer, Cucumber Beetle (adult), Corn Rootworm Beetle (adult), Flea Beetle (adult), Blister Beetle, Bean Leaf Beetle, Japanese Beetle (adult), Leafhopper spp., Fleas Hopper spp., Three-Cornered Alfalfa Hopper, Meadow Spittle Bug, Stink Bug spp., Plant Bug spp., Grasshopper, Thrips spp., Aphid spp.

For use on Lettuce (Head and Leaf) to control [kill]: Alfalfa Looper, Cabbage Looper, Imported Cabbageworm, Cutworm spp., Saltmarsh Caterpillar, Green Cloverworm, Armyworm, Beet Armyworm, Fall Armyworm, Corn Earworm, European Corn Borer, Flea Beetle spp., Japanese Beetle (adult), Vegetable Beetle (adult), Grasshopper, Leafhopper, Plant Bug, Stink Bug, Meadow Spittlebug.

NUMBER OF DAYS TO WA	NIT BETWEEN THE PAST APPLICATION AN	D
VEGETABLES		PHI
Broccoli, Chinese Cabbage, Chi Leaf)	age, Cavalo Broccoli, Cauliflower, Chinese nese Mustard, Kohlrabi, Lettuce (Head &	1
Tomato, Tomatillo, Peppers (bell and Pepino	and nonbell) ဥဋဋဌgclant, Ground Cherry இத	5
Onions (bulb), Garlic		14

Corn (sweet and pop)	21
Asparagus beans, Snap Beans, Wax Beans Black-eyed Peas, Garden Peas, Green Peas and Sugar Snap Peas	7
Lettuce (Leaf and Head)	1
PHI = Pre-Harvest Interval	

#### **FRUIT AND NUT TREES**

Apply when insects appear or when damage occurs. Spray to wet all infested plant surfaces. Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to spray underside of leaves and penetrate dense foliage. 1 gt. will treat 19.200 sq. ft. of fruit and nut trees.

For Fruit Trees do not apply more than 9 times per year, and only apply 5 times after blooms have formed. For Nut Trees do not apply more than 6 times per year, and only apply 5 times after blooms have formed.

For use on Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear and Quince to control [kill]: Leafhopper spp., Codling Moth, Tufted Apple Budworm, Oriental Fruit Moth, Lesser Appleworm, Green Fruitworm, Tent Caterpillar spp., Tentiform Leaf Miner spp., Apple Maggot (adult), Pear Sawfly, Plum Curculio, Japanese Beetle, Plant Bug, Stink Bug, Leafhopper, Periodical Cicada, Apple Aphid, Rosy Apple Aphid, Pear Psylla, San Jose Scale (on fruit only).

For use on Apricot, Sweet Cherry, Tart Cherry, Nectarine, Peach, Plum (Chickasaw, Damson, Japanese). Plumcot and Prune to control [kill]: Leafroller spp., Peach Twig Borer, Oriental Fruit Moth, Peachtree Borer spp., Green Fruitworm, Tent Caterpillar spp., American Plum Borer, Cherry Fruit Fly spp., (adult), Plum Curculio, Rose Chafer, Japanese Beetle, Plant Bug, Stink Bug, Leafhopper spp., Periodical Cicada, Black Cherry Aphid.

For use on Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut, Walnut (Black and English) and Pecan to control [kill]: Leafroller spp., Navel Orangeworm, Codling moth, Filbertworm, Peach Twig Borer, Walnut Husk Fly spp. (adult), Ants. Plant Bug, Stink Bug, Chinch Bug, Leaffooted Bug, Walnut Aphid, Hickory Shuckworm, Pecan Casebearer spp., Pecan Weevil, Pecan Aphid spp., Pecan Spittlebug, Pecan Phylloxera spp.

NUMBER OF DAYS TO WAIT BETWEEN THE LAST APPLICATION AND HARVEST						
Fruit and Nut Trees	PHI					
Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear, Quince	21					
Apricot, Sweet Cherry, Tart Cherry, Nectarine, Peach, Plum (Chickasaw,	14					
Damson, Japanese), Plumcot and Prune						

Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin,	14
Filbert (Hazelnut), Hickory Nut, Macadamia Nut, Walnut (Black and	·
English) and Pecan	
PHI = Pre-Harvest Interval	

#### STORAGE AND DISPOSAL

**Storage:** Store in a cool dry area away from heat or open flame. **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

**CAUTION:** Harmful if swallowed. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

#### **FIRST AID**

If Swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a Poison Control Center or doctor for treatment advice.

#### **Environmental Hazards**

This product is extremely toxic to fish. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring greecy

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

NOTICE: Buyer assumes all responsibility for safety and fish inclining accordance with directions.

[Questions or comments? Call 1-800-332-5553]

[\*If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.]

Made in the USA for Chemsico, Division of United Industries Corporation PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No.: 9688-237

EPA Est. No.: 9688-MO-1

© 2006

MAY OF THE MAY

