



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

March 21, 2024

Michael Sarli
Sr. Director Regulatory
Chemsico, A Div. Of United Industries Corp.
P.O. Box 142642
St. Louis, MO 63114

Subject: Label Amendment - Registration Review Mitigation for MCPA, Quinclorac, & Sulfentrazone
Product Name: Chemsico RTU Herbicide 1462
EPA Registration Number: 9688-280
Application Dates: December 3, 2021; January 19, 2022; & July 14, 2023
Decision Numbers: 580499; 581287; & 592917

Dear Michael Sarli:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the MCPA, Quinclorac, and Sulfentrazone Interim Decisions, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling

before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Caleb Carr by phone at (202) 566-0636, or via email at carr.caleb@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington", with a stylized flourish at the end.


Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

Chemsico RTU Herbicide 1462

Two sublabels represent the entire master label of Chemsico RTU Herbicide 1462. Please refer to the appropriate section of the labeling as shown as follows:

Section
Label Language Common to All Sublabels
Sublabel 1
Sublabel 2
Appendix



Label Language Common to All Sublabels

Chemsico RTU Herbicide 1462

ACTIVE INGREDIENT:

MCPA, dimethylamine salt.....	0.257%
Quinclorac	0.118%
Dicamba, dimethylamine salt	0.029%
Sulfentrazone	0.015%
INERT INGREDIENTS:	<u>99.581%</u>
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- 0.018 lb 2-methyl-4-chlorophenoxyacetic acid equivalent per gallon or 0.210%
- 0.010 lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 0.118%
- 0.002 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.024%
- 0.001 lb N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per gallon or 0.015%

[NOTE: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then we ask to add a referral statement and present the substatement on the back panel: See back panel for substatement of ingredient statement.]

KEEP OUT OF REACH OF CHILDREN CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid	
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to aquatic invertebrates and may be toxic to fish and aquatic plants. Do not apply directly to water. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 48 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. Drift and runoff may be hazardous to plants in adjacent areas.

NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

This product is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water.

This product is classified as having high potential for reaching surface water via runoff for several months or more after application.

SUBLABEL 1

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift.
- Take precautions to minimize drift.
- Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- This product may only be used to control listed broadleaf and grass weeds on residential lawns.
- Do not apply more than 75 gallons of product per acre per application (equivalent to 0.75 lbs of Quinclorac/acre, 1.35 lbs of MCPA dimethylamine salt acid/acre, 0.15 lbs of Dicamba dimethylamine salt acid/acre and 0.075 lbs of Sulfentrazone/acre).
- For all application methods, the maximum number of treatments is limited to 2 per year with a minimum of 21 days between applications.
- Do not apply more than 150 gallons of product per acre per year (equivalent to 1.5 lbs of Quinclorac/acre, 2.7 lbs of MCPA dimethylamine salt acid/acre, 0.30 lbs of Dicamba dimethylamine salt acid/acre and 0.15 lbs of Sulfentrazone/acre).

For home lawn care:

Chemsico RTU Herbicide 1462 is a ready-to-use (RTU) product intended for "spot" treatment of individual broadleaf and grass weeds by homeowners on residential lawns. This product should be applied to lawns containing the following species of grass: Kentucky bluegrass, perennial ryegrass, fescue species (including tall, red, and fine leaf varieties), bermudagrass¹, buffalograss and zoysiagrass.

¹Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.

This product should **not** be used on lawns containing bentgrass, St Augustinegrass, bahiagrass, centipedegrass, or carpetgrass. Do **not** use on dichondra or lawns with desirable clovers or legumes, on vegetable gardens, fruits, or ornamentals (flowers, trees, groundcovers, hedges, landscape/flower beds and shrubs)

[options for different size containers]

- This 32 fl.oz. (1.0 quart) retail container size treats approximately 250 weeds or up to 250 sq.ft.
- This 128 fl.oz. (1.0 gallon) retail container size treats approximately 1,000 weeds or up to 1,000 sq.ft.
- This 1.3 gallon retail container size treats approximately 1,300 weeds or up to 1,300 sq.ft.

WEEDS CONTROLLED: This product will control or suppress the following broadleaf and grass weeds. Partial weed lists are presented below:

Broadleaf Weeds			
Bedstraw	Ground ivy	Morningglory	Redweed
Black medic	Healall	Old world diamond flower	Sheep sorrel
Buckhorn	Henbit	Peppergrass	Shepherdspurse
Burdock	Innocence (Blue-eyed	Pigweed	Speedwell
Chicory	Mary)	Plantain	Spurge
Chickweed	Knotweed	Poison ivy	Wild carrot
Clover	Lambsquarters	Poison oak	Wild garlic
Corn speedwell	Lawn burweed	Purple cudweed	Wild lettuce
Dandelion	Lespedeza	Purslane	Wild onion
Dock	Mallow	Ragweed	Yarrow

Grassy Weeds		
Crabgrass (large and smooth)	Foxtail (green, giant and yellow)	Yellow nutsedge

[HAND SPRAYER PRODUCT]
(Directions & Storage and Disposal)

1. Shake well.

2. Connect trigger sprayer. Unscrew and discard the white cap on the bottle. Insert the hose into the bottle and tighten the cap securely.
3. Prime Sprayer: Turn the sprayer nozzle one-half turn counter-clockwise. Hold the sprayer lower than the bottle and squeeze the trigger several times until it sprays. Adjust the nozzle to coarse spray pattern to minimize wind drift. (graphic)
4. Spray just enough [Chemsico RTU Herbicide 1462] [product] to lightly cover [broadleaf and grass] weeds. **Excessive spraying can injure some lawns.** Do not apply when daytime temperatures are above 90° F as lawn injury can occur. (graphic) Spray only the weeds, not the entire lawn.
5. Weeds will begin wilting and browning overnight.

USE INFORMATION

- Some hard-to-kill weeds may require re-treatment. Wait at least 21-days between applications - some weeds can take that long to die.
- Do not allow spray to drift onto desirable plants since injury may result.
- Spot treatment is defined as a treatment area no greater than 1,000 sq. ft. per acre.
- The maximum number of spot treatments is limited to 2 per year with a minimum of 21 days between applications.
- Not for use on turf being grown for sale or other commercial use such as sod, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Turn the nozzle to "OFF" position and store with sprayer above the level of liquid in container to prevent leakage. Store in original container in an area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

[TRIGGER SPRAYER PRODUCT]

(Directions & Storage and Disposal)

1. Shake well.
2. Adjust nozzle to coarse spray pattern to minimize wind drift. (graphic)
3. Spray just enough [Chemsico RTU Herbicide 1462] [product] to lightly cover [grass and broadleaf] weeds. **Excessive spraying can injure some lawns.** Do not apply when daytime temperatures are above 90°F as lawn injury can occur. (graphic) Spray only the weeds, not the entire lawn.
4. Weeds will begin wilting and browning overnight.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Turn the nozzle to "OFF" position and store with sprayer above the level of liquid in container to prevent leakage. Store in original container in an area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE INFORMATION

- Some hard-to-kill weeds may require re-treatment. Wait at least 21-days between applications - some weeds can take that long to die.
- Do not allow spray to drift onto desirable plants since injury may result.
- Spot treatment is defined as a treatment area no greater than 1,000 sq. ft. per acre.
- The maximum number of spot treatments is limited to 2 per year with a minimum of 21 days between applications.

- Not for use on turf being grown for sale or other commercial use such as sod, or for commercial seed production, or for research purposes.

[CAPTIVE SPRAYER PRODUCT]

(Directions & Storage and Disposal)

1. Shake well before each use
2. Peel sticker from flip-top cap and open cap. Discard sticker.
3. Remove sprayer from **its** slot beneath the bottle handle and remove and discard plastic wrap. Pull black hose to completely extend tubing.
4. Firmly press black hose lip into opening in cap until it snaps into place
5. Hold sprayer below the level of the container and prime by pumping trigger several times until product is dispensed.
6. Adjust the nozzle to a coarse spray pattern to avoid sprays that may drift and harm other plants.
7. Spray just enough [product] to lightly cover [broadleaf] weeds. **Excessive spraying can injure some lawns.** Do not apply when daytime temperatures are above 85°F as lawn injury can occur. (graphic) Spray only the weeds, not the entire lawn.
8. Weeds will begin wilting and browning overnight.
9. To store sprayer after use turn sprayer nozzle off
10. DO NOT disconnect hose
11. Return sprayer to its slot. Hose can be pushed back into sprayer handle.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Turn the nozzle to "OFF" position and store with sprayer above the level of liquid in container to prevent leakage. Store in original container in an area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.weosite.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE INFORMATION

- Some hard-to-kill weeds may require re-treatment. Wait at least 30 days between applications - some weeds can take that long to die.
- Do not allow spray to drift onto desirable plants since injury may result.
 - Do not use on 'Floritam' or other improved varieties of St. Augustinegrass grown in Florida.
 - Spot treatment is defined as a treatment area no greater than 1,000 sq. ft. per acre.
 - The maximum number of spot treatments is limited to 2 per year with a minimum of 21 days between applications.
 - Not for use on turf being grown for sale or other commercial use such as sod, or for commercial seed production, or for research purposes.

[Optional Directions for Use]

DIRECTIONS FOR USE for EzSPRAY(TM) applicator:

EzSPRAY(TM) EASY SET UP

1. Remove sprayer from side holster. Unwrap the tubing (Graphical representation)
2. Remove seal from top of cap. Unplug cap. (Graphical representation)
3. Securely snap the [color] plug into hole in cap. (Graphical representation)
4. Remove plastic tab near bottom of sprayer to activate batteries. (Graphical representation)
5. Twist nozzle to adjust spray from CLOSE to SPRAY or STREAM. (Graphical representation)
6. When priming the sprayer, hold sprayer level to the ground. Sprayer will not prime if it is held at an angle. Once primed it will spray in any direction (Graphical representation)

HOW TO USE THE EzSPRAY(TM)[@]

- A. Aim toward target weeds/plants, and away from people and pets.
- B. Press the trigger to spray.
- C. Release the trigger to stop spray.

After use:

1. Twist nozzle completely to CLOSE position
2. Secure sprayer upright in holder.
3. **Failure to adjust nozzle completely to CLOSE position and properly secure sprayer**

in holder may result in leakage and damage to property or injury to people or animals.

To replace batteries:

1. Push release button near nozzle and lift up on battery hatch.
2. Remove used batteries and insert two new AA batteries **in correct positions** as marked per diagram inside of battery compartment. Never insert the positive end where the negative end belongs and vice versa. **Always** use a complete set of new batteries of the same type when replacing batteries. **Never** mix alkaline, carbon-zinc or rechargeable batteries.
3. Securely close battery compartment door.
4. **Always** follow manufacturer's recommendations for use and disposal of batteries.

Additional operating notes:

- Do not submerge in water
- Before each use inspect sprayer carefully-makes sure hose is flexible and not kinked, worn or cracked, and that all connections are tight.
- When storing sprayer for prolonged periods remove batteries.

[REFILL DIRECTIONS

Refill only with Chemsico Herbicide Concentrate 1464 [brand name]. Remove cap. Carefully measure and mix 8.5 fl.oz. (17 Tbsp.) in one gallon of water. Replace and tightly close cap. Wipe off any spilled product.

Additional advertising items to be included on **EzSPRAY(TM)** container label:

- Battery Powered [with graphic]
- **EzSPRAY(TM) NO PUMPING - NO PULLING - NO FATIGUE!**
- 2 AA Batteries Included! (with graphic)]

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Turn the nozzle to "CLOSE" position. To prevent leakage, attach sprayer to clip. Store this product only in its original container in a secure storage area inaccessible to children and pets. Keep from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable Container. Do not reuse container except to refill (refer to REFILL DIRECTIONS). **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

***GUARANTEED RESULTS or your money back.** If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.

SUBLABEL 2

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift.
- Take precautions to minimize spray drift.
- Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- This product may only be used to control listed broadleaf and grass weeds on residential lawns.
- Do not apply more than 75 gallons of product per acre per application (equivalent to 0.75 lb. of Quinclorac/acre 1.35 lbs. of MCPA dimethylamine salt acid/acre, 0.15 lbs Dicamba, dimethylamine salt/acre and 0.075 lbs of Sulfentrazone/acre).
- For all application methods, the maximum number of treatments is limited to 2 per year with a minimum of 21 days between applications
- Do not apply more than 150 gallons of product per acre per application (equivalent to 1.5 lbs of Quinclorac/acre, 2.70 lbs of MCPA dimethylamine salt acid/acre, 0.3 lbs of Dicamba dimethylamine salt/acre and 0.15 lbs of Sulfentrazone/acre).

WHERE TO USE:

On Lawns: This product is intended for use by homeowners as a spot treatment on lawns. This product should be applied only to the following species of turfgrass:

Cool season turfgrass	Warm season turfgrass
Kentucky bluegrass Perennial ryegrass Fescue spp., including tall, red and fine leaf fescues	Bermudagrass ¹ Buffalograss Zoysiagrass
¹ Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.	

DO NOT use Chemsico RTU Herbicide 1462 for the following:

Cool season turfgrass	Warm season turfgrass	Other areas
Bentgrass, creeping Bentgrass, 'Seaside' Bentgrass, colonial	St. Augustinegrass Bahigrass Centipedegrass Carpetgrass Seashore paspalum	Lawns with desirable clovers or legumes Dichondra Vegetable gardens or fruits (food crops) Ornamentals (flowers, trees, groundcovers, hedges, landscape/flower beds and shrubs)

WHEN TO USE

Generally, apply this product to actively-growing weeds during the early growth stages for the best results. Two or more applications at 21 day intervals throughout the season may be required for grass weeds. Apply when daytime temperatures are between 45'and 90' F.

SPRING AND EARLY SUMMER:

- **Crabgrass (large and smooth):** Apply this product from crabgrass emergence until the 6 leaf stage (2 tillers) or when the crabgrass is 3 to 4 inches tall. This period begins the first time that you can see crabgrass in your lawn. Use two applications with a 14-day interval between applications.
- **Foxtail (green, giant and yellow):** Apply this product to foxtail during the early growth stages or when the foxtail is 1 to 4 inches tall. Use two applications with a 14-day interval between applications.
- **Yellow nutsedge:** Two applications are required when the yellow nutsedge plants are 3 to 6 inches tall. The second application should follow the first application 21 days later.
- **Dandelion, plantain, buttercup, speedwell, knotweed, pennywort (dollarweed) and other broadleaf weeds:** Apply when broadleaf weeds are young and actively growing with adequate soil moisture.

FALL:

- **Henbit, chickweed, white clover, ground ivy, lawn burweed, pearlwort, Carolina geranium and other broadleaf weeds:** Apply when broadleaf weeds are young and actively growing with adequate

soil moisture for best results.

Note for weeds controlled: Supplemental registrants may select broadleaf weeds from the list presented in the appendix and may include additional species on the container labels for different geographic regions.

HOW TO USE:

Ready-To-Use: No mixing. No measuring. No fuss.

For Containers with Triggers Sprayers: Chemsico RTU Herbicide 1462 is ready-to-use and is applied directly from the container fitted with the trigger sprayer. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Usually, two (2) pumps of this trigger sprayer will provide adequate coverage of individual weeds. Excessive spraying can injure some lawns.

For Refill Containers: Chemsico RTU Herbicide 1462 is ready-to-use and is applied directly from the container. Remove the cap from the container and attach the trigger sprayer. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Excessive spraying can injure some lawns.

USE PRECAUTIONS:

- Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants.
- Avoid applications of this product when turfgrasses are under stress since injury may result

CULTURAL TIPS:

For newly seeded areas of Kentucky bluegrass, fine fescue blends and perennial ryegrass: Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.

For newly sodded, sprigged, or plugged areas of bermudagrass and zoysiagrass: The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

Seeding: Turf species listed on this label can be seeded into the treated areas at four (4) weeks after the application of this product.

Mowing: Delay mowing 2 days before and until 2 days after the application of this product.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area. Keep from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents:
- EPA Est. No.

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

SUBLABEL 1:

- For use on Many Northern and Southern Grasses including Bluegrass, Fescue, Zoysia, Bermuda, and Perennial Ryegrass
- Kills the Weeds, not the Lawn
- Kills xx+ [Listed] Weed Types [As Listed] Roots and All.
- Kills weed roots
- Kills xx+ [Listed] Weeds [As Listed]
- Kills Dandelion, Chickweed, Wild onion, Plantain, Poison ivy and many others. Weed details on back. See Results in Hours.
- Overnight results • Rainproof in hours • Rainproof in as little as 3 hours - won't wash away.
- Water-Based
- Especially effective on all types of hard-to-kill weeds
- Kills xx+ [listed] types of weeds [as listed] • Kills weed roots! • Kills major broadleaf weeds
- Clear-view measuring window
- For home lawn care.
- Kills over xx [listed] weed types [as listed] including dandelion, chickweed, wild onion, plantain and poison ivy
- Kills broadleaf weeds in lawns
- Kills major broadleaf weeds, roots and all. And [brand name] is formulated to kill only the weeds, not the lawn. Use as directed to kill major broadleaf weeds in your lawn.
- Kills dandelion and many other broadleaf weeds
- [Brand name] controls more than xx troublesome weeds including dandelion, clover, plantain and wild onion. It kills even woody weeds like poison ivy and poison oak in both northern and southern lawns.
- Kills the Weeds! As listed on our Weed Stop product.
- Use on Lawns to Kill the Toughest Weeds, Like Clover and Dandelion.
- includes crabgrass control]
- All-in-one [complete] [total] [2-in-1] weed control [for lawns]
- kills [controls] crabgrass [yellow nutsedge] [foxtail]
- Kills [controls] types of [lawn] weeds
- Targets the toughest [stubborn] weeds [crabgrass] [yellow nutsedge].
- 100% root kill
- Kills the toughest weeds - down to the root
- [Kills the root so] weeds won't come back
- Lawn- formula
- Weed deadly. Lawn friendly.
- Kills weeds [including crabgrass] - not the lawn.
- For a weed-free lawn.
- Weed-free. Worry-free.
- No weeds. No worry.
- Starts working immediately
- Fast acting [formula]
- Destroys weeds [dandelions] [clover] [oxalis] [crabgrass] [yellow nutsedge] completely - [down to the root].
- Long lasting [formula]
- [Works fast] on [hard-to-kill] weeds
- [Even] works in cold [and hot] weather [temperatures]
- Let [Weed Stop] [the liquid] do the weeding

- Kills on contact
- A simple [an easy] step towards a weed free lawn [yard]
- kills types of weeds - [grassy or broadleaf]
- Won't Harm Lawns (when used as directed).

EFFICACY

- Tough weed killer
- Kills even the toughest broadleaf weeds [like (clover) (dandelion) (oxalis)]
- Kills even tough broadleaf weeds [like (clover) (dandelion) (oxalis)]
- Treats Many Northern and Southern Grasses
- Treats Bluegrass, Fescue, Zoysia, Bermuda, Perennial ryegrass
- Destroys the hard-to-kill broadleaf weeds [like (clover) (dandelion) (oxalis)]
- Kills major broadleaf weeds [like (clover) (dandelion) (oxalis)]
- Kills dandelion, chickweed, wild onion, oxalis, plantain, poison ivy and many others. [See Weed details on back.]
- Kills [major] broadleaf weeds
- Kills broadleaf weeds in lawns
- Kills poison ivy and poison oak
- Provides broadleaf control
- [Controls][Kills] clover, dandelion, plantain, chickweed and [other broadleaf weeds] [others as listed]
- Clobbers clover
- [Complete] clover control [as little as one week]
- Controls clover in one week
- Kills XX [listed] [types of] [brush and] weeds [as listed] [roots and all]
- Kills weed roots
- Kills the root
- Kills weed roots!
- Kills weeds
- Kills tough lawn weeds

SPEED

- Overnight results
- [See] results in hours
- Consistently fast on tough weeds like clover, plantain, ground ivy and spurge [various other weeds listed]
- Starts working on contact

WEATHER

- [Cool][all] weather performance
- [cool] weather performance
- Performs in both hot and cold weather

RAINPROOF

- Rainproof - [For control that] won't wash away
- Can't be washed [away][off]
- Can't be washed [off by rain]
- [Rainfast][Rainproof] [in hours [Won't wash away]]
- Rainproof - Won't wash away

OTHER

- Water-based
- Guaranteed [results] or your money back
- Clear-view measuring window
- For home lawn care
- For home lawns
- For a beautiful, weed-free lawn
- Helps keep lawns weed-free
- Ideal for spot treating individual lawn weeds
- Kills the weeds, not the lawn
- Kills weeds, not lawn grasses

- Kills [down to] the root, [not the lawn]
- For use on lawns
- Targets weeds, leaves [grasses] [the lawn] unharmed
- Kills yellow nutsedge
- Visible results in 8 hours
- Kills both broadleaf and grassy weeds
- Kills types of weeds
- Rainfast in 3 hours

EQUAL MEASURES:

- 1 fl.oz. = 2 Tablespoons = 6 Teaspoons (tsp.)
- 2 fl.oz. = 4 Tablespoons
- 4 fl.oz. = 1/2 cup
- 8 fl.oz. = 1 cup
- 16 fl.oz. = 1 pint
- 32 fl.oz. = 1 quart

PRODUCT FACTS What it Does: [optional info box]

What It Does: Kills major broadleaf and troublesome grass weeds Including dandelion, chickweed and clover as well as crabgrass, foxtail, and yellow nutsedge. (See inside for complete list)

Where to Use: (icon weeds)

Lawns: Kills the weeds not the lawn

When to Use:

Apply when daytime temperatures are between 45° and 90°F.

Questions & Comments: (icon phone)(icon computer)

Call 1-800-xxx-xxxx or Visit our Website at www.companyname.com

WEEDS			
Amaranth, purple	Cupid shaving brush	Mugwort	Speedwell, common
Amaranth, slender	Cudweed	Mustard, ball	Speedwell, corn
Aster, heath	Daisy, English	Mustard, black	Speedwell, ivyleaf
Aster, white heath	Daisy, oxeye	Mustard, blue	Speedwell, Persian
Aster, purple	Dandelion, common	Mustard, hedge	Speedwell, purslane
Aster, white prairie	Dandelion, false	Mustard, Indian,	Speedwell, slender
Artichoke	Day flower	Mustard, tansy	Speedwell, snow
Austrian fieldcress	Deadnettle	Mustard, tumble	Speedwell, thymeleaf
Bedstraw, catchweed	Dichondra	Mustard, white	Spiny amaranth
Bedstraw, smooth	Dock, broadleaf	Mustard, wormseed	Spurge, spotted
Bedstraw, northern	Dock, curly	Nettle, stinging {burning}	Spurge, prostrate
Bedstraw, pineywoods	Dogbane	Nettle, tall	Spurweed
Beggarticks, hairy	Dogfennel	Nettle, wood	Star of Bethlehem
Betony, Florida	Dollarweed	Parsley-piert	Strawberry, Indian mock
Bindweed, Field	{pennywort}	Parsnip	Tansy ragwort
Bindweed, hedge	Doveweed	Pearlwort	Tanweed
Bitter Wintercress	Elderberry	Pennycress (Frenchweed)	Thistle, blessed
Bittercress, hairy	Falseflax	Pepperweed, perennial	Thistle, bull
Bitterweed	False sunflower	Pepperweed, Virginia	Thistle, Canada
Black-eyed Susan	Fiddleneck	{peppergrass}	Thistle, flodman {prairie}
Black medic	Filaree, redstem	Pigweed, prostate	Thistle, musk
Blood flower milkweed	Filaree, whitestem	Pigweed, redroot	Thistle, platte
Blue lettuce	Fleabane, daisy	Pigweed, Russian	Thistle, plumeless
Brassbuttons	{annual}	Pigweed, smooth	Thistle, Russian
Bristly oxtongue	Fleabane, rough	Pigweed, tumble	Thistle, Scotch
Broomweed	Galinsoga, hairy	Pineappleweed	Thistle, wavyleaf
Buckhorn	Galinsoga, smallflower	Plains coreopsis {tickseed}	Thistle, yellow {horrible}
Bullnettle	Goldenrod	Plantain, blackseed	Trailing crownvetch
Burclover	Ground ivy	Plantain, bracted	Velvetleaf

WEEDS			
Burdock	Groundsel	Plantain, broadleaf	Vervain, blue
Burweed, lawn (spurweed)	Gumweed	Plantain, buckhorn	Vervain, hoary
Buttercupbulbous	Hairy fleabane	Plantain, hoary	Vervain, prostrate
Buttercup, creeping	Hawkweed, meadow	Plantain, narrowleaf	Vervain, tall
Buttonweed, common	Hawkweed, mouseear	Plantain, slender	Vetch, bird
(poorjoe)	Hawkweed, orange	Plantain, woolly	Vetch, hairy
Buttonweed, Virginia	Hawkweed, yellow	Poison ivy	Vetch, milk
Carolina geranium (wild)	Healall	Poison oak	Vetch, narrowleaf
Carpetweed	Heartleaf drymary	Pokeweed	Vetch, wild
Catnip	Hemp	Prairie sunflower	Virginia creeper
Catsear, spotted (false dandelion)	Henbit	Prickly lettuce (compassplant)	Western clematis
Chamber bitter	Hoary cress	Prickly sida	Western salsify
Chickweed, common	Horsenettle	Prostrate knotweed	Wild aster
Chickweed, mouseear	Horseweed	Puncturevine	Wild buckweed
Chickweed, sticky	Jimsonweed	Purslane, common	Wild carrot
Chicory	Johnny-Jumpup violet	Pusley, Brazil	Wild four-o'clock
Cinquefoil	Knawel	Pusley, Florida	Wild garlic
Clover, crimson	Knotweed, prostrate	Pusley, large flower	Wild lettuce
Clover, alsike	Kochia	Ragweed, bur	Wild marigold
Clover, hop	Lambsquarters	Ragweed, common	Wild mustard (yellow rocket)
Clover, rabbit foot	Lespedeza	Ragweed, lanceleaf	Wild onion
Clover, red	Mallow, alkali	Ragweed, western	Wild parsnip
Clover, strawberry	Mallow, bristly	Redstem filaree	Wild radish
Clover, yellow sweet	Mallow, common	Rough cinquefoil	Wild rape
Clover, white sweet	Mallow, dwarf	Scarlet pimpernel	Wild strawberry
Clover, white	Mallow, Venice	Shepherdspurse	Wild sweet potato
Cockle	Marcela	Smartweed, ladythumb	Wild violet
Cocklebur	Matchweed	Smartweed, pale	Woodsorrel, creeping
Cocklebur, oriental	Mexicanweed	Smartweed, Pennsylvania	Woodsorrel, violet
Cocklebur, spiny	Moneywort	Smooth chaff-flower	Woodsorrel, yellow (Oxalis)
Common mullein	Morningglory, bigroot	Smooth dock	Woolly croton
Com spurry	Morningglory, ivyleaf	Sorrel, red (sheep)	Wormseed
Creeping beggarweed	Momingglory, Mexican	Sowthistle, annual	Yarrow, common
Creeping Jenny (creeping Charlie)	Momingglory red	Sowthistle, spiny	Yarrow, western
	Momingglory tall	Spanish needles	Yarrow, yellow
	Momingglory woolly	Speedwell, bi-lobed	

SUBLABEL 2 :

PRODUCT FACTS [optional info box]

What It Does: Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass, foxtail, and yellow nutsedge. (See inside for complete list)

Where to Use: (icon weeds)

Lawns: Kills the weeds not the lawn

When to Use:

Apply when daytime temperatures are between 45°F and 90°F.

Questions & Comments: (icon phone) (icon computer)

Call 1-800-xxx-xxxx or Visit our Website at www.companyname.com

Chemsico RTU Herbicide 1462 will control or suppress the following broadleaf weeds. A partial list is presented below:

Aster, white heath & white prairie	Dollarweed (*pennywort)	Lawn burweed	Ragweed
Bedstraw	False dandelion ('spotted catsear & common catsear)	Lespedeza, common	Redweed
Beggarweed, creeping	Field bindweed	Mallow, common	Red sorrel (*sheep sorrel)
Bindweed	(*morningglory & creeping jenny)	Matchweed	Shepherdspurse
Black medic	Field oxeye-daisy	Mouseear chickweed	Spurge
Broadleaf plantain	(*creeping oxeye)	Old world diamond flower	Thistle
Buckhorn plantain	Filaree, whitestem & redstem	Oxa/is (*yellow woodsorrel & creeping woodsorrel)	Virginia buttonweed
Bull thistle	Florida betony	Parsley-piert	White clover ("Dutch clover, honeysuckle clover, white trefoil, & purplewort)
Burdock, common	Florida pusley	Pennsylvania smartweed	Wild carrot
Buttercup, creeping	Ground ivy	Pepperweed	Wild garlic
Carpetweed	Groundsel	Pigweed	Wild geranium
Chickweed, common	Hawkweed	Pineappleweed	Wild lettuce
Chicory	Healall	Plantain	Wild mustard
Cinque/oil	Henbit	Poison ivy	Wild onion
Clover	Innocence (Blue-eyed Mary)	Poison oak	Wild strawberry
Curly dock	Knotweed	Prickly lettuce ('compass plant)	Yarrow
Dandelion	Lambsquarters	Puncturevine	Yellow rocket
Dayflower		Purple cudweed	and many more broadleaf weeds
Deadnettle		Purslane	
Dock			
Dogfennel			
*Synonyms			

NOTICE: To the extent allowed by applicable law buyer assumes all responsibility for safety and use not in accordance with directions.

If you are not satisfied with this product for any reason, send your written request for a refund to the address at left, together with the original dated store receipt showing the price you paid. Please allow 6 to 8 weeks for processing.

Questions or comments? Call toll free 1-800-xxx-xxxx.

[telephone number, address, website information for supplemental registrant or distributor].

Chemsico, A Division of United Industries Corp
P.O. Box 142642
St.Louis MO 63114-0642

EPA Reg. No 9688-280

EPA Est No 9688-M0-1

©2024