66-290

23 2012 2



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

FEB 2.3 2012

Jessica Goldberg Chemsico Division of United Industries Corp. P.O. Box 142642 St. Louis, MO 63114

Subject: Notification per PR Notice 98-10 – Add Optional Marketing Statements Chemsico Pesticide Concentrate WI-S EPA Reg. No. 9688-290 Application Dated – February 3, 2012

Dear Ms. Goldberg:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the subject product.

The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been date-stamped "Notification" and will be placed in our records.

If you have any questions regarding this letter, please contact Maggie Rudick at (703) 347-0257 or <u>rudick.maggie@epa.gov</u>.

Sincerely,

Kathryn Montague, Product Manager 23 Herbicide Branch Registration Division (7505P)

, 1))	2099
			d and a second s	
Please read instructions on rev	United States Environmental Protection Office of Pesticide Programs (H750 Washington, DC 20460	n Agency D5C)	OMB No. 2070-0060. App Registration	OPP Identifier Number
	Application for Pesti] Other	
1. Company/Product Number	Section	2.Product Manager	3. Proposed Classifica	ation
9688-290		K. Montague	5. Floposed Classifica	
4. Company/Product (Name)		PM#	X None	Restricted
Chemsico Pesticide Conce	ntrate WI-S	23		
5. Name and Address of Applicant Chemsico, Div. Of United In P.O. Box 142642 St. Louis, MO 63114	(include ZIP Code)		I accordance with FIFRA lentical in composition	
Check if this is a r	new address	Product Name		······
	Section	11		
Amendment - Explain be Resubmission in respons Notification - Explain be	se to Agency letter dated	Final printed label in resp "ME TOO" Application. X Other - explain below	onse to Agency letter date	31
Explanation: Use additional page Label notification per PRN 98-10 t	to add optional marketing claims (see cover letter for details).	(Certificati On co	on statement orer letter)
1. Material in Which This Product W	Section Section			
Child-Resistant Packaging Yes * No No Submitted	Unit Packaging Yes No If "YES," No. per Unit package wgt container	Water-Soluble Packaging Yes No If "YES," No. per Package wgt container	2. Type of Container Metal Plastic Glass Paper Other (Sp	ecify)
3. Location of Net Contents Informatic	n 4. Size(s)	of Retail Container	5. Location of Label D	Directions
Label	Container		On Label On Labeling acco	ompanying product
6. Manner in Which Label Is Affixed T	Paper glu Stenciled	led	Other	
1. Contact Point (Complete items	Section directly below for identification of the		cessant to process this	application
Name	Title		Telephone No. (incl	
Jessica Goldberg	Regulato	ory Affairs Specialist	314-683-2438	33 3 7 3 8 3 3 3
	Certification have made on this form and all attac ingly false or misleading statement n		isonment or	6 Date Application Received (Stamped)
4. Typed Name		5. Date		د د د د د د د د
Jessica Goldberg EPA Form 8570-1 (Rev.12-90)	Previous editions are ob	February 3, 2012	ile Copy (original)	Yellow - Applicant copy

Chemsico

Division of United Industries P.O. Box 142642 St. Louis, MO 63114-0642

February 3, 2012

Document Processing Desk (NOTIF) Office of Pesticide Programs United States Environmental Protection Agency Room S4900, One Potomac Yard 2777 S. Crystal Drive Arlington, VA 22202-4501

- Attention: Ms. Kathryn Montague Product Manager Team 23
- Subject: NOTIFICATION: Label Notification per PRN 98-10 Chemsico Pesticide Concentrate WI-S EPA Reg. No. 9688-290

Enclosed please find our notification per 98-10 to add optional marketing claims. All changes have been highlighted on the enclosed label.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA **NOTIFICATION**

FEB 2 3 2012

The following documents are enclosed in support of this submission:

- 1. Application for Pesticide (EPA Form 8570-1)
- 2. One copy of the proposed label with changes highlighted
- 3. One clean copy of the proposed label

Please call me at 1-800-242-1166, Ext. 2438 or via email: jessica.goldberg@spectrumbrands.com, if you have questions or need further information. Thank you in advance for your time.

Sincerely,

Jessica Goldberg Regulatory Affairs Specialist



Chemsico Pesticide Concentrate WI-S



Active Ingredients:	
* 2,4-D, dimethylamine salt	2.37%
†Mecoprop-p, dimethylamine salt	
‡Dicamba, dimethylamine salt	0.26%
Gamma-cyhalothrin	0.08%
Other Ingredients	


* 0.167 lb. 2,4-D acid equivalent per gallon or 1.96%. † 0.040 lb. Mecoprop-p acid equivalent per gallon or 0.470%. ‡ 0.018 lb. Dicamba acid equivalent per gallon or 0.217%. Contains the single isomer form of Mecoprop-p.

KEEP OUT OF REACH OF CHILDREN

CAUTION See back [panel] [booklet] for additional precautionary statements

NET CONTENTS 32 FL. OZ. (1 QUART)

[OPTIONAL PRODUCT FACT BOX]



*Do not use on Foratam variety of St. Augustinegrass in Florida.

[STOP. READ ALL INSTRUCTIONS INSIDE BEFORE USING.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING.



Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Do not allow people or pets to enter the treated area until sprays have dried.

HOW TO MIX

Choose from one of the application methods listed below.

[Pump-Up][Pressurized] Sprayer or Trigger Bottle Sprayer

- 1. Start with a clean sprayer.
- 2. Add the amount of concentrate specified below.
- 3. Replace sprayer top and shake well before spraying.
- 4. For pressure sprayers pump to pressurize according to sprayer instructions

DIAL-STYLE HOSE END SPRAYER

- 1. Start with a clean sprayer.
- 2. Set sprayer to the "fl oz/gallon" level specified below.
- 3. Use only enough product for the area to be treated
- 4. Close sprayer and attach to hose prior to application

CONCENTRATE PLUS WATER-STYLE SPRAYER

- 1. Start with a clean sprayer.
- 2. Determine the amount of concentrate required below.
- 3. Pour concentrate into sprayer jar to the required "fl. oz." level.
- 4. Then add the required amount of water to the "gals." level.
- 5. Replace sprayer top on jar and shake well before spraying.

MEASUREMENT CONVERSIONS

1 Tablespoon = 3 teaspoons, 1 fl. oz. = 6 teaspoons or 2 tablespoons, 6 fl. oz. = $\frac{3}{4}$ cup

1 ½ Tablespoons in 3 gallons of water = 1½ teaspoons in 1 gallon of water

2 Tablespoons in 3 gallons of water = 2 teaspoons in 1 gallon of water

SURFACE INSECTS: ants (Fire, Argentine, Carpenter, Southern, Field, Allegheny mound, Cornfield, Honey, Pavement, Red harvester, Odorous, Pyramid, Black turf, White-footed, Crazy, Little black, Small honey, Ghost, Thief, Acrobat, Citronella and Big-headed), armyworms (Fall, Yellow-striped), cockroaches, crickets (House, Common short-tailed, Indies short-tailed, Arizona, Tropical, Variable field, Fall field, Southeastern field, Texas field, Spring field, Northern wood, Western striped, Eastern striped, Sand field, Vocal field, Southern wood) cutworms (Black, Bronzed, Granulate, Variegated), earwigs, fleas, Iawn moths (sod webworms), millipedes, mosquitoes, palmetto bugs, sowbugs, mites, spittlebugs, and waterbugs: Thoroughly wet down grass a few hours before applying. Mix 1.5 fl.oz. (3 Tablespoons) in 1 gallon of water to cover [treat] 240 sq. ft. [1 Qt of Chemsico Pesticide Concentrate WI-S will treat 5120 sq ft.] Delay additional watering or mowing for 24 hours after application for optimum control. Use this product on St. Augustine*, Bahia, Centipede, Bermuda, Bentgrass, and zoysia varieties only.

This application will also kill broadleaf weeds that have emerged at the time of application. See enclosed Table for list of weeds controlled.

WHEN TO TREAT: Treat when insects are present in the area to be treated.

*Do not use on Floratam variety of St. Augustinegrass in Florida.

USE PRECAUTIONS

- 1) For best results, mow before treating weeds.
- 2) For new lawns, treat within 24 hours after seeding. On established lawns, wait 3 weeks after treatment before re-seeding. Once new grass has started to grow, wait until it has reached a height of 2 inches before treating;
- 3) Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage other plants.

USE RESTRICTIONS

- Do not treat when air temperatures exceed 85° F or damage to grass may occura
- Do not water within 6 hours after treatment. Poor weed control may result if rainfall occurs within 6 hours of application.
- Do not allow spray to get on vegetables or ornamental plants since injury may result. Do not spray around roots of trees and ornamentals. Clover will be damaged if sprayed.
- Do not make more than 2 applications per year with a minimum retreatment interval of 30 days.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not water the treated area to the point of run-off
- Do not make applications during rain.
- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 - 1. Treatment to soil or vegetation around structures
 - 2. Applications to lawns, turf, and other vegetation
- All outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

In New York State, this product may not be applied to lawns within 100 ft of a coastal marsh, or stream that drains directly into a coastal marsh.

OPTIONAL READY SPRAY NOZZLE DIRECTIONS

STOP. READ ENTIRE LABEL BEFORE USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Do not allow people or pets to enter the treated area until sprays have dried.

HOW TO APPLY

Determine the size of the spraying job. For lawns measure the area to be sprayed. For trees, shrubs and flowers, apply as a thorough cover spray. Begin treating when insects first appear. Applications should be repeated only as directed to maintain control. Apply as directed [in the rate chart]. Remember, thorough coverage is important. Be sure to follow the important use limitations for the specific area you are treating, including the amount of spray that can be applied.

USING THE READY SPRAY NOZZLE (Optional subheading)

- 1. Shake container well before using.
- 2. Connect a garden hose to the Ready Spray nozzle. Make sure the dial on the nozzle is in the "OFF" position with the [green] safety tab in the valve notch.
- 3. Turn on water at faucet. Extend hose to the farthest area to be treated and work back toward the faucet so you don't come in contact with the treated area.
- 4. To BEGIN spraying, point nozzle toward treatment site and a.) bend the [green] safety tab back (located at right of yellow dial) with your thumb, b.) hold while turning yellow dial clockwise until it stops. Water will automatically mix with the product.
- 5. Spray until wet to control insects. Walk at a steady pace while spraying using an even sweeping motion, slightly overlapping treated areas.
- 6. To STOP spraying, QUICKLY turn the dial counter clockwise until it stops and the safety tab engages into the notch on the valve. Turn water off at faucet. To relieve pressure before removing nozzle from hose, bend the safety tab back and turn dial "ON" until water stops spraying.
- 7. To STORE unused product, make sure the dial is in the "OFF" position with [green] safety tab in the valve notch. Place in cool area away from heat, sunlight or open flame.

SURFACE INSECTS: ants (Fire, Argentine, Carpenter, Southern, Field, Allegheny Mound, Cornfield, Honey, Payement, Red Harvester, Odorous, Pyramid, Black Turf, White-footed, Crazy, Little Black, Small Honey, Ghost, Thief, Acrobat, Citronella and Big-headed), armyworms (Fall, Yellow-striped), cockroaches, crickets (House, Common short-tailed, Indies short-tailed, Arizona, Tropical, Variable field, Fall field, Southeastern field, Texas field, Spring field, Northerr, wood, Western striped, Eastern striped, Sand field, Vocal field, Southern wood) cutworms (Black, Bronzed, Granulate, Variegated), earwigs, fleas, lawn moths (sod webworms), millipedes, mosquitoes, palmetto bugs, sowbugs, mites, spittlebugs, and waterbugs:

Thoroughly wet down grass a few hours before applying. 1 Qt of Chemsico Pesticide Concentrate WI-S will treat 5120 sq ft. Delay additional watering or mowing for 24 hours after application for optimum control. Use this product on St. Augustine*, Bahia, Centipede, Bermuda, Bentgrass and zoysia varieties only

This application will also kill broadleaf weeds that have emerged at the time of application. See enclosed Table for list of weeds controlled.

WHEN TO TREAT: Treat when insects are present in the area to be treated.

*Do not use on Floratam variety of St Augustinegrass in Florida

USE PRECAUTIONS

- 1) For best results, mow before treating weeds.
- 2) **For new lawns,** treat within 24 hours after seeding. **On established lawns,** wait 3 weeks after treatment before re-seeding. Once new grass has started to grow, wait until it has reached a height of 2 inches before treating.
- 3) Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage other plants.

USE RESTRICTIONS

- Do not treat when air temperatures exceed 85° F or damage to grass may occur.
- Do not water within 6 hours after treatment. Poor weed control may result if rainfall occurs within 6 hours of application.
- Do not allow spray to get on vegetables or ornamental plants since injury may result. Do not spray around roots of trees and ornamentals. Clover will be damaged if sprayed.
- Do not make more than 2 applications per year with a minimum retreatment interval of 30 days.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff

6000

or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only.

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not water the treated area to the point of run-off
- Do not make applications during rain.
- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 - 3. Treatment to soil or vegetation around structures
 - 4. Applications to lawns, turf, and other vegetation
- All outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

In New York State, this product may not be applied to lawns within 100 ft of a coastal marsh, or stream that drains directly into a coastal marsh.

----- END OF OPTIONAL READY SPRAY NOZZLE DIRECTIONS ------

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a cool, dry area. Always store pesticides in the original container. Store away from food and pet food. **Pesticide Disposal and Container Handling**: Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling, if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID

If swallowed:	 Call a Poison Control Center or doctor immediately for treatment advice. 	
	 Have person sip a glass of water if able to swallow. 	
	 Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by 	

	mouth to an unconscious person.
lf in Eyes:	 Hold eye open and rinse slowly and gently with water for 15-29 minutes. Remove contact lenses, if present, gently after the first 5 minutes, then continue rinsing eye.
	Call a Poison Control Center or doctor Tor treatmant advice.
Have the product containe or doctor, or going for treat	r or label with you wijen calling a Poison Control Center tment. You may also contact 1-80ງ-zzz-zzzz for

emergency medical treatment information.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect nontarget plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

Questions or comments? Call 1-800-332-5553

[*If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.]

Made in the USA for Chemsico Division of United Industries Corporation P.O. Box 142642, St. Louis, MO 63114-0642

EPA Reg. No. 9688-290 EPA Est. No. 9688-290 © 2011

R

Chemsico Pesticide Concentrate WI-S EPA Reg. No. 9688-290 FN: 9688-290-FPL-N1-Highlighted.doc

Optional Marketing Claims

- 1. [Kills][Controls] the weeds, not the lawn
- 2. [Kills][Controls] 200+ listed weed types to the root
- 3. [Kills][Controls] 200+ listed weed types, including the roots
- 4. [Kills][Controls] 200+ listed weed types [including dandelion, chickweed, wild onion, clover and poison ivy]
- 5. [Kills][Controls] Dandelion, Chickweed, Plantain, Wild onion. Poison lvv land other listed weeds]
- 6. [Kills][Controls] even the toughest broadleaf weeds [as listed] [including clover, dandelion and oxalis]
- 7. [Kills][Controls] hard to kill broadleaf weeds [as listed] [including clover, dandelion, oxalis]
- 8. [Kills][Controls] Broadleaf Lawn Weeds [and insects] as listed
- 9. Effective on hard to kill weeds [as listed]
- 10. Treats 5120 sq ft [for weed and insect control]
- 11. [See] Visible results in hours
- 12. Controls tough lawn [weeds][insects][weeds and insects] as listed
- 13. [Kills][Controls] Insects [as listed] in 24 hours
- 14. Dual action formula
- 15. 2 in 1 [weed and insect control] lawn treatment
- 16. Guaranteed Results or your money back*
- 17. Money Back Guarantee*
- 18. Apply year Round [when temperatures are below 85°F]
- 19. See Results in 24 Hours
- 20. Rainfast in 3 Hours
- 21. Rainproof in 3 Hours [Won't wash away]
- 22. Dilutable Concentrate
- 23. For home lawn care
- 24. Non-staining
- 25. Low Odor
- 26. For Use on Home Lawns
- 27. For use on Southern Grasses
- 28. Won't harm Southern grasses
- 29. Protects lawns from listed [weeds][insects][weeds & insects]
- 30. 2 Products in 1 for [easy][time-saving] lawn treatment
- 31. Lawn Weed and Insect Control [in 1 Formula]
- 32. [Kills][Controls] up to 55 Listed insects by contact
- 33. [Kills][Controls] over 50+ listed insects
- 34. [Kills][Controls] 50+ listed insects in 24 hours
- 35. For use on Bahia, Bentgrass, Bermuda, Centipede, St. Augustine* and Zoysia lawns



د ر ر ر	ں پ	<u>ب</u> ن
2222	ر ر.	ت ن
-		ر ن
ر ب ب		د ب ر ر
ູ ມ	5	
و ر ر	دَ	

EPA Reg. No. 9688-290 FN: 9688-290-FPL.doc

5/27/11

9099

- BROADLEAF WEEDS -						
Annual yellow sweetclover	Cockle	Musk thistle	Spiny amaranth			
Aster	Cocklebur	Mustard	Spiny cocklebur			
Austrian fieldcress	Common mullein	Narrowleaf plantain	Spiny sowthistle			
Bèdstraw	Creeping Jenny (Creeping Charlie)	Narrowleaf vetch	Spotted catsear			
Boggarticks 📄 🦷	Cudweed	Nettle	Spotted spurge			
Betony, Florida	Curly dock	Orange hawkweed	Spurweed			
Bindweed, field	Daisy, English	Oriental cocklebur	Stinging nettle			
Bird vetch	Daisy fleabane	Oxalis	Strawberry, India mock			
Bitter wintercies	Daisy, oxeye	Parsley-piert	Tall nettle			
Bitteroress, hairy	Dandelion	Parsnip	Tall vervain			
Bitterweed	Dichondra	Pearlwort	Tansy ragwort			
Black-eyed Susan	Dogbane	Pennycress	Tansy mustard			
Black medic	Dogfennel	Pennywort	Tanweed			
Black mustard	Dollarweed	Peppergrass	Thistle			
Blackseed plantain	Elderberry	Pepperweed	Trailing crownvetch			
Blessed thistle	False dandelion	Pigweed	Tumble mustard			
Bloodflower milkweed	Falseflax	Pineywoods bedstraw				
Blue lettuce	False sunflower	Plains coreopsis (tickseed)	Tumble pigweed Velvetleaf			
Blue vervain			Venice mallow			
	Fiddleneck	Plantain Deises inte				
Bracted plantain	Florida pusley	Poison ivy	Virginia buttonweed			
Brassbuttons	Frenchweed	Poison oak	Virginia creeper			
Bristly oxtongue	Galinsoga	Pokeweed	Virginia pepperweed			
Broadleaf dock	Goathead	Poorjoe	Water pennywort			
Broadleaf plantain	Goldenrod	Prairie sunflower	Wavyleaf bullthistle			
Broomweed	Ground ivy	Prickly lettuce	Western clematis			
Buckhorn	Gumweed	Prickly sida	Western salsify			
Buckhorn plantain	Hairy fleabane	Prostrate knotweed	White mustard			
Bulbous buttercup	Hawkweed	Prostrate pigweed	Wild aster			
Bull thistle	Healall	Prostrate spurge	Wild buckwheat			
Bullnettle	Heartleaf drymary	Prostrate vervain	Wild carrot			
Burclover	Heathaster	Puncturevine	Wild four-o'clock			
Burdock	Hedge bindweed	Purslane, common	Wild garlic			
Burning nettle	Hedge mustard	Ragweed	Wild geranium			
Burweed	Hemp	Red sorrel	Wild lettuce			
Buttercup	Henbit	Redroot pigweed	Wild marigold			
Buttonweed	Hoary cress	Redstem filaree	Wild mustard			
Canada thistle	Hoary plantain	Rough cinquefoil	Wild onion			
Carolina geranium	Hoary vervain	Rough fleabane	Wild parsnip			
Carpetweed	Horsenettle	Russian pigweed	Wild radish			
Catchweed bedstraw	Jimsonweed	Russian thistle	Wild rape			
Catnip	Knawel	Scarlet pimpernel	Wild strawberry			
Catsear	Knotweed	Scotch thistle	Wild sweet potato			
Chickweed, common	Kochia	Sheep sorrel	Wild vetch			
Chickweed, mouseear	Lambsquarters	Shepherdspurse	Woodsorrel			
Chicory	Lespedeza	Slender plantain	Woolly croton			
Cinquefoil	Mallow	Smallflower galinsoga	Woolly morningglory			
Clover, crimson	Matchweed	Smooth dock	Woolly plantain			
Clover, hop	Mexicanweed	Smooth pigweed	Wormseed			
Clover, red	Milk vetch	Sorrel	Yarrow			
Clover, strawberry	Morningglory	Sowthistle	Yellow rocket			
Clover, sweet	Mouseear hawkweed	Spanishneedles	Yellowflower pepperweed			
Clover, white	Mugwort	Speedwell	Cranesbill			
Deadnettle	English Daisy	Fleabane	Fillaree			
Soliva	Wild Violet		and other broadleaf weed			

Page 6 of 6

,

.