

ADDITIONAL CAUTIONS

READ ENTIRE LABEL BEFORE USING THIS PRODUCT:
Use of this material in a manner, or at any time other than in accordance with the directions, may produce plant injury, excessive residues, or other undesirable results.

CAUTION: Avoid contact with the skin. Avoid breathing spray mist. Keep away from children and domestic animals. Avoid contamination of feed and foodstuff. Keep animals off treated areas. Wash hands after using. Harmful if swallowed.

ANTIDOTE: If taken internally, induce vomiting and call a physician at once.

Rinse empty container thoroughly with water and discard it.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

MANUFACTURED BY:
GRO-LYFE, a division of
SUNGRO CHEMICALS INCORPORATED
P.O. BOX 24632 LOS ANGELES, CALIFORNIA 90024

SUNGRO

LIQUID DICHONDRA AND LAWN FOOD WITH

CRABGRASS KILLER

Kills The Pest — Feeds The Rest

COVERS 1,000 SQUARE FEET

INGREDIENT STATEMENT

As a Plant Food:
ORGANIC NITROGEN, Derived from Urea10%
PHOSPHORIC ACID 0%
POTASH0%

As a Herbicide:
ACTIVE INGREDIENT:
Monosodium Acid Methanearsonate 0.52%
INERT INGREDIENTS 99.48%
TOTAL 100%

TOTAL ARSENIC, all in water soluble form,
expressed as elemental 0.24%


E.P.A. Reg. No. 9754-2 E.P.A. Estab. No. 11474CA1

CAUTION:

KEEP OUT OF REACH OF CHILDREN

(SEE SIDE PANEL FOR ADDITIONAL CAUTIONS)

NET CONTENTS:.....GAL.



**A COMBINATION PLANT FOOD AND
CRABGRASS KILLER**



DIRECTIONS: When using 80 to 1 hose type fertilizer gun, use full strength. Moisten soil before and after application. Apply at the rate of one (1) gallon for each 1000 sq. ft. Hose type fertilizer gun dilutes at proper rate of dilution.

DICHONDRA AND GRASS LAWNS: Use at rate of one (1) gallon to 1000 sq. ft. ratio with gun type applicator. Apply Sungro Crabgrass Killer to lawns, dichondra or for the control of crabgrass and other annual grasses such as foxtail, witch grass and barnyard grass, Johnson grass, Dallis grass and goose grass. Avoid treating during hot weather (above 90 degrees F.).

Repeat at seven day intervals for one month. Fertilizer content will feed desirable grasses, while crabgrass turns brown and dies. If new crabgrass appears due to old seed germination, repeat application. This product will not permanently injure desirable grasses or dichondra. Newly seeded lawns should not be treated until well established. Don't use on St. Augustine grass

PRODUCT OF THE U.S.A.

