

T-CHEM

PARATHION E LB.
Emulsion Concentrate

Contains 8 lbs. of Parathion per gallon.

**DANGER: KEEP OUT OF REACH OF CHILDREN.
READ ALL DIRECTIONS CAREFULLY BEFORE USING.
SEE ANTIDOTE STATEMENT AND OTHER WARNINGS ON THE BACK/SIDE PANELS.**

X
P O I S O N
X

ACTIVE INGREDIENTS:

Parathion O,O-diethyl O-p-nitrophenyl thiophosphate..... 80.02
Xylene..... 5.82

INERT INGREDIENTS:

..... 14.24
TOTAL 100.08

NET CONTENTS:

USDA REG. /

TRITCHER CHEMICAL COMPANY
Salt Lake City, Utah

For control of aphids, spider mites, lygus bugs and alfalfa weevil on alfalfa,
grasshoppers and army worms - mix 1/4 pint of Parathion Emulsion Concentrate (4
ounces of Parathion) with sufficient water to properly cover the crop. Treated by
the methods used with high volume sprayers this will be approximately 100 gallons
of water, low volume sprayers 20 gallons, and aircraft from 2 to 12 gallons per
acre.

On small grains, such as, wheat, oats, barley for control of mites and grasshoppers,
one or more applications may be necessary, but none should be applied after the heads
begin to form.

On alfalfa and clover for control of aphids, spider mites, grasshoppers, and army
worms, no application should be made later than 15 days prior to cutting or pasturing.

of
On field peas for control of aphids, one or more applications may be necessary, but
none later than 21 days prior to harvest.



ACCEPTED
JUN 13 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
AND ECONOMIC POISON REGISTERED
ED UNDER NO. 9713-23 SUBJECT
TO ATTACHED COMMENTS

DANGER: POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN.
Repeated exposures may be increasingly hazardous. Rapidly absorbed through the skin.
Do not breathe vapors.
Do not get into eyes, on skin or on clothing. If spilled on skin, immediately wash thoroughly with soap and water.
Wear protective clothing, natural rubber gloves and goggles.
When filling spray tanks and during mixing and application wear a respirator passed by the U.S. Dept. of Agriculture for Parathion protection. Airplane pilots should wear full-face canister type masks.
Wash hands, arms and face thoroughly with soap and water before eating or smoking.
Bathe promptly with soap and hot water after spraying.
Wash clothing with soap and hot water daily before reuse.
Keep all persons out of the operating area or vicinity where there may be danger of drift.
Vacated areas should not be re-entered until insecticidal residue has dissipated.
This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on the label.
DO NOT STORE NEAR NOR CONTAMINATE FEED OR FOODSTUFFS!

DANGER: Keep away from heat or open flame!
Destroy the container when empty. Wash out thoroughly with water. Puncture full of holes or crush so that it cannot be used for any purpose. Dispose of it where it cannot be salvaged for any purpose.

ATROPINE IS ANTIDOTAL - KEEP AN EMERGENCY SUPPLY ON HAND. CALL A PHYSICIAN AT ONCE.

FIRST AID TREATMENT: DO NOT WAIT FOR A DOCTOR. Give 2 Atropine tablets at once if patient has blurred vision or pin-point pupils.

INTERNAL: If swallowed, give tablespoon of salt in warm water and repeat until vomit fluid is clear.

EXTERNAL: If spilled on body, immediately remove contaminated clothing and wash skin thoroughly with soap and water.

INHALATION: Remove patient from contaminated area and keep under continual observation until doctor arrives.

NOTE TO PHYSICIANS: Warning symptoms include weakness, headache, chest pain, blurred vision, pin point pupils, rabid salivation, sweating, nausea, vomiting, diarrhea, and cramps.

TREATMENT: Give atropine, preferably by injection, grains 1/100 three or more tablets at once and parenterally or orally every hour until pupils dilate. Thirty or more tablets may be required in the first 24 hours. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may without warning cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.

Parathion Emulsion Concentrate is for use only by thoroughly trained operators familiar with the handling of hazardous insecticides and is to be applied only by aircraft and power sprayers.

