RIVERSIDE, LADUSTRIES

Marks; Mississippi

MARKET 194 1945 FAR 74

AUG 23 1967

RAVERSIDE CHEMICAL COMPANY

Sico)

Champion Chemical Company

APORE/AMERICAN CARMAN

IN YOR YOUR ALL

፲ﯘﯨﭙﯘﭘﻼ*ﻟﯩﺪﻩ ﺳﯘ*ﻻ-Დ₭₭₭₭

55% PREMIUM GRADE MALATHION Contains a pounds actual Malathion per gallon

ACTIVE INGREDIENTS:
Malathion*
INERG INGREDIENTS:

55% 15%

TOTAL

__15% 100%

*O, O-Dimethyl Dithiophosphate of Diethyl Mercaptosuccinate.
CAUTION: KEEF OUT OF REACH OF CHILDREN. SEE OTHER CAUTIONS ON RIGHT PANEL.
DIRECTIONS

STORED GRAIN - For the protection of stored graves such as wheat, oats, rice, corn, rye, barley, grain sorghum, peanuts, and field or garden seeds, against confused flour beetle, rice weevil, granary weevil, sawtoothed grain beetle, flat grain beetle, red flour beetle, rusty grain beetle, lesser grain borer, and Indian meal moth, apply 1 pint of premium grade Malathion 55% emulsifiable liquid in 2 to 5 gallons of water per 1,000 bushels. Apply as the grain is being loaded or turned into final storage.

Field and garden seeds can be protected against the above grain pests with a dosage of $\frac{1}{2}$ pint of Malathion premium grade 55% emulsifiable liquid in 1 to $2^{1}{2}$ gallons of water per 500 bushels of seed.

To protect stored grains from attack by Indian Meal Moth, apply premium grade Malathion 55% emulsifiable liquid to the surface of clean or uninfested grain at the rate of ½ pint in 1 to 2 gallons of water per 1000 square feet of grain surface area. Apply the spray evenly over the surface of the grain. Apply immediately after grain is loaded into storage and repeat if necessary.

For a residual wall, floor, and machinery spray in grain elevators, in treating truck beds, box cars, and ships holds before loading grain, apply I gallon of premium grade Malathion 55% emulsifiable liquid per 25 gallons of water making thorough application. Refore applying spray, clean elevators, box cars, etc. thoroughly. Remove and burn all sweepings and debris.

Malathion may be used under the requirements of the Khapra heetle quarantine, where water or diesel oil emulsions are indicated, as prescribed by the current quarantine instructions.

FLIES: For use in and around buildings which house domestic animals, use 2 gallons in 100 gallons of water and apply at the rate of 1 to 2 gallons per $1,000~\mathrm{su}$, ft, where flies alight or congregate. Do not use in milk processing rooms.

CATTLE & SWINE: Lice and ticks on beef animals, and lice on swine - Use I gallon in 100 gallons of water. Apply a complete coverage spray.

AGRICULTURAL

CHEMICALS

RIVERSIDE CHEMICAL COMPANY



Champion Chemical Company

Marks, Mississippi Beale

イズ 大イグン ない / しゅんんこれ

おおおり まなか まみのれてはずじ

Repeat applications at 2-week intervals if needed. Do not use on dairy anim, 's. Do not treat calves under one month of age. When applying sprays, avoid contamination of feed, food containers, and watering troughs.

CAUTION

Harmful if swallowed, avoid prolonged breathing of spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoil contamination of feed and foodstuffs.

Highly toxic to fish and wildlife. Keep out of lakes, streams and ponds. Do not apply when conditions favor drift from the crop area treated. Do not use for any purpose other than stated on this label.

Note to Physicians: Malathion, upon repeated, prolonged or careless use, may cause cholinesterase inhibition. Atropine is antidotal.

Rinse equipment and containers and dispose of wastes or spillage in non-crop lands away from water supplies. Containers should be disposed by peaching holes in them and burying with wastes.

FLAMMABLE: KEEP CONTAINER CLOSED AND AWAY FROM HEAT AND OPEN FLAME.

NET CONTENTS: 1, 5, 30 and 55 Gallons

USDA REG. NO. \$\$77\&\$ 977' -

Manufactured by: CHAMPIONY CHEMICALY COMPANY RIVERSIDE CHEMICAL COMPANY

CARTONY MISSISSIPPI TIRKS, MISSISSIPPI