


LLON.

18 15%
22 40%
54 50%
4 50%
00 0%

GI
ER
OF
H OF
EN



ATED
if vapors or
exposed to
cups. Protec-
from your
Service

GENERAL INFORMATION

RIVERSIDE 162 is a systemic insecticide for control of sucking insects and other pests of cotton, tobacco, and other crops.

RIVERSIDE 162 is a systemic insecticide for control of sucking insects and other pests of cotton, tobacco, and other crops. It is a colorless, odorless liquid which is easily absorbed by the plant and moves through the vascular system to the leaves and stems.

When used as a systemic insecticide, RIVERSIDE 162 should be applied to the soil around the base of the plant. It is absorbed by the roots and moves up through the stem to the leaves and stems. It is effective against a wide range of sucking insects, including aphids, beet beetles, and whiteflies.

When used as a contact insecticide, RIVERSIDE 162 should be applied to the leaves and stems of the plant. It is effective against a wide range of sucking insects, including aphids, beet beetles, and whiteflies.

For control of sucking insects on cotton, tobacco, and other crops, RIVERSIDE 162 should be applied to the soil around the base of the plant. It is absorbed by the roots and moves up through the stem to the leaves and stems.

For control of sucking insects on cotton, tobacco, and other crops, RIVERSIDE 162 should be applied to the leaves and stems of the plant. It is effective against a wide range of sucking insects, including aphids, beet beetles, and whiteflies.

DIRECTIONS FOR USE

COTTON
Apply RIVERSIDE 162 to the soil around the base of the plant. It is absorbed by the roots and moves up through the stem to the leaves and stems. It is effective against a wide range of sucking insects, including aphids, beet beetles, and whiteflies.

SMALL GRAINS (wheat, oats, rye, barley)
Apply RIVERSIDE 162 to the soil around the base of the plant. It is absorbed by the roots and moves up through the stem to the leaves and stems. It is effective against a wide range of sucking insects, including aphids, beet beetles, and whiteflies.

Do not apply within 45 days of harvest. Do not feed threshings to livestock. Do not feed threshings to livestock.

DISPOSAL OF CONTAINER AND WASTES

This container should not be reused for ANY purpose. When empty, it should be rendered unusable by punching large holes in the top, bottom and sides with an instrument such as an axe. It should then be placed as far as possible and buried underground as deep as practical, with at least 18 inches of soil covering. This should be done in an isolated area away from habitation, and in a location which would not drain into, or cause pollution of, lakes, streams, or ponds.

Any spray waste or spillage resulting from the application of this material or the destruction of the container should be neutralized with a caustic solution and buried in a pit with the disposed container.

NOTICE

Riverside Chemical Company warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to Riverside Chemical Company, and buyer assumes the risk of any such use.


EPA REG. NO. 97796

NET CONTENTS



RIVERSIDE CHEMICAL CO.

Riverside serves agriculture. Agriculture serves everyone.



METHYL PARATHION MIXTURE, LIQUID