

# RIVERSIDE TOXAPHENE 6

AN EMULSIFIABLE CONCENTRATE CONTAINING 6 POUNDS OF TOXAPHENE PER GALLON.

USEFUL FOR THE CONTROL OF CERTAIN INSECT PESTS OF BEANS (Dry, Green, Lima), CABBAGE AND OTHER COLE CROPS, CORN, COTTON, COWPEAS, OKRA, PEAS, PEANUTS, PIMENTOS, CERTAIN SMALL GRAINS (Barley, Oats, Rice, Rye, Wheat), SORGHUM, SOYBEANS, SPINACH, STRAWBERRIES, AND TOMATOES.

#### ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene-chlorine content 67% to 69%)	58.7%
Aromatic Petroleum Solvent	38.3%
INERT INGREDIENTS:	3.0%

TOTAL 100.0%

Contains 6 pounds of Toxaphene per gallon

## WARNING: KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed. Do not get on skin, in eyes, or on clothing. Can be absorbed through skin. Wash thoroughly after using. Do not breathe spray mist.

Avoid contamination of feed or foodstuffs. Do not use on dairy animals or in dairy barns. Do not use on cucurbits.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the areas.

#### ANTIDOTE

INTERNAL: Give sodium chloride 15 g. (one tablespoonful of table salt) in glass of warm water and repeat until vomit is clear. Call a doctor immediately.

EXTERNAL: Wash with soap and water.

#### DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

**READ ENTIRE LABEL BEFORE USING THIS PRODUCT:** Use of this product in a manner not in accordance with the information on this label may injure or kill humans, domestic animals, wildlife, aquatic life, desirable plants, create excessive and unlawful chemical residues, or produce other undesirable results.

#### GENERAL INFORMATION

RIVERSIDE TOXAPHENE 6 is useful for the control of certain insect pests of beans (dry, green, lima), cabbage and other cole crops, corn, cotton, cowpeas, certain small grains (barley, oats, rice, rye, wheat), okra, peas, peanuts, pimentos, sorghum, soybeans, spinach, strawberries and tomatoes.

RIVERSIDE TOXAPHENE 6 may be applied by ground sprayers or by airplanes, mixing the RIVERSIDE TOXAPHENE 6 in water to form an emulsion spray solution. To make the spray solution, half fill the spray tank with water, agitate, add the proper amount of RIVERSIDE TOXAPHENE 6 and then fill the spray tank with water while still agitating. Continue to agitate during use.

When applying with ground sprayers calibrate so as to apply the recommended amount of RIVERSIDE TOXAPHENE 6 in not less than 3 gallons of water per acre. Airplanes should apply the recommended amount of RIVERSIDE TOXAPHENE 6 in not less than 1 gallon of water per acre. Always use sufficient water for adequate coverage of the crop foliage.

Where a dosage range is given use the lower rate on small or young plants or with light insect infestations, use the higher rate on large or mature plants or with heavy insect infestations.

The hazards and precautions for handling RIVERSIDE TOXAPHENE 6 in this container are equally applicable to it after dilution with water for spray application.

#### DIRECTIONS FOR USE

**BEANS (Dry, Green, Lima):** For control of Mexican Bean Beetle and Thrips apply 1½ to 2 pints of RIVERSIDE TOXAPHENE 6 per acre per application; for control of Corn Earworms, Grasshoppers, and Serpentine Leaf Miner apply 3 to 4 pints per acre; for control of Bean Lycaenid apply 3½ to 4½ pints per acre; for control of Waxbean Beetle apply 3 to 6 pints per acre. Repeat applications as necessary. Do not apply after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter.

**CABBAGE AND OTHER COLE CROPS (Also including Broccoli, Brussels Sprouts, Cauliflower, Collards, and Kale):** For control of Armyworms, Fall Armyworms, Cabbage Aphid, and Surface Feeding Cutworms apply 3 pints of RIVERSIDE TOXAPHENE 6 per acre per application; for control of Imported Cabbageworm, Green Peach Aphid, Serpentine Leaf Miner, and Thrips apply 2½ pints per acre; for control of Corn Earworm, and Diamondback Moth apply 2 to 5½ pints per acre; for control of Cabbage Looper apply 2½ to 5½ pints per acre. Repeat applications as necessary. Do not apply after edible parts begin to form or within 35 days of harvest.

Plants treated with RIVERSIDE TOXAPHENE 6 must not be fed to dairy animals or animals being finished for slaughter.

**CORN:** For control of Chinch Bugs and Corn Billbug apply 2½ pints of RIVERSIDE TOXAPHENE 6 per acre per application; for control of Armyworm and Budworm apply 3½ pints per acre; for control of Corn Leafminer and Lesser Cornstalk Borer apply 5½ to 6 pints per acre. Repeat applications as necessary.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

**COTTON:** For control of Thrips apply 1 to 2 pints of RIVERSIDE TOXAPHENE 6 per application; for control of Fleahoppers apply 1½ to 4 pints per acre; for control of Armyworm, Beet Armyworm, Fall Armyworm, Boll Weevil, Cotton Leafworm, Cutworms, Flea Beetles, Grasshoppers, Lygus Bugs, Plant Bugs and other Mirids, and Yellow Striped Armyworm apply 1½ to 5½ pints per acre; for control of Western Yellow Striped Armyworm, Cotton Leaf Perforator, and Garden Webworm apply 5½ to 6 pints per acre. Repeat applications as necessary.

Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

**COWPEAS:** For control of Cowpea Curculio apply 2 to 4 pints of RIVERSIDE TOXAPHENE 6 per acre per application. Repeat applications as necessary. Do not apply to cowpeas to be used as green snaps after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

**OKRA:** For control of Serpentine Leaf Miner apply 2½ to 4 pints of RIVERSIDE TOXAPHENE 6 per acre per application. Repeat applications as necessary.

Do not apply after pods begin to form.

**PEAS:** For control of Lygus Bugs apply 2½ to 4 pints of RIVERSIDE TOXAPHENE 6 per acre per application. Repeat applications as necessary. Do not apply after pods begin to form if pods are to be used as food, or within 7 days if pods are not to be used as food. Do not feed vines to dairy animals or animals being fattened for slaughter.

**PEANUTS:** For control of Thrips apply 2½ to 3 pints of RIVERSIDE TOXAPHENE 6 per acre per application; for control of Fleahoppers and Leafworm apply 5½ to 6 pints per acre; for control of Velvet Bean Caterpillar apply 2 to 4 pints per acre; for control of Fall Armyworm apply 2½ to 5½ pints per acre; for control of Southern Stink Bug apply 5½ to 6½ pints per acre. Repeat applications as necessary.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

**PIMENTOS:** For control of Flea Beetles, Hornworm, Leafminers, and Pepper Weevil apply 1½ to 2½ pints of RIVERSIDE TOXAPHENE 6 per acre per application. Repeat applications as necessary. Do not apply later than 2 days prior to harvest.

**SMALL GRAINS (Barley, Oats, Rice, Rye, Wheat):** For control of Armyworms, Cutworms (Climbing and Surface), and Grasshoppers apply 2½ to 3 pints of RIVERSIDE TOXAPHENE 6 per acre per application; for control of Chinch Bugs and Rice Stink Bug apply 2½ pints per acre. Repeat applications as necessary.

**SORGHUM:** Apply RIVERSIDE TOXAPHENE 6 the same as for CORN, but do not apply more than 4 pints per acre. Do not apply later than 28 days before harvest or more than once after heads begin to form. Do not graze dairy animals or animals being finished for slaughter on treated fields. Do not ensilage treated forage.

**SOYBEANS:** For control of Armyworms, Bean Leaf Beetles, Blister Beetle, Corn Earworm (Bollworm), Grasshoppers, Green Clover Worm, and Velvet Bean Caterpillar apply 3 to 5½ pints of RIVERSIDE TOXAPHENE 6 per acre per application. Repeat applications as necessary.

Do not apply later than 21 days before harvest. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

**SPINACH:** For control of Armyworms, Fall Armyworms, Climbing and Surface Feeding Cutworms apply 3 pints of RIVERSIDE TOXAPHENE 6 per acre per application; for control of Imported Cabbageworm, Green Peach Aphid, Serpentine Leaf Miner, and Thrips apply 2½ pints per acre; for control of Diamondback Moth and Cabbage Looper apply 3 to 5½ pints per acre.

Do not apply more than once per season, or later than 21 days before harvest.

**STRAWBERRIES:** For control of Crown Borer and Strawberry Weevil apply 2½ to 4 pints of RIVERSIDE TOXAPHENE 6 per acre per application. Do not apply after fruit starts to form.

**TOMATOES:** For control of Armyworm, Blister Beetle, Cabbage Looper, Colorado Potato Beetle, Cutworms, Fleabeetles, Green Chrysanthemum Aphid, Hornworm, Pinworm, Russet Mite, Serpentine Leafminer, and Tomato Fruitworm apply 2½ to 6½ pints of RIVERSIDE TOXAPHENE 6 per acre per application. Repeat applications as necessary.

Do not apply later than 3 days before harvest.

#### DISPOSAL OF CONTAINER AND WASTES

Destroy all containers when empty by punching holes in them and burying with wastes. Do not burn. Do not reuse containers for other purposes. Rinse equipment and containers and dispose of wastes by burying in noncrop lands away from water supplies.

#### NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instruction, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA REG. NO. 9779-20

NET CONTENTS

EPA Est. 9779-AL-1,2, AR-1,2, GA-1, LA-1, MS 1,2,3, TN-1, TX-1



MANUFACTURED BY

**RIVERSIDE CHEMICAL CO.**

A SUBSIDIARY OF COOK INDUSTRIES INC., MEMPHIS, TN 38117

Riverside serves agriculture. Agriculture serves everyone.