

Effective for the control of Leaf Spot (*Cerospora Arachidicola*).

WHEN TO BEGIN

WHEN FIRST SPOTS APPEAR ON BASAL LEAVES.

NUMBER OF APPLICATIONS

USUALLY FOUR, APPLIED AT ABOUT 10-DAY INTERVALS, THE FIRST DUSTING SHOULD BE ABOUT THE TIME OF LAST CULTIVATION.

AMOUNT OF MATERIAL

FROM 20 TO 30 POUNDS OF DUST PER ACRE FOR EACH DUSTING DEPENDING ON SIZE OF PLANTS.



PEANUT FUNGICIDE Copper - Sulphur - Peanut Dust (10-90)

COMPOSITION

ACTIVE INGREDIENTS:	
Copper (From Basic Copper Sulphate) Expressed as Metallic	3.4%
Sulphur	63.0%
(Sulphur 93% passed 325 mesh)	
INERT INGREDIENTS: Not More than	
TOTAL	100.0%

WARNING

Keep Out Of Reach Of Children

Manufactured by
Agricultural Chemical Service Co.
Montgomery, Ala.

NET CONTENTS: 50 Pounds


This product is toxic to fish and wildlife. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label. Do not apply when weather conditions favor drift of spray from areas treated. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

Harmful if taken internally. ANTIDOTE: Emetic of Mustard or Hydrated Oxide of Iron and Magnesia; a cupful, followed with Olive Oil or White of EGGS, Mucilaginous. AVOID BREATHING DUST. AVOID CONTAMINATION OF FOODSTUFFS. AVOID CONTACT WITH SKIN AND EYES. In case of skin contact, wash with plenty of soap and water. For eyes, flush with plenty of water. To protect fish and wild life, do not contaminate streams, lakes, or ponds with this material.

NON-WARRANTY—Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the Risk of any such use.

Manufactured by:
AGRICULTURAL CHEMICAL SERVICE CO.
MONTGOMERY, ALA.

USDA Reg. No. 857-20



ACCEPTED
Apr. 7, 1972
 UNDER THE FEDERAL INSECTICIDE ACT
 FOR ECONOMIC CONTROL OF PESTS
 REG. NO. *1074-32*
 NO ATTACHED COMMENTS.

WOOD PRESERVATIVE

This Wood Preservative is widely used for the pressure treatments of lumber, poles and other wood products. When pressure treated lumber is not available, this Preservative can be effectively used to treat both old and new structures. When the wood is not in contact with the ground, as in the case of frames, sub-floors, sills, joists, etc. two applications of this Preservative should be made by swabbing or brushing in an erected structure. If individual members are being treated as in the case of a new building it is best to soak the wood in this Preservative for at least three minutes. All joints and ends of boards should be given thorough treatment.

Fence posts should be barked, seasoned and soaked for at least 24 hours and preferably 48 hours in a vertical position with butts downward. The best treatment is a full length coverage. If a butt treatment only is used, the posts should be treated to a length of three feet when the post is set two feet in the ground. Pines, red oak, basswood, black gum, and ash take treatment well; aspen and cottonwood do not treat well.

Forestal Wood Preservative is especially designed for the treatment of existing wooden structures or those already in place, for the protection of wood against such pests as woodrot, powder post beetles, lyctus beetles and carpenter ants. It is also an excellent soil poison for termites.

Treatment of wood after it is already in place would not be effective against termites as they tunnel in the wood below the surface in those areas not reached by the treatment.

INFLAMMABLE
 —Keep Away from Fire—

NET CONTENTS

GALLONS

TERMITE CONTROL

SO-PRO-CO

FORESTAL

WOOD PRESERVATIVE

READY TO USE

**An Effective Wood Preservative
 and Soil Poison**

ACTIVE INGREDIENTS	99.8%
†Pentachlorophenol	4.3%
†Other chlorophenols	.5%
Petroleum hydrocarbons	95.0%
†INERT INGREDIENTS	0.2%
†Equivalent to 5% Technical Pentachlorophenol (Fed. Spec. TT-W-570)	

CAUTION
KEEP OUT OF THE REACH OF CHILDREN
 See caution statement side panel

EPA REG. NO. 1074-32

SOUTHERN PRODUCTS COMPANY
 1301 Heistan Pl. S. 6th & Davidson St.
 Memphis, Tenn. Nashville, Tenn.

Dig a trench 30 inches deep all the way around foundations but do not dig below the top of the footing if this is shallower. Spray or soak the soil, which has been removed from around the foundation of the building using 1 gallon of preservative per linear foot of soil to a depth of 30 inches. Replace the treated dirt respraying at intervals as the earth is replaced. Shallow trenches (up to 15 inches) can be treated at a rate of 5 gallons forestal per 10 linear feet. Decayed or termite attacked wooden sills should be replaced with new ones. Wood preservative treated. Do not spray on plants. Grass, trees and shrubs within 3 feet of the trench should be removed or protected by Tar Paper.

NON-WARRANTY

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

WARNING: Harmful if swallowed or absorbed through skin. Causes skin irritation. Do not breathe vapor or spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling. Wear Protective clothing when handling freshly treated lumber.

In case of contact with skin, wash at once with plenty of warm water and soap.

Do not take internally, if accidentally swallowed give a teaspoon of salt in a glass of warm water and repeat until vomit fluid is clear. Give milk or egg whites. Have patient lie down. **CALL A DOCTOR.** Do not reuse container. Destroy When Empty.

FOR INDUSTRIAL USE

This product is toxic to fish and wildlife. Do not contaminate water by cleaning of equipment, or disposal of wastes.