

9777-32  
SEP 20 1978

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. \_\_\_\_\_ SUBJECT  
TO ATTACHED COMMENTS

# RIVERSIDE EPN 4

AN EMULSIFIABLE CONCENTRATE  
USEFUL FOR THE CONTROL OF  
FRUITS AND NUTS (ALMONDS,  
GRAPES, NECTARINES, PEACHES,  
WALNUTS) AND CERTAIN FIELD

### USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING

(1) Wear waterproof pants, coat, hat, rubber boots or rubber overshoes. (2) Wear safety goggles. (3) Wear mask or respirator approved by the U.S. Bureau of Mines for EPN protection. (4) Wear heavy duty, natural rubber gloves.

### WORK SAFETY RULES

Keep all unprotected persons and children away from treated area or where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way STOP work and get help right away. Call a doctor (physician, clinic or hospital) immediately. Explain that the victim has been exposed to EPN and describe his condition. After first aid is given (see First Aid Treatment Section) and if a doctor cannot come, take victim to clinic or hospital.

**IMPORTANT!** Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smoking, eating or drinking.

**AFTER WORK,** take off all work clothes and shoes. Shower with soap and water. Wash only clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirator should be cleaned and filter replaced according to instructions included with respirator.

### POISON SIGNS (Symptoms)

EPN is a very strong poison that rapidly enters the body on contact with an skin surface and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment if they may die.

Some of the signs and symptoms of poisoning are: headache, nausea, vomiting, cramps, weakness, blurred vision, pin-point pupils, tightness in chest, labored breathing, nervousness, sweating, watering of eyes, drooping or frothing of mouth and nose, muscle spasms and coma.

### FIRST AID TREATMENT

Call a doctor (physician, clinic or hospital) immediately. Explain that the victim has been exposed to EPN and describe his condition.

**IF BREATHING HAS STOPPED,** start artificial respiration immediately and maintain until doctor sees victim.

**IF SWALLOWED** and victim is awake (conscious) make him vomit quickly. Induce vomiting by sticking finger down throat or by giving soupy or strong salty water to drink. Repeat until vomit is clear. Never give anything by mouth to an unconscious person. Have victim lie down and keep quiet. See doctor immediately.

**IN CASE OF CONTACT,** immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. See doctor immediately.

### NOTE TO PHYSICIAN

Antidote — For patients believed to be suffering from EPN poisoning administer atropine sulfate in large doses. TWO TO FOUR mg. intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 10 minute intervals until signs of chloroform appear. 2-PAM chloride is also an antidote and may be administered in conjunction with atropine. **DO NOT GIVE MORPHINE OR TRANQUILIZERS.** EPN is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal reactions may occur after initial improvement. **VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.**

### POST TREATED AREA

Consult your State Agricultural Extension Service or Experiment Station regarding posting treated areas.

### ACTIVE INGREDIENTS:

O-ethyl O-p nitrophenyl phenylphosphonothioate ..... 40.0%

Aromatic Petroleum Solvent ..... 60.0%

**INERT INGREDIENTS:** ..... 100.0%

\*EPN Contains 4 pounds of EPN per gallon.


**STOP — READ LABEL BEFORE USING!**

**POISON**  
SEE SIDE  
PANELS FOR  
ANTIDOTE AND  
PRECAUTIONS




**DANGER**  
KEEP OUT OF  
THE REACH OF  
CHILDREN

**NOT FOR HOME USE**




### CAN KILL IF SWALLOWED

This product can kill you if swallowed even in small amounts; spray mist may be fatal if swallowed.



### CAN KILL BY SKIN CONTACT

This product can kill you if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes).



### CAN KILL IF BREATHED

This product can kill you if spray mist are breathed.

### POISONOUS TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur.

### HIGHLY TOXIC TO BEES

This product is highly toxic to bees exposed to direct treatment or residues on crops. For more information may be obtained from your Cooperative Agricultural Extension Service.

### GENERAL INFORMATION

Where a dosage range is given use the lower rate on small or young with light insect infestations; use the higher rate on large or mature with heavy insect infestations.

The hazards and precautions for handling RIVERSIDE EPN 4 in this are equally applicable to it after dilution with water for spray application.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR FLAME.**

**READ ENTIRE LABEL BEFORE USING THIS PRODUCT;** Use of this in a manner not in accordance with the information on this label may kill humans, domestic animals, wildlife, aquatic life, desirable plants, excessive and unlawful chemical residues, or produce other undesirable results.

### DIRECTIONS FOR USE — FRUITS AND NUTS

RIVERSIDE EPN 4 is useful for the control of certain insect pests of apples, apricots, cherries, citrus, grapes, nectarines, peaches, pears, plums, and walnuts.

RIVERSIDE EPN 4 will normally be applied to fruit and nut fields, orchards and groves with spray equipment. The use of conventional sprayers increases the hazard of use. If higher concentrations are used with equipment such as Spread Sprayers, the usual rigid precautions contact with the spray mixture must be followed. The use of RIVERSIDE EPN 4 in mist blower or thermal aerosol type (fog) equipment is not recommended. Treatment should begin with the first signs of infestation; for fruits the first spray should be made following bloom. Apply one spray and retreat as necessary. Do not apply during bloom of fruit or later than the following days prior to picking, or during citrus — 30 days after fruit and nut harvest.

For control of **Apple** (Red Spotted, European Red, Fall Webworm, Spotted, and Scab) on **Almonds**, **Apricots**, **Cherries**, **Grapes**, **Peaches**, **Pears**, **Plums**, **Prunes**, and **Walnuts** apply 1/2 pint to 1/4 pint of RIVERSIDE EPN 4.

For insect control on **apples**, **cherries**, **grapes**, **plums** and **prunes** apply 1/2 pint to 1/4 pint of RIVERSIDE EPN 4 for Plum Curculio and Oriental Fruit Moth; for Peach Tree Borer and Lesser Peach Tree Borer apply 1/2 to 1/4 pint; for Locust Borer (borers) and Pear Borer apply 1/2 pint (draw off) apply 1/2 pint for Fruit Tree Leafroller on cherries and borers on prunes apply 1/2 pint.

For insect control on **apples** and **pears** apply 1/2 pint to 1/4 pint of RIVERSIDE EPN 4 for Plum Curculio or Oriental Fruit Moth apply 1/2 pint, for Fruit Tree Leafroller apply 1/2 pint for Pear Psylla apply 1/2 pint. Do not use on Maine tested varieties of apples.

For insect control on **walnuts** apply 1/2 pint of RIVERSIDE EPN 4 for Moth and Walnut Aphid; for Fruit Tree Leafroller apply 1/2 pint.

For insect control of **grapes** and **peaches** apply 1/2 pint of RIVERSIDE EPN 4 for Grape Berry Moth; for Pecan Nut Casebearer apply 1/2 pint. Do not use on treated pecan groves.

For insect control on **citrus** fruits apply 1/2 pint to 1/4 pint of RIVERSIDE EPN 4 for Citrus Thrips. Apply 3 to 4 pints per acre — coverage for Fruit Tree Leafroller and Orange Tortix apply 1 1/2 to 4 1/2 pints.

The above figures for amounts of RIVERSIDE EPN 4 to apply are for use only if applied by high pressure hydraulic equipment.

Do not use RIVERSIDE EPN 4 in combination with Bordeaux or zinc lime spray.

Do not use RIVERSIDE EPN 4 in home plantings.

EPA Reg. No. 9779-32

NET CONTENTS \_\_\_\_\_



MANUFACTURED BY  
**RIVERSIDE CHEMICAL CO.**  
A SUBSIDIARY OF COOK INDUSTRIES, INC. MEMPHIS, TN 38116

Riverside serves agriculture. Agriculture serves everyone.

LABEL PRINTS IN THREE COLORS: GREEN (1PI INK #CF/3Y3/03) RED (1PI INK #RED) AND BLACK. (See figure overlay for color distribution.)

REPRODUCE

# RIVERSIDE EPN 4

AN EMULSIFIABLE CONCENTRATE CONTAINING 4 POUNDS OF EPN PER GALLON. USEFUL FOR THE CONTROL OF CERTAIN INSECT PESTS OF CERTAIN FRUITS AND NUTS (ALMONDS, APPLES, APRICOTS, CHERRIES, CITRUS, GRAPES, NECTARINES, PEACHES, PEARS, PECANS, PLUMS, AND WALNUTS) AND CERTAIN FIELD CROPS (BEANS, CORN, COTTON AND TOMATOES).

INGREDIENTS:	
Diethyl phosphorothioate	46.0%
Diethylamine	44.9%
Diethylamine	8.1%
<b>TOTAL</b>	<b>100.0%</b>

4 pounds of EPN per gallon.


**READ LABEL BEFORE USING!**

**KEEP OUT OF THE REACH OF CHILDREN**




**DANGER**

**NOT FOR HOME USE**



**DO NOT TOUCH**

**CAN KILL BY SKIN CONTACT**  
This product can kill you if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes).



**DO NOT BREATHE**

**CAN KILL IF BREATHED**  
This product can kill you if spray mist are breathed.

**HIGHLY TOXIC TO BEES**  
This product is highly toxic to bees exposed to direct treatment or residues on crops. For more information contact your local Cooperative Agricultural Extension Service.

**GENERAL INFORMATION**

Where a dosage range is given use the lower rate on small or young plants or with light insect infestations; use the higher rate on large or mature plants or with heavy insect infestations.

The hazards and precautions for handling RIVERSIDE EPN 4 in this container are equally applicable to it after dilution with water for spray application.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

**READ ENTIRE LABEL BEFORE USING THIS PRODUCT:** Use of this product in a manner not in accordance with the information on this label may injure or kill humans, domestic animals, wildlife, aquatic life, desirable plants, create excessive and unwanted chemical residues, or produce other undesirable results.

**DIRECTIONS FOR USE — FRUITS AND NUTS**

RIVERSIDE EPN 4 is useful for the control of certain insect pests of almonds, apples, apricots, cherries, citrus, grapes, nectarines, peaches, pears, pecans, plums, and walnuts.

RIVERSIDE EPN 4 will normally be applied to fruit and nut fields by high pressure hydraulic or hand type spray equipment. The use of concentrations of RIVERSIDE EPN 4 greater than recommended for conventional hydraulic sprayers increases the hazards of use. If higher concentrations are employed with equipment such as Ground Sprayers, the most rigid precautions to avoid contact with the spray mixture must be followed. The use of RIVERSIDE EPN 4 in mist blower or thermal aerosol type (fog) equipment is not recommended.

Treatment should begin with the first signs of infestation; for fruits begin with the first cover spray following bloom. Apply one spray, and retreat as necessary. Do not apply during bloom or within 14 days of later than the following number of days prior to picking, pruning, etc. — 30 days, other fruits and nuts — 21 days.

For control of European Red Spider, European Red Flycatcher, Two-spotted Willow-leaf, and European Pear on almonds, apples, cherries, grapes, peaches, pears, pecans, plums, and walnuts apply 1/4 pint to 1/2 pint of RIVERSIDE EPN 4.

For insect control on apples, cherries, plums and peaches apply 1/2 pint to 3/4 pint of RIVERSIDE EPN 4 for Plum Curculionid and Quince Fruit Moth; for Peach Tree Borer and Lesser Peach Tree Borer apply 1/2 to 1 1/2 pints; for Lecanium Scale (Crawley) and Pear Psyllid apply 1/2 pint; for Fruit Tree Leafroller on cherries and branches on plums apply 1/2 pint.

For insect control on apples and pears apply 1/2 pint to 3/4 pint of RIVERSIDE EPN 4 for Plum Curculionid for Codling Moth apply 1/2 pint; for Fruit Tree Leafroller apply 1/2 pint; for Pear Psyllid apply 1/2 pint. Do not use on McIntosh and related varieties of apples.

For insect control on walnuts apply 1/2 pint of RIVERSIDE EPN 4 for Codling Moth and Walnut Aphid; for Fruit Tree Leafroller apply 1/2 pint.

For insect control on grapes and pecans apply 1/2 pint of RIVERSIDE EPN 4 for Grape Berry Moth; for Pecan Nut Casebearer apply 1/2 pint. Do not graze livestock in treated pecan groves.

For insect control on citrus fruits apply 2 1/2 pints to 3 pints of RIVERSIDE EPN 4 for Citrus Thrips; for Citrus Thrips apply 3 to 4 pints per acre — outside coverage; for Fruit Tree Leafroller and Orange Tortix apply 1 1/2 to 4 1/2 pints per acre.

The above figures for amounts of RIVERSIDE EPN 4 to apply are per 100 gallons of spray as applied by high pressure hydraulic equipment.

Do not use RIVERSIDE EPN 4 in combination with Bordeaux or zinc sulfide — lime spray.

Do not use RIVERSIDE EPN 4 in home plantings.

**DIRECTIONS FOR USE — FIELD CROPS**

RIVERSIDE EPN 4 is useful for the control of certain insect pests of beans, corn, cotton, and tomatoes.

RIVERSIDE EPN 4 may be applied by ground sprayers or by airplanes, mixing the RIVERSIDE EPN 4 in water to form an emulsion spray solution. To make the spray solution, half fill the spray tank with water, agitate, add the proper amount of RIVERSIDE EPN 4, and then fill the spray tank with water while still agitating. Continue to agitate during use.

When applying to field crops by ground sprayers calibrate so as to apply the recommended amount of RIVERSIDE EPN 4 in not less than 3 gallons of water per acre. Airplanes should apply the recommended amount of RIVERSIDE EPN 4 in not less than 1 gallon of water per acre. Always use sufficient water for adequate coverage of the crop foliage.

**BEANS** — For control of Mexican Bean Beetle make two to four applications of 1/2 to 1 pint of RIVERSIDE EPN 4 per acre per application after eggs are laid. Do not apply RIVERSIDE EPN 4 to beans within 21 days of harvest.

**CORN** — For control of European Corn Borer on sweet corn apply 1/2 pint of RIVERSIDE EPN 4 per acre 5 to 7 days after eggs are laid; repeat 7 to 10 days later at intervals of 5 to 7 days during periods when moths are actively depositing eggs.

For control of European Corn Borer in field corn apply 1/2 pint of RIVERSIDE EPN 4 per acre 10 days after eggs first hatch or when 75% of plants show evidence of feeding in leaf whorls; repeat 7 to 10 days later if new evidence of leaf feeding is present. For second brood borers, apply when eggs begin to hatch.

Do not apply RIVERSIDE EPN 4 to corn within 14 days of harvest.

**COTTON** — For control of Cotton Fleahopper, Cotton Leafworm, Thrips, Yellow-striped Armyworm and Red Spider apply 1/2 to 1 pint of RIVERSIDE EPN 4 per acre per application; for control of Bollworm apply 1 to 1 1/2 pints per acre; for control of Bollworm or Tobacco Budworm apply 2 pints per acre.

Do not apply RIVERSIDE EPN 4 within 3 days of handpicking or harvest.

**TOMATOES** — For control of Potato Aphid, Serpentine Leaf Miner, Southern Armyworm, and Tomato Fruitworm apply 1/2 to 1 pint of RIVERSIDE EPN 4 per acre per application at 7 to 10 day intervals as needed.

Do not apply RIVERSIDE EPN 4 later than 3 days before harvest.

**DISPOSAL OF CONTAINER AND WASTES**

This container should not be reused for ANY purpose. When empty, it should be rendered useless by punching large holes in the top, bottom, and sides with an instrument such as an axe. It should then be mashed as flat as possible and buried underground 2 feet deep as practical, with at least 18 inches of soil covering. This should be done in an isolated area, away from habitation, and in a location which would not drain into, or cause pollution of lakes, streams, or ponds.

Any spray wastes or spillage resulting from the application of this material or the destruction of the container should be neutralized with a caustic solution and buried in a pit with the disposed container.

**NOTICE**

Riverside Chemical Company warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label, when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Riverside Chemical Company, and buyer assumes the risk of any such use.

**AN EMULSIFIABLE CONCENTRATE CONTAINING 4 POUNDS OF EPN PER GALLON. USEFUL FOR THE CONTROL OF CERTAIN INSECT PESTS OF CERTAIN FRUITS AND NUTS (ALMONDS, APPLES, APRICOTS, CHERRIES, CITRUS, GRAPES, NECTARINES, PEACHES, PEARS, PECANS, PLUMS, AND WALNUTS) AND CERTAIN FIELD CROPS (BEANS, CORN, COTTON AND TOMATOES).**

**GENERAL INFORMATION**

Where a dosage range is given use the lower rate on small or young plants or with light insect infestations. Use the higher rate on large or mature plants or with heavy insect infestation.

The hazards and precautions for handling RIVERSIDE EPN 4 in this container are equally applicable to it after dilution with water for spray application.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

**READ ENTIRE LABEL BEFORE USING THIS PRODUCT:** Use of this product in a manner not in accordance with the information on this label may injure or kill humans, domestic animals, wildlife, aquatic life, desirable plants, create excessive and unlawful chemical residues, or produce other undesirable results.

**DIRECTIONS FOR USE—FRUITS AND NUTS**

RIVERSIDE EPN 4 is useful for the control of certain insect pests of almonds, apples, apricots, cherries, citrus, grapes, nectarines, peaches, pears, pecans, plums, and walnuts.

RIVERSIDE EPN 4 will normally be applied to fruit and nut trees by high pressure hydraulic or other type spray equipment. The use of low concentrations of RIVERSIDE EPN 4 greater than recommended for conventional hydraulic sprayers increases the hazards of use. If higher concentrations are employed with equipment such as Ground Sprayers, the most rigid precautions to avoid contact with the spray mixture must be followed. The use of RIVERSIDE EPN 4 in air blower or thermal aerosol type fog equipment is not recommended.

Treatment should begin with the first signs of infestation. For fruits begin with the first opportunity for a good spray. Apply one spray and retreat as necessary. Do not apply during bloom of fruit or later than the following number of days prior to picking, or during citrus — 30 days, other fruits and nuts — 21 days.

For control of ~~Apple Red Spider, European Red, Pear, and Two spotted~~ ~~Worm~~ ~~and~~ ~~Apple~~ ~~on~~ ~~almonds,~~ ~~apples,~~ ~~cherries,~~ ~~citrus,~~ ~~grapes,~~ ~~nectarines,~~ ~~peaches,~~ ~~pears,~~ ~~pecans,~~ ~~plums,~~ ~~and~~ ~~walnuts~~ apply 1/2 pint to 1/4 pint of RIVERSIDE EPN 4.

For insect control on apricots, cherries, grapes, plums and pears apply 1/2 pint to 1/4 pint of RIVERSIDE EPN 4 for Plum Curculio; for Codling Moth apply 1/2 pint; for Fruit Tree Leafhopper apply 1/2 pint; for Pear Psylla apply 1/2 pint. Do not use on Mountain and related varieties of apples.

For insect control on apples and pears apply 1/2 pint to 1/4 pint of RIVERSIDE EPN 4 for Fruit Curculio; for Codling Moth apply 1/2 pint; for Fruit Tree Leafhopper apply 1/2 pint; for Pear Psylla apply 1/2 pint. Do not use on Mountain and related varieties of apples.

For insect control on walnuts apply 1/2 pint of RIVERSIDE EPN 4 for Codling Moth and Walnut Aphid; for Fruit Tree Leafhopper apply 1/2 pint.

For insect control on grapes and pecans apply 1/2 pint of RIVERSIDE EPN 4 for Grape Berry Moth; for Pecan Nut Casebearer apply 1/2 pint. Do not graze livestock in treated pecan groves.

For insect control on citrus fruits apply 2 1/2 pints to 3 pints of RIVERSIDE EPN 4 for Citrus Thrips; for Citrus Thrips apply 3 to 4 pints per acre — outdoor coverage; for Fruit Tree Leafhopper and Orange Tortix apply 1 1/2 to 4 1/2 pints per acre.

The above figures for amounts of RIVERSIDE EPN 4 to apply are per 100 gallons of spray as applied by high pressure hydraulic equipment.

Do not use RIVERSIDE EPN 4 in combination with Bordeaux or zinc sulfate — 1000 ppm.

Do not use RIVERSIDE EPN 4 in home plantings.

**DIRECTIONS FOR USE—FIELD CROPS**

RIVERSIDE EPN 4 is useful for the control of certain insect pests of beans, corn, cotton, and tomatoes.

RIVERSIDE EPN 4 may be applied by ground sprayers or by airplanes, mixing the RIVERSIDE EPN 4 in water to form an emulsion spray solution. To make the spray solution, half fill the spray tank with water, agitate, add the proper amount of RIVERSIDE EPN 4, and then fill the spray tank with water while still agitating. Continue to agitate during use.

When applying to field crops by ground sprayers calibrate so as to apply the recommended amount of RIVERSIDE EPN 4 in not less than 3 gallons of water per acre. Airplanes should apply the recommended amount of RIVERSIDE EPN 4 in not less than 1 gallon of water per acre. Always use sufficient water for adequate coverage of the crop foliage.

**BEANS** — For control of Mexican Bean Beetle make two to four applications of 1 to 1 1/2 pint of RIVERSIDE EPN 4 per acre per application after eggs are laid. Do not apply RIVERSIDE EPN 4 to beans within 21 days of harvest.

**CORN** — For control of European Corn Borer on sweet corn apply 1/2 pint of RIVERSIDE EPN 4 per acre 5 to 7 days after eggs are laid; repeat 7 to 10 days later and at intervals of 5 to 7 days during periods when moths are actively depositing eggs.

For control of European Corn Borer in field corn apply 1/2 pint of RIVERSIDE EPN 4 per acre 10 days after eggs first hatch or when 75% of plants show evidence of feeding in leaf whorls; repeat 7 to 10 days later if new evidence of leaf feeding is present. For second brood borers, apply when eggs begin to hatch.

Do not apply RIVERSIDE EPN 4 to corn within 14 days of harvest.

**COTTON** — For control of Cotton Fleahopper, Cotton Leafworm, Thrips, Yellow-shouldered Weevil and Red Spinger apply 1/2 to 1 pint of RIVERSIDE EPN 4 per acre per application for control of Boll Weevil apply 1 to 1 1/2 pints per acre; for control of Bollworm or Tobacco Budworm apply 2 pints per acre.

Do not apply RIVERSIDE EPN 4 within 3 days of handpicking or harvest.

**TOMATOES** — For control of Potato Aphid, Serpentine Leaf Miner, Southern Armyworm, and Tomato Fruitworm apply 1/4 to 1/2 pint of RIVERSIDE EPN 4 per acre per application at 7 to 10 day intervals as needed.

Do not apply RIVERSIDE EPN 4 later than 3 days before harvest.

**DISPOSAL OF CONTAINER AND WASTES**

This container should not be reused for ANY purpose. When empty, it should be rendered useless by punching large holes in the top, bottom, and sides with a instrument such as an axe; it should then be mashed as flat as possible and buried underground as deep as practical, with at least 18 inches of soil covering. This should be done in an isolated area, away from habitation, and in a location which would not drain into, or cause pollution of lakes, streams, or ponds.

Any spray wastes or spillage resulting from the application of this material or the destruction of the container should be neutralized with a caustic solution and buried in a pit with the disposed container.

**NOTICE**

Riverside Chemical Company warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Riverside Chemical Company, and buyer assumes the risk of any such use.

