

RIVERSIDE CHLORDANE 4

ACCEPTED

APR 6 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 9779-33

CONTAINING 4 POUNDS CHLORDANE PER GALLON.
CONTAIN INSECT PESTS OF LAWNS, HOMES, GARDENS, GOLF GREENS, AND CERTAIN CROPLANDS.

44.0%
53.0%
3.0%
100.0%
done and

When
the skin, can
see of spillage
minimization of
apply in living

ing on treated
plication rates
stant resources.
off is likely to
or disposal of
areas treated.

is or to animals
EAT

of this product
label may injure
sirable plants,
other undesir-

mole crickets,
the rate of 3
r to cover 1,000
at as necessary.

silverfish, and
(3) tablespoons
such as cracks,
s hiding places.
DE 4, thoroughly
es around door
Care should be
food and cook-
ts are for spot

SUBTERRANEAN TERMITES (Directions for Non-Professional Use): Use RIVERSIDE CHLORDANE 4 in a 2% water emulsion prepared by mixing 1 quart RIVERSIDE CHLORDANE 4 in 25 gallons of water

BUILDINGS HAVING BASEMENTS OR CRAWL SPACES: Dig a trench along the outside of the foundation walls. In brick or hollow block or concrete foundations, dig a trench to, but not below, the footing. Then as the trench is refilled, treat the soil at the rate of 4 gallons per 10 linear feet for each foot of depth. A trench 3 feet deep would require 12 gallons per 10 linear feet.

The objective is to establish a treated soil barrier which will prevent termite entry. Other critical areas may also require treatment. It is suggested that you consult the State Extension Service for advice.

GRASSHOPPERS: In early season when grasshoppers are small and concentrated in hatching areas, in field margins and along fence rows apply 1 pint of RIVERSIDE CHLORDANE 4 per acre in sufficient water for coverage. Later in the season when grasshoppers are in the late stages of growth and vegetation is tall and dense, increase the rate to 1 quart of RIVERSIDE CHLORDANE 4 in sufficient water for coverage.

STRAWBERRY CROWN BOBBER: Apply to foliage at the rate of 1 to 2 quarts of RIVERSIDE CHLORDANE 4 per acre in sufficient water for coverage. Do not apply after fruit begins to form.

CUTWORMS, MOLE CRICKETS: On beets, celery, collards, kale, lettuce, okra, peppers and potatoes apply 1 to 2 quarts of RIVERSIDE CHLORDANE 4 per acre in sufficient water for coverage. Apply to soil around the bases of plants. Do not apply after edible parts of plants begin to form. Do not apply to beets after seedling stage if tops are to be used for food.

WIREWORMS: BROADCAST TREATMENT - Crop land to be planted to beans, beets, blackberries, broccoli, Brussels sprouts, cabbage, cauliflower, celery, corn, grapes, kohlrabi, loganberries, peanuts, peppers, potatoes, raspberries, rutabaga, strawberries, sweet potatoes, and turnips can be treated prior to planting time. RIVERSIDE CHLORDANE 4 should be applied to the surface of the soil and disced in to a depth of 4 to 9 inches. Application rates vary from 2 1/2 to 5 quarts of RIVERSIDE CHLORDANE 4 per acre depending upon the species of wireworms present and type of soil. Consult Cooperative Agricultural Extension Service specialists for proper amount of RIVERSIDE CHLORDANE 4 to use. The average clay or loam soils are usually treated at the rate of 2 1/2 to 3 quarts of RIVERSIDE CHLORDANE 4 per acre in sufficient water for coverage.

SOUTHERN CORN ROOTWORM: On corn and peanuts apply 4 quarts of RIVERSIDE CHLORDANE 4 in sufficient water for coverage. Apply broadcast treatments as directed under WIREWORMS.

SEED CORN MAGGOT: On corn, beans and cotton apply 2 1/2 quarts of RIVERSIDE CHLORDANE 4 per acre in sufficient water for coverage. Apply uniformly to the soil surface before planting.

WHITE GRUBS, JAPANESE BEETLE LARVAE, WHITE-FRINGED BEETLE LARVAE: On golf greens, lawns, and infested cropland listed under WIREWORMS apply at the rate of 5 to 10 quarts of RIVERSIDE CHLORDANE 4 per acre (1/2 to 1 cup in 5 gallons of water applied to 1,000 square feet) in sufficient water for coverage. On crop land disc into the soil before planting the crop. On lawns and golf greens water thoroughly after application. Do not apply to edible parts of food crops. Do not allow dairy animals or animals being finished for slaughter to graze treated feed or forage crops.

DISPOSAL OF CONTAINER AND WASTES

This container should not be reused for ANY purpose. When empty, it should be rendered useless by punching large holes in the top, bottom, and sides with an instrument such as an axe. It should then be mashed as flat as possible and buried underground as deep as practical, with at least 18 inches of soil covering. This should be done in an isolated area, away from habitation, and in a location which would not drain into, or cause pollution of lakes, streams, or ponds. Any spray wastes or spillage resulting from the application of this material or the destruction of the container should be neutralized with a caustic solution and buried in a pit with the disposed container.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instruction, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA REG. NO. 9779-33

NET CONTENTS _____



MANUFACTURED BY
RIVERSIDE CHEMICAL CO.
DIVISION OF COOK INDUSTRIES, INC., MEMPHIS, TENN. 38116
Riverside serves agriculture. Agriculture serves everyone.

COLORS: GREEN, WHITE-PI. INK No. 4444 CF/3y9/03
BLACK.