

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING

FIRST AID TREATMENT

WORK SAFETY RULES

IF BREATHING HAS STOPPED


IF SWALLOWED

IN CASE OF CONTACT

NOTE TO PHYSICIAN

POISON SIGNS (Symptoms)

POST TREATED AREA



ACTIVE INGREDIENTS	
1. 45.4%	2. 50.6%
INERT INGREDIENTS 4.0%	
TOTAL 100.0%	


STOP - READ LABEL BEFORE USING!

POISON
SEE SIDE
PANELS FOR
ANTIDOTE AND
PRECAUTIONS



DANGER
KEEP OUT OF
THE REACH OF
CHILDREN

NOT FOR HOME USE



DO NOT SWALLOW



DO NOT TOUCH



DO NOT BREATHE

POISONOUS TO FISH AND WILDLIFE

HIGHLY TOXIC TO BEES



BEST AVAILABLE COPY

GENERAL INFORMATION

RIVERSIDE METHYL PARATHION 4 is useful for the control of certain insect pests of alfalfa, beans, corn, grapes, and cole crops (broccoli, cauliflower, cabbage, etc.). It is also useful for the control of grasshoppers, beet beetles, and other insects on alfalfa, beans, peppers, potatoes, and other small grains (barley, oats, etc.). It is also useful for the control of certain insects on sweet potatoes, tobacco, and tomatoes.

RIVERSIDE METHYL PARATHION 4 may be applied by ground sprayers or by airplanes using the RIVERSIDE METHYL PARATHION 4 in water to form a emulsion spray solution. To make the spray solution, add the spray rate with water, agitate, add the proper amount of RIVERSIDE METHYL PARATHION 4, and then fill the spray tank or the sprayer while still agitating. Continue to agitate during use.

When applying with ground sprayers, calculate so as to apply the recommended amount of RIVERSIDE METHYL PARATHION 4 in not less than 3 gallons of water per acre. Airplanes should apply the recommended amount of RIVERSIDE METHYL PARATHION 4 in not less than 1 gallon of water per acre. Where a dosage range is given use the lower rate on small or young plants or with light insect infestations; use the higher rate on large or mature plants or with heavy insect infestations.

The amounts to use given in DIRECTIONS FOR USE are the amounts of RIVERSIDE METHYL PARATHION 4 to apply per acre per application. Repeat applications as necessary unless stated otherwise in DIRECTIONS FOR USE. The hazards and precautions for RIVERSIDE METHYL PARATHION 4 in this container are equally applicable to other solutions with water to which application.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT. Use of this product in a manner not in accordance with the information on this label may cause or contribute to the death of humans, domestic animals, aquatic life, desirable plants, crops, excessive and unwanted chemical residues, or produce other undesirable results.

DIRECTIONS FOR USE

ALFALFA For control of Aphids including Pea Aphid and Spined Alfalfa Aphid, Armyworms, Cabbage Cutworms, Flea Beetles and Leafhoppers, apply 1 to 1 1/2 pints for Legume Bugs, Mites, and Stem-bugs; apply 1 to 1 1/2 pints for Clover Leaf Beetles and Weevils; apply 1 pint for Grasshoppers and Thrips; apply 1 1/2 pints for Egyptian Alfalfa Weevil; apply 1 to 2 pints for Bean Aweil, Alfalfa Weevil, and Yellow-striped Armyworm; apply 1 1/2 pints.

Do not apply later than 20 days prior to harvest, cutting, or forage use. Do not apply during bloom period to avoid injury to honeybees.

CLOVER For control of Alfalfa Weevil, apply 1 to 1 1/2 pints when 75% of the clover plants are in bud and shoots appear ragged or when the green leaf reaches 25 per cent. For Aphids, Armyworms, and Leafhoppers, apply 1 to 1 1/2 pints for Alfalfa Weevil, adults; Clover Leaf Weevil, and Weevils; apply 1 pint for Legume Bugs and Mites; apply 1 to 1 1/2 pints for Stem-bugs; apply 1 to 1 1/2 pints for Egyptian Alfalfa Weevil; apply 1 to 1 1/2 pints for Cabbage Cutworm and Cabbage Loopworm; apply 1 to 1 1/2 pints.

Do not apply later than 20 days prior to harvest, cutting, or forage use. Do not apply during bloom period to avoid injury to honeybees.

BEANS (Dry Green Lima) For control of Aphids, Armyworm, Flea Beetles, Leafhoppers, Mites, and Plant Bugs, apply 1 to 1 1/2 pints for Legume Bugs, apply 1 to 1 1/2 pints for Egyptian Alfalfa Weevil; apply 1 to 1 1/2 pints for Cabbage Cutworm and Cabbage Loopworm; apply 1 to 1 1/2 pints for Stem-bugs; 2 to 3 pints.

Do not apply later than 20 days prior to harvest, cutting, or forage use. Do not apply during bloom period to avoid injury to honeybees.

CABBAGE AND COLE CROPS (Broccoli, Brussels Sprouts, Cauliflower) For control of Aphids, Armyworms, and Flea Beetles, apply 1 to 1 1/2 pints for Cabbage Cutworm, Flea Beetles, Grasshoppers, and Mites; apply 1 to 1 1/2 pints for Imported Cabbageworm, Cabbage Loopworm, and Stem-bugs; 2 to 3 pints for Plant Bugs; 3 pints.

Do not apply later than 20 days prior to harvest, cutting, or forage use. Do not apply during bloom period to avoid injury to honeybees.

COLLARD RALE AND MUSTARD GREENS For control of Aphids, Armyworms, Flea Beetles, Leafhoppers and Mites apply 1 to 2 1/2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Bugs apply 3 pints.

Do not apply more than 10 days before harvest.

CORN For control of Aphids including Corn Leaf Aphid, Flea Beetles, and Mites, including Western Corn Rootworm, Armyworm, Flea Beetles, and Strawbugs apply 1 pint.

Do not apply more than 10 days before harvest.

Workers entering treated fields within 24 hours after application should wear protective clothing.

COTTON For control of Fleas, apply 1 to 1 1/2 pints per acre during the 4 leaf stage and repeat 1 to 3 times at 7 to 10 day intervals.

For early season control of Fleas, including Flea Beetles, apply 1 to 2 pints as early as the 2 to 4 leaf stage and repeat 1 to 3 times at 7 to 10 day intervals. For mid season control apply 1 to 2 pints at the start of square formation to 25% of the first squares the plants and repeat at 7 to 10 day intervals as needed to control the Fleas in the full green. Repeat at 7 to 10 days during the major flowering period and periods of heavy insect pressure. Up to 6 pints may be applied as needed.

For control of the leafminer, Garden Webworm, Flea Beetles, and other pests, including Corn Rootworm, Flea Beetles, including Flea Beetles, and Mites, apply 1 to 2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest. Workers entering treated fields within 24 hours after application should wear protective clothing.

GRASS Hay Pasture For control of Fleas, including Flea Beetles, apply 1 to 2 pints.

Do not apply more than 10 days before harvest.

PEAS For control of Fleas, including Flea Beetles, Leafhoppers, and Mites, apply 1 to 2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest.

PUMPKINS For control of Fleas, including Flea Beetles, Leafhoppers, and Mites, apply 1 to 2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest.

POTATOES For control of Fleas, including Flea Beetles, Leafhoppers, and Mites, apply 1 to 2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest.

RAPE For control of Fleas, including Flea Beetles, Leafhoppers, and Mites, apply 1 to 2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest.

SMALL GRAINS (Barley, Oats, Rye, Wheat) For control of Fleas, including Flea Beetles, Leafhoppers, and Mites, apply 1 to 2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest.

SORGHUM For control of Fleas, including Flea Beetles, Leafhoppers, and Mites, apply 1 to 2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest.

Leafhoppers, Flea Beetles, and Mites, apply 1 to 2 1/2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest.

SOYBEANS For control of Fleas, including Flea Beetles, Leafhoppers, and Mites, apply 1 to 2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest.

Workers entering treated fields within 24 hours after application should wear protective clothing.

SPINACH For control of Fleas, including Flea Beetles, Leafhoppers, and Mites, apply 1 to 2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest.

SUGAR BEETS For control of Fleas, including Flea Beetles, Leafhoppers, and Mites, apply 1 to 2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest.

Workers entering treated fields within 24 hours after application should wear protective clothing.

SWEET POTATOES For control of Fleas, including Flea Beetles, Leafhoppers, and Mites, apply 1 to 2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest.

TOBACCO For control of Fleas, including Flea Beetles, Leafhoppers, and Mites, apply 1 to 2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest.

Workers entering treated fields within 24 hours after application should wear protective clothing.

TOMATOES For control of Fleas, including Flea Beetles, Leafhoppers, and Mites, apply 1 to 2 pints of Methyl Parathion (Cabbage, Turnip, Strawbug, and Flea Beetles) 2 to 3 pints for Flea Beetles, and Legume Bugs 2 to 3 pints for Leafhoppers, Flea Beetles, and Flea Beetles 1 pint for Armyworms 2 pints for Bombyx 2 to 3 pints.

Do not apply more than 10 days before harvest.

DISPOSAL OF CONTAINER AND WASTES This material is extremely toxic and should be disposed of in accordance with the instructions on the label. Do not use this material for any purpose other than that for which it is intended. Do not use this material for any purpose other than that for which it is intended. Do not use this material for any purpose other than that for which it is intended.

NOTICE This material is extremely toxic and should be disposed of in accordance with the instructions on the label. Do not use this material for any purpose other than that for which it is intended. Do not use this material for any purpose other than that for which it is intended. Do not use this material for any purpose other than that for which it is intended.



METHYL PARATHION MIXTURE, LIQUID

BEST AVAILABLE COPY