

Right label

RIVERSIDE BRAND 50% CONCENTRATE 100%  
INSECTICIDE

Contains DDT as active ingredient  
FOR THE CONTROL OF INSECTS ON COTTON

Active Ingredient:

Dibutyl (1-naphthyl N-methylcarbamate) . . . . . 50% w/w  
Inert ingredients . . . . . 50% w/w

U. S. Pat. Nos. 2,903,478; 3,009,055

CAUTION: KEEP OUT OF REACH OF CHILDREN.

~~XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX~~

Manufactured by

RIVERSIDE CHEMICAL COMPANY  
Macon, Miss.

AC  
1277-01

Left label

Directions for Use

For early-season control, apply 1 to 2 pounds of Riverside Brand 50% Concentrate per acre.

For cotton leafworm and striped blister beetle:

Apply 1 to 2 pounds of Riverside Brand 50% Concentrate per acre at the first appearance of insects.

For fly- and web-worm control on all varieties of cotton:

Cotton leaf miner, leaf rollers, webworms, and bollworms:

plant bug. Also for fly- and web-worm control on all varieties of cotton

in western irrigated cotton.

Apply 1 to 2 pounds of Riverside Brand 50% Concentrate per acre, depending on the size of the cotton and insect population and the level of infestation. Applications should be made every 7 to 10 days until control is achieved.

For pink bollworm:

Apply 3 to 5 pounds of Riverside Brand 50% Concentrate per acre, depending on the degree of infestation.

For stink bug and cotton-nose weevil control:

Apply 1 pound of Riverside Brand 50% Concentrate per acre.

For grasshopper:

Apply 2 to 3 pounds of Riverside Brand 50% Concentrate per acre at the first appearance of insects and their feeding damage. A higher dosage should be used when grasshoppers are present in large numbers. This insecticide is applied on partly defoliated plants.

Aphid populations will be controlled by application of this insecticide.

~~Left Panel~~ CAUTION: HARMFUL IF INHALED OR SWALLOWED.

Avoid prolonged or repeated breathing of dust.

Do not take internally.

Skin contact with any insecticide may be harmful.

Avoid unnecessary contact. Wash hands and face before eating.

Take shower or bath after work.

Wear regular long sleeved work clothing. Change to clean clothing daily.

Note for physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of food, feed, feeding troughs and watering receptacles.

Keep out of reach of children and animals.

Right Panel

1. Calibrate ground equipment or aerial dust apparatus accurately. After properly adjusting equipment to deliver the required low volume, maintain constant speed throughout application of 50% SEVIN Dust to insure even coverage. Strictly avoid excessive application.
2. To effectively apply 50% SEVIN Dust by aircraft use standard venturi type equipment with spreader vanes or Swathmaster type spreaders. Fly at normal height (4 to 6 feet above the top of the plants is suggested) and maintain regular swath width (usually 30 to 35 feet is indicated). For application with ground equipment use tractor mounted power dusters with a metering device that feeds the 50% SEVIN Dust through one nozzle per row (with deflector) or similarly designed electronic dusters.
3. For optimum results apply 50% SEVIN Dust only when weather conditions are favorable. Wind and rising air currents may cause undesirable drift. Strictly avoid excessive dust drift away from crop being treated.
4. SEVIN does not control spider mites. Where infestations are encountered, use a recommended miticide.
5. SEVIN may be applied after bolls open. Allow 7 days between last application and grazing treated fields or harvesting crop residue for use as feed or bedding for dairy or meat animals.
6. For protection of honeybees avoid use, if possible, during periods when honeybees are visiting the crop or neighboring crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance (beyond bee flight range) until one week after application.

ACCEPTED

WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Net Contents: 50 Pounds

USDA Reg. No. 9779-

