

# RIVERSIDE 20% TOXAPHENE

USEFUL FOR THE CONTROL OF CERTAIN INSECT PESTS OF BEANS (Dry, Green, Lima), CABBAGE AND OTHER COLE CROPS, COWPEAS, CERTAIN SMALL GRAINS (Barley, Oats, Rice, Rye, Wheat), OKRA, PEAS, PEANUTS, PIMENTOS, SORGHUM, SOYBEANS, STRAWBERRIES, AND TOMATOES.

<b>ACTIVE INGREDIENT:</b>	
Toxaphene (Technical Chlorinated Camphene-chlorine content: 67% to 69%)	20.0%
<b>INERT INGREDIENTS</b>	80.0%
TOTAL	100.0%

## WARNING: KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed. Do not get on skin, in eyes, or on clothing. Wash thoroughly after using. Do not breathe dust.

Avoid contamination of feed or foodstuffs. Do not use on dairy animals or in dairy barns. Do not use on cucurbits.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the areas.

### ANTIDOTE

**INTERNAL:** Give sodium chloride 15 g (one tablespoonful of table salt) in glass of warm water and repeat until vomit is clear. Call a doctor immediately.

**EXTERNAL:** Wash with soap and water.

**READ ENTIRE LABEL BEFORE USING THIS PRODUCT:** Use of this product in a manner not in accordance with the information on this label may injure or kill humans, domestic animals, wildlife, aquatic life, desirable plants, create excessive and unlawful chemical residues, or produce other undesirable results.

### GENERAL INFORMATION

RIVERSIDE 20% TOXAPHENE DUST is useful for the control of certain insect pests of beans (dry, green, lima), cabbage and other cole crops, corn, cotton, cowpeas, okra, peas, peanuts, pimientos, certain small grains (barley, oats, rice, rye, wheat), sorghum, soybeans, spinach, strawberries and tomatoes.

The dosages given under DIRECTIONS FOR USE are the amounts of RIVERSIDE 20% TOXAPHENE DUST to be applied per acre. Treat per application. Repeat applications as necessary, unless otherwise stated in DIRECTIONS FOR USE.

Where a dosage range is given use the lower rate on small or young plants or with light insect infestations; use the higher rate on large or mature plants or with heavy insect infestations.

### DIRECTIONS FOR USE

**BEANS (Dry, Green, Lima):** For control of Mexican Bean Beetle and Thrips apply 6 to 10 pounds; for control of Corn Earworms, Grasshoppers, and Serpentine Leaf Miner apply 10 to 15 pounds; for control of Bean Lycaenid apply 12 to 18 pounds; for control of Waxbean Beetle apply 12 to 25 pounds.

Do not feed treated vines to dairy animals or animals being finished for slaughter.

**CABBAGE AND OTHER COLE CROPS (Also including Broccoli, Brussels Sprouts, Cauliflower, Collards and Kale):** For control of Armyworms, Fall Armyworms, Cabbage Aphid, Climbing and Surface Feeding Cutworms apply 8 to 10 pounds; for control of Imported Cabbageworms, Green Peach Aphid, Serpentine Leaf Miner, and Thrips apply 10 to 12 pounds; for control of Corn Earworm and Diamondback Moth apply 8 to 20 pounds; for control of Cabbage Looper apply 15 to 20 pounds.

Do not apply after edible parts begin to form or within 35 days of harvest. Treated plants must not be fed to dairy animals or animals being finished for slaughter.

**CORN:** For control of Chinch Bugs and Corn Billbug apply 10 pounds; for control of Armyworm and Budworm apply 15 pounds; for control of Corn Leafminer and Lesser Cornstalk Borer apply 20 to 23 pounds.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

**COTTON:** For control of Thrips apply 5 to 8 pounds; for control of Fleahoppers apply 5 to 15 pounds; for control of Armyworm, Beet Armyworm, Fall Armyworm, Boll Weevil, Cotton Leafworm, Cutworms, Flea Beetle, Grasshoppers, Lygus Bugs and other Mites, Plant Bugs, Yellow Striped Armyworm, and Western Yellow Striped Armyworm apply 10 to 20 pounds; for control of Cotton Leaf Perforator and Garden Webworm apply 20 to 23 pounds.

Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

**COWPEAS:** For control of Cowpea Curculio apply 8 to 15 pounds. Do not apply to cowpeas to be used as green snaps after pods begin to form.

Do not feed treated vines to dairy animals or animals being finished for slaughter.

**OKRA:** For control of Serpentine Leaf Miner apply 12 to 15 pounds. Do not apply after pods begin to form.

**PEAS:** For control of Leafminer and Lygus Bugs apply 12 to 15 pounds. Do not apply after pods begin to form if pods are to be used as food or within 7 days if pods are not to be used as food.

Do not feed vines to dairy animals or animals being finished for slaughter.

**PEANUTS:** For control of Thrips apply 10 to 12 pounds; for control of Fleahoppers and Leafworm apply 20 to 25 pounds; for control of Velvet Bean Caterpillar apply 8 to 15 pounds; for control of Fall Armyworm apply 10 to 20 pounds; for control of Southern Stink Bug apply 20 to 25 pounds.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

**PIMENTOS:** For control of Flea Beetles, Hornworm, Leafminers, and Pepper Weevil apply 7 to 10 pounds.

Do not apply later than 2 days prior to harvest.

**SMALL GRAINS (Barley, Oats, Rice, Rye, Wheat):** For control of Armyworms, Cutworms (climbing and surface), and Grasshoppers apply 10 to 12 pounds; for control of Chinch Bugs and Rice Stink Bug apply 10 pounds.

Do not feed treated forage to dairy animals or animals being finished for

**SORGHUM:** Apply the same as for CORN.

Do not apply later than 28 days before harvest to begin to term. Do not graze dairy animals on treated fields. Do not ensilage.

**SOYBEANS:** For control of Armyworms, Bollworm, Corn Earworm, Grasshoppers, Green Caterpillar apply 12 to 20 pounds.

Do not apply later than 21 days before harvest to ensilage made from treated plants to poultry finished for slaughter. Do not feed soybean to

**SPINACH:** Apply the same as for COLE CROPS.

Do not apply later than 21 days before harvest per season.

**STRAWBERRIES:** For control of Crown Borer apply 10 to 15 pounds.

Do not apply after fruit starts to form.

**TOMATOES:** For control of Armyworm, Blister Beetle, Potato Beetle, Cutworms, Fleabeetle, Hornworm, Pinworm, Russet Mites, Serpentine worm apply 10 to 25 pounds.

Do not apply later than 3 days before harvest.

### DISPOSAL OF CONTAINER AND WASTE

Destroy all containers when empty by puncturing, crushing, or by burning. Stay out of smoke and fumes for other purposes. Dispose of wastes by burying or by water supplies.

### NOTICE

Seller warrants that the product conforms substantially to the purposes stated on the label with directions under normal conditions of use. Any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product under abnormal conditions, or for a purpose not intended, or for a use not foreseeable to seller, and buyer assumes the risk.

EPA REG. NO. 9779-70



MANUFACTURED BY  
**RIVERSIDE**  
DIVISION OF COOK INDUSTRIES  
Riverside serves agriculture

# 20% TOXAPHENE DUST

CERTAIN INSECT PESTS OF BEANS (Dry, Green, Lima), CABBAGE AND OTHER COLE CROPS, CORN, COTTON, PEAS (Barley, Oats, Rice, Rye, Wheat), OKRA, PEAS, PEANUTS, PIMENTOS, SORGHUM, SOYBEANS, SPINACH,

ACCEPTED  
1973  
9779-70

20.0%  
80.0%  
100.0%

Do not feed treated vines to dairy animals or animals being finished for slaughter.

**CABBAGE AND OTHER COLE CROPS (Also including Broccoli, Brussels Sprouts, Cauliflower, Collards and Kale):** For control of Armyworms, Fall Armyworms, Cabbage Aphid, Climbing and Surface Feeding Cutworms apply 8 to 10 pounds, for control of Imported Cabbageworms, Green Peach Aphid, Serpentine Leaf Miner, and Thrips apply 10 to 12 pounds, for control of Corn Earworm and Diamondback Moth apply 8 to 20 pounds, for control of Cabbage Looper apply 15 to 20 pounds.

Do not apply after edible parts begin to form or within 35 days of harvest. Treated plants must not be fed to dairy animals or animals being finished for slaughter.

**CORN:** For control of Chinch Bugs and Corn Billbug apply 10 pounds, for control of Armyworm and Budworm apply 15 pounds, for control of Corn Lanternfly and Lesser Cornstalk Borer apply 20 to 23 pounds.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

**COTTON:** For control of Thrips apply 5 to 8 pounds, for control of Fleahoppers apply 5 to 15 pounds, for control of Armyworm, Beet Armyworm, Fall Armyworm, Boll Weevil, Cotton Leafworm, Cutworms, Flea Beetles, Grasshoppers, Lygus Bugs and other Mites, Plant Bugs, Yellow Striped Armyworm, and Western Yellow Striped Armyworm apply 10 to 20 pounds, for control of Cotton Leaf Perforator and Garden Webworm apply 20 to 23 pounds.

Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

**COWPEAS:** For control of Cowpea Curculio apply 8 to 15 pounds. Do not apply to cowpeas to be used as green snaps after pods begin to form.

Do not feed treated vines to dairy animals or animals being finished for slaughter.

**OKRA:** For control of Serpentine Leaf Miner apply 12 to 15 pounds. Do not apply after pods begin to form.

**PEAS:** For control of Leafminer and Lygus Bugs apply 12 to 15 pounds.

Do not apply after pods begin to form if pods are to be used as food, or within 7 days if pods are not to be used as food.

Do not feed vines to dairy animals or animals being fattened for slaughter.

**PEANUTS:** For control of Thrips apply 10 to 12 pounds, for control of Fleahoppers and Leafworm apply 20 to 23 pounds, for control of Velvet Bean Caterpillar apply 8 to 15 pounds, for control of Fall Armyworm apply 10 to 20 pounds, for control of Southern Stink Bug apply 20 to 25 pounds.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

**PIMENTOS:** For control of Flea Beetles, Hornworm, Leafminers, and Pepper Weevil apply 7 to 10 pounds.

Do not apply later than 2 days prior to harvest.

**SMALL GRAINS (Barley, Oats, Rice, Rye, Wheat):** For control of Armyworms, Cutworms (climbing and surface), and Grasshoppers apply 10 to 12 pounds, for control of Chinch Bugs and Rice Stink Bug apply 10 pounds.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

**SORGHUM:** Apply the same as for CORN.

Do not apply later than 28 days before harvest or more than once after heads begin to form. Do not graze dairy animals or animals being finished for slaughter on treated fields. Do not ensilage treated forage.

**SOYBEANS:** For control of Armyworms, Bean Leaf Beetles, Blister Beetle, Bollworm, Corn Earworm, Grasshoppers, Green Clover Worm, and Velvet Bean Caterpillar apply 12 to 20 pounds.

Do not apply later than 21 days before harvest. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

**SPINACH:** Apply the same as for COLE CROPS.

Do not apply later than 21 days before harvest. Do not apply more than once per season.

**STRAWBERRIES:** For control of Crown Borer and Strawberry Weevil apply 10 to 15 pounds.

Do not apply after fruit starts to form.

**TOMATOES:** For control of Armyworm, Blister Beetle, Cabbage Looper, Colorado Potato Beetle, Cutworms, Fleabeetles, Green Chrysanthemum Aphid, Hornworm, Pinworm, Russet Mites, Serpentine Leaf Miner, and Tomato Fruitworm apply 10 to 25 pounds.

Do not apply later than 3 days before harvest.

### DISPOSAL OF CONTAINER AND WASTES

Destroy all containers when empty by punching holes in them and burying with wastes, or by burning. Stay out of smoke and fumes. Do not reuse containers for other purposes. Dispose of wastes by burying in non-crop lands away from water supplies.

### NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instruction, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA REG. NO. 9779-70

NET CONTENTS



MANUFACTURED BY  
**RIVERSIDE CHEMICAL CO.**  
DIVISION OF COOK INDUSTRIES, INC., MEMPHIS, TENN. 38116  
*Riverside serves agriculture. Agriculture serves everyone.*

ACCEPTED  
1973  
9779-70