

RIVERSIDE 5% SEVIN DUST

USEFUL FOR THE CONTROL OF CERTAIN INSECT PESTS OF COTTON, CERTAIN SMALL FRUITS, CERTAIN VEGETABLES, TOBACCO, CERTAIN FORAGE AND FIELD CROPS, GRASSHOPPERS, POULTRY INSECT CONTROL, AND PET INSECT CONTROL.

| | |
|---|------|
| ACTIVE INGREDIENT: | |
| Carbaryl (1-naphthyl N-Methylcarbamate) | 5% |
| INERT INGREDIENTS: | 95% |
| Total | 100% |

CAUTION: KEEP OUT OF REACH OF CHILDREN

HARMFUL IF INHALED OR SWALLOWED. Avoid Breathing of Dust. Do Not Take Internally. Avoid Contact with Skin. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS, AND WATER RECEPTACLES.

Apply this product only as specified on this label.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

NOTE

1. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.
2. This product does not control spider mites. Where infestations are encountered use a recommended miticide.
3. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT: Use of this product in a manner not in accordance with the information on this label may injure or kill humans, domestic animals, wildlife, aquatic life, desirable plants, create excessive and unlawful chemical residues, or produce other undesirable results.

GENERAL INFORMATION

Suggested dosage refers to pounds of RIVERSIDE 5% SEVIN DUST to use per acre of crop treated. Where a dosage rate is indicated use the low rate on small or young plants and the higher rate on large or mature plants. Always use sufficient dust to obtain good coverage of the crop parts to be treated. Apply when insects or their damage appear and repeat applications as necessary or at 7 to 10 day intervals unless shorter intervals are specified in "DIRECTIONS FOR USE."

DIRECTIONS FOR USE

COTTON: For early season control of Thrips, Flea Beetle, Cotton Fleahopper, Cotton Leafworm, and Striped Flea Beetle apply 10 to 20 pounds.

For mid- and late season control of Boll Weevil, Earworm, Fall Armyworm, Cotton Leaf Perforator, Leaf Rollers, Leafhoppers, and Tarnished Plant Bug apply 20 to 40 pounds. Make applications every 5 to 7 days as long as control is necessary.

For control of Pink Egglworm apply 30 to 50 pounds on a 5 to 7 day schedule.

For control of Stink Bugs and Salt Marsh Caterpillar apply 40 pounds.

Aphid populations will be suppressed by repeated applications.

SMALL FRUITS: Blackberries, Boysenberries, Raspberries, Logberries, and Raspberries. Use 40 pounds for Japanese Beetle, Leaf Rollers, Leafhoppers and Raspberry Aphid. Do not apply later than 7 days before harvesting berries.

Grapes: Use 20 to 40 pounds for Grape Leaf Folder, Grape Leafhopper and Grape Leaf Skeletonizer. Apply just before first brood Leaf Folder larvae

Where Cabbage Looper is the principal problem, use an alternative pesticide that is specifically recommended. For control of small Cabbage Loopers after edible portions begin to form use 20 to 40 pounds in a 5 to 7 day schedule.

Do not apply to Chinese cabbage, Collards, Dandelion, Endive (Escarole), Garden Beets, Kale, Leaf Lettuce, Mustard Greens, Parsley, Salsify, Spinach, Swiss Chard or Turnips later than 14 days before harvest, or to Horseradish, Parsnips, Radishes, and Rutabagas later than 3 days before harvest.

Cabbage, Broccoli, Brussels Sprouts, Cauliflower, and Kohlrabi: Use 10 to 20 pounds for Flea Beetles and Harlequin Bug.

Use 20 to 40 pounds for Armyworms, Imported Cabbageworm and Corn Earworm.

Where Cabbage Looper is the principal problem, use an alternative pesticide that is specifically recommended. For control of small Cabbage Loopers after edible portions begin to form use in a 5 to 7 day schedule. Do not apply to these crops later than 3 days before harvest.

TOBACCO: In plant beds (for Tobacco Flea Beetle) - Apply 1 to 2 pounds per 100 square yards when insects appear. Avoid excessive application; plant injury may result. Repeat as needed at 10 to 14 day intervals and just before transplanting.

In fields (broadcast applications for Hornworms and Tobacco Flea Beetles) - Apply 15 to 25 pounds per acre depending on size of the plants. Good coverage is essential. Use the highest dosage only on large plants and repeat as needed. Heavier applications may cause plant injury.

If late applications are necessary allow 3 days before pruning or cutting.

Treated fields may be entered immediately after dust has settled. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

FORAGE AND FIELD CROPS: Alfalfa, Clovers, Cowpeas, Peanuts, and Soybeans - Use 10 to 20 pounds for Blister Beetles and Mexican Bean Beetle.

Use 20 pounds for Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetle, Green Cloverworm, Japanese Beetle, Leafhoppers, Three-toothed Alfalfa Hopper, Thrips and Velvetbean Caterpillar.

Use 20 to 30 pounds for Armyworms, Corn Earworm, Stink Bugs, and Wireworms.

To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.

Forage Grasses and Pasture: Use 20 to 30 pounds for Armyworms and Thrips. For optimum thrips control, water the pasture well before dust application.

Rice (In the Mississippi Delta and Texas): Use 20 to 30 pounds for Armyworms and Stink Bugs. Warn of Dust toxicity to birds and mammals by providing calcium E-3 for the flock or water supply and poultry or animal dust. Do not apply later than 14 days before harvesting rice.

Sorghums (Including Milo and Grain Sorghums): Use 20 to 40 pounds for Armyworms, Corn Earworm, Stink Bugs, and Wireworms. Do not apply to sorghum Midge. For optimum insect control on grain, start making applications to the forming heads. Treatment for Sorghum Midges should be made 3 to 4 days after heads have emerged from boot. Do not apply later than 21 days before harvesting sorghums.

GRASSHOPPERS: Apply 10 to 30 pounds, as often as necessary, to control grasshoppers on the above crops. A 10 to 20 pound dosage is suggested for nymphs on small plants or sparse vegetation in watered areas, field ditchbanks, and borders. A 20 to 30 pound dosage should be used when grasshoppers are mature or material is applied to crops requiring greater coverage. Do not apply

DIRECTIONS FOR USE

COTTON - Use 20 to 40 pounds for Armyworm, Corn Earworm, Stink Bug, and Ear-rolling Plant Bug. Do not apply later than 1 day before harvest.

SMALL FRUITS - Use 20 to 40 pounds for Japanese Beetle, Leafhoppers, and Raspberry Aphid. Do not apply later than 1 day before harvest.

Grapes - Use 20 to 40 pounds for Grape Leaf Folder, Grape Leafhopper, and Grape Leaf Skeletonizer. Apply just before first bud break.

Strawberries - Use 20 to 40 pounds for Meadow Spittlebug, Strawberry Leaf Roller, and Strawberry Weevil. Do not apply later than 1 day before harvesting berries.

VEGETABLES: Asparagus - Use 10 to 20 pounds for Asparagus Beetle on seedlings or sprouts. Do not apply later than 1 day before harvest.

Beans - Use 10 to 20 pounds for Bean Beetles, Bean Leaf Beetle, and Bean Pod Miner. Do not apply later than 1 day before harvest.

Corn - Use 20 to 40 pounds for Corn Earworm, Corn Flea, and Corn Rootworm. Do not apply later than 1 day before harvest.

Cucumbers, Melons - Use 20 to 40 pounds for Cucumber Beetle, Leafhoppers, and Velvetbean Caterpillar. Do not apply later than 1 day before harvest.

Garden Fruits - Use 20 to 40 pounds for Japanese Beetle, Leafhoppers, and Raspberry Aphid. Do not apply later than 1 day before harvest.

Potato Tomatoes - Use 20 to 40 pounds for Colorado Potato Beetle, Leafhoppers, and Colorado Potato Beetle. Do not apply later than 1 day before harvest.

Okra - Use 20 to 40 pounds for Colorado Potato Beetle, Leafhoppers, and Colorado Potato Beetle. Do not apply later than 1 day before harvest.

Carrots, Celery - Use 20 to 40 pounds for Colorado Potato Beetle, Leafhoppers, and Colorado Potato Beetle. Do not apply later than 1 day before harvest.

Spinach - Use 20 to 40 pounds for Colorado Potato Beetle, Leafhoppers, and Colorado Potato Beetle. Do not apply later than 1 day before harvest.

Peas - Use 20 to 40 pounds for Colorado Potato Beetle, Leafhoppers, and Colorado Potato Beetle. Do not apply later than 1 day before harvest.

Onions - Use 20 to 40 pounds for Colorado Potato Beetle, Leafhoppers, and Colorado Potato Beetle. Do not apply later than 1 day before harvest.

Garlic - Use 20 to 40 pounds for Colorado Potato Beetle, Leafhoppers, and Colorado Potato Beetle. Do not apply later than 1 day before harvest.

Herbs - Use 20 to 40 pounds for Colorado Potato Beetle, Leafhoppers, and Colorado Potato Beetle. Do not apply later than 1 day before harvest.

Flowers - Use 20 to 40 pounds for Colorado Potato Beetle, Leafhoppers, and Colorado Potato Beetle. Do not apply later than 1 day before harvest.

77777
ACCEPTED
APR 10 1973
Applied at
per 100 w. apr
Cucumber, Melons
and GARLIC II.

For optimum Triaps control in grasses grown for seed, dust into boot

Rice (In the Mississippi Delta and Texas) - Use 20 to 30 pounds for Armyworms and Stink Bugs. Do not apply later than 1 day before harvest.

Sorghums (Including Milo and Grain Sorghum) - Use 20 to 40 pounds for Armyworm, Corn Earworm, Stink Bug, and Vast Worms. Do not apply later than 1 day before harvest.

GRASSHOPPERS: Apply 10 to 30 pounds as often as necessary to control grasshoppers on the above crops. A 10 to 20 pound rate is suggested for nymphs on small plants or sparse vegetation in waste and range land, ditchbanks, and borders.

POULTRY (On Chickens, Turkeys, Ducks, Geese, Game Birds, and Pigeons) - Litter Treatment: Apply 1 pound per 40 sq. ft. to control Northern Fowl Mite, Chicken Mite, Lice, Fleas, and Bird Louse.

Hand Application to Birds - Use 1 pound to treat 100 birds by means of a sprayer or dust applicator. Do not apply later than 1 day before harvest.

Dust-bath-boxes - Use 2 1/2 pounds per box for 1 to 50 birds. Do not apply later than 1 day before harvest.

PLTS (On Dogs and Cats) - For Ticks and Fleas dust liberally over the entire area including the eyes and ears. Do not apply later than 1 day before harvest.

In or Around Dog and Cat Quarters - For Ticks and Fleas dust liberally in and around dog and cat quarters, kennels, and dog houses by means of a sprayer or dust applicator.

DISPOSAL OF CONTAINER AND WASTES - Empty containers should be disposed of properly.

NOTICE - This product is a restricted use pesticide. It is not to be used on food crops.

NOTICE - This product is a restricted use pesticide. It is not to be used on food crops.

NOTICE - This product is a restricted use pesticide. It is not to be used on food crops.

NOTICE - This product is a restricted use pesticide. It is not to be used on food crops.

NOTICE - This product is a restricted use pesticide. It is not to be used on food crops.

NOTICE - This product is a restricted use pesticide. It is not to be used on food crops.

NOTICE - This product is a restricted use pesticide. It is not to be used on food crops.

NOTICE - This product is a restricted use pesticide. It is not to be used on food crops.

NOTICE - This product is a restricted use pesticide. It is not to be used on food crops.

NOTICE - This product is a restricted use pesticide. It is not to be used on food crops.



RIVERSIDE CHEMICAL CO.