Federal Register / Vol 61 No 144 / Thursday, July 25, 1996 / Notices

registration number product names active ingredients and the specific uses

deleted Users of these products who

desire continued use on crops or sites

38732

being deleted should contact the applicable registrant before October 23 — 1996 to discuss withdrawal of the applications for amendment. This 90

day period will also permit interested members of the public to intercede with registrants prior to the Agency approval of the deleuon.

TABLE 1 — REGISTRATIONS WITH REQUESTS FOR AVENDMENTS TO DELETE USES IN CERTAIN PESTICIDE REGISTRATIONS

EPA Reg stration No	Product Name	Active Ing edient	Delete From Label
000004-00337	Bon de Insect Fog	Resmelhan	Campsites uses
0002600324	SEVIN Brand 90% Technical Carbaryl Innecticide	Carbaryi	Avocados grass for seed maple trees for sap okra, Oyster beds prickly pear cactus
~ 600#35 ~ 60e3a ₃	SBP 1382 Liquid Insectode Spray 0.25% Formula III	Resmethra	Outdoor thermal application in yards patios picnic areas campsites drive-ins horse stables
4000802 2004 423	Lilly/Miller Sevin 5% Dust	Carbaryt	Dog & cat uses
002270-00707	Excelcide Cold Fog	N-Octyl bicycloheptene dicarboximide Piperonyl butoxide Pyrethrins	Mushroom production and proc- essing
003125-00404	DYLOX Technical insectionde	Trichlorion	Livestock uses
004581-00280	TOPSIN M Technical	Thiophanate-methyl	Sugarcane
00458100322	TOPSIN-M 70WP	Thiophanate-methyl	Sugarcane
004581-00352	TOPSIN M 4.5 Turf & Ornamentals Fungicide	Throphanate-methyl	Sugarcane
004581-00372	TOPSIN M 85 WDG	Throphanate-methyl	Sugarcane
004581-00377	TOPSIN-M WSB	Thiophanate-methyl	Sugarcane
	Riverside 5% Sevin Dust	Carbaryl	Use on pe.s
010182-00169	Vernam 10-G Selective Heroxide	Vemolate	Soybeans
010182-00221	Vernam 7-E Selective Herbicide	Vernolate	Soybeans
	Misty Aqua Kill Insec ticide	Resmethan	Greenhouses
0 56228 2 00002	Cas Cartnoge	Carbon, Sodium Nitrate	Pocket gopher

The following Table 2 includes the names and addresses of record for all registrants of the products in Table 1, in sequence by EPA company number

TABLE 2.—REGISTRANTS REQUESTING AMENDMENTS TO DELETE USES IN CERTAIN PESTICIDE REGISTRATIONS

Com- pany No	- Company Name and Address	
000004	Bonide Products Inc., 2 Wurz Avenue, Yorkville NY 13495	
000264	Rhone-Poulenc Ag Company P O Box 12014 T W Alexander Drive Research Triangle Park, NC 27709	ميا ف
000432	AgrEvo Environmental Health, 95 Chestnut Ridge Road, Montvale NJ 07645	
000802	The Chas. H Lilly Co., P.O. Box 83179 Portland OR 97283.	
-002270	The Huge Company Inc., do F.P.J., 1902 Tomahawk Ridge. New Lenox, IL 60451	
003125	Bayer Corporation, P.O. Box 4913, 8400 Hawthorn Rd. Kansas City. MO 64120.	
004581	Elf Atochem North America, Inc. 200 Market Street, Philadelphia, PA 19103.	
009779	Riverside/Terra Corp P O Box 6000 600 4th Street, Sioux City IA 51102	×
010182	Zeneca Ag Products, 1800 Concord Pike P O Box 15458 Wilmington DE 19850	
010807	Amrep, Inc., 990 Industrial Park Drive Marietta, GA 30062.	
056228	U.S Dept. of Agriculture Animal & Plant Health Inspection Service 4700 River Road, Unit 150 Riverdale, MD 20737	

III. Existing Stocks Provisions

The Agency has authorized registrants to sell or distribute product under the previously approved labeling for a period of 18 months after approval of the revision, unless other restrictions

have been imposed as in special review actions

List of Subjects

Environmental protection Pesticides and pesus Product registrations.

Dated July 11 1996.

Frank Sanders

Director Program Management and Support Division Office of Pesticide Programs

(FR Doc. 90-18845 Filed 7-24-96 8 45 am) Billing code 6560-50-F

BEST COPY AVAILABLE

20f6

o EDA		ited States			TORIG APP		Registration				Number
⊗EPA	Environmental Protection Agency Washington, DC 20460 Amendment Other						nt <u></u>	220595			
Application for Pesticide - Section I											
1. Company/Product Number				2. EPA Product Manager 3.					Proposed Classification		
	9779-7	1_	Historian - Conf	D.	Edw	aro	1.		None		Restricted
4. Company/Product (Name) Riverside 5% Sevin Dust				PM# 19							
5. Name and Address of Applicant Include ZIP Code! Riverside / Terra Corp 600 4th St. 7-0. Box 6000: Sioux City, IA 51102				6. Expedited Reveiw., In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No.							
	is a new address			Product Name							
			Sect	tion - I	ſ						
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below.				"Me Too" Application. Other - Explain below.							
Explanation: Use additional page(s) if necessary. (For section I and Section II.) Deletion of Use on Pets and updated label for WPS											
			Sect	ion - i			******			••••	
1. Material This Product Will	Be Packaged In:		STLEMEN NO. C	terms of the second	mayer average of the		THE SECTION OF THE	<u>0. €.12</u>	التسأه غفايا	in sets the t	an er er er er
Child-Resistant Packaging Yes No Ves No			-	Yes No					- ·		
* Certification must } ;ubmitted	If "Yes" Unit Packaging wgt.	No. per container	Packag	je wgt	No, per containe	r	(Other (S	pecify)_		<u> </u>
3. Location of Net Contents	Information	4. Size(s) Reta	il Contair	ner		5. Lo	cation of Label	Directio	ns		
X Label C	. :	2.5, 4, 3				==	on 1	laho 1	A - A	. 4	1
6. Manner in Which Label is Affixed to Product Container				ph Other 5/epse							
			Sect	ion - l'	/					_	
1. Contact Point (Complete	items directly below fo	r identification	of indivi	idual to b	e contacted,	if nec	essary, to proce	ess this	applicati	on.)	ar name komment kom
Name Steven E. Rogosheske Sr. Registration Specialist (7)						lephone (7/6	No. (Inc	lude A	rea Code) 8716		
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowlingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. 6. Date Application Received (Stamped)							-				
2. Signature	B. Title	Registration Specialist				.					
4. Typed Name 5 Steven E. Rogosheske				May 24, 1996							

R°

5% Sevin Dust

N-Methyl Carbamate

Useful for the Control of Certain Insect Pests of Cotton, Certain Small Fruits, Certain Vegetables, Tobacco, Certain Forage and Field Crops, Grasshoppers, and Poultry Insect Control.

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl <u>N</u>-Methylcarbamate)

INERT INGREDIENTS

TOTAL

TOTAL

TOTAL

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS CAUTION HAZARDS TO HUMANS & DOMESTIC ANIMALS

HARMFUL IF INHALED OR SWALLOWED. Avoid breathing of dust. Do not take internally. Avoid contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS, AND WATER RECEPTACLES.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Personal Protective Equipment

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application. This product does not control spider mites. Where infestations are encountered, use a recommended miticide.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT: Use of this product in a manner not in accordance with the information on this label may injure or kill humans, domestic animals, wildlife, aquatic life, desirable plants, create excessive and unlawful chemical residues, or produce other undesirable results.

EPA Reg. No. 9779-74

EPA Est. No. 9779-AL-1

Manufactured by:
RIVERSIDE/TERRA CORPORATION
P.O. Box 6000, Sioux City, IA 51102-6000
Riverside Serves Agriculture. Agriculture Serves Everyone.

NET CONTENTS

9/D03/6

4016

DIRECTIONS FOR USE

it is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statement on this label about personal protective equipment (PPE). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, waterproof gloves, and shoes plus socks.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry place. In case of minor spills, follow all precautions and clean up immediately. Sweep up and dispose of wastes and broken or empty containers in a landfill approved for pesticide use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely emply bag into application equipment. Then dispose of emply bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Suggested dosages refer to pounds of RIVERSIDE 5% SEVIN DUST to use per acre of crop treated. Where a dosage range is indicated, use the lower rate on small or young plants and the higher rate on large or mature plants. Always use sufficient dust to obtain good coverage of the crop plants being treated. Apply when insects or their damage appear and repeat applications as necessary, or at 7 to 10 day intervals, unless shorter intervals are specified in 'DIRECTIONS FOR USE."

DIRECTIONS FOR USE

Cotton: For early-season control of Thrips, Flea beetles, Cotton fleahopper, Cotton leafworm, and Striped blister beetle, apply 10 to 20 pounds.

For mid- and late-season control of Boll weevil, Bollworm, Fall armyworm, Cotton leaf perforator, Leaf rollers, Leafhoppers, and Tarnished plant bug, apply 20 to 40 pounds. Make applications every 5 to 7 days as long as control is necessary.

For control of Pink bollworm, apply 30 to 50 pounds on a 5 to 7 day schedule.

For control of Stink bugs and Salt-marsh caterpillar, apply 40 pounds.

Aphid populations will be suppressed by repeated applications.

SMALL FRUITS

Blackberries, Boysenberries, Dewberries, Loganberries and Raspberries: Use 40 pounds for Japanese beetles, Leaf rollers, Leafhoppers and Raspberry aphid. Do not apply later than 7 days before harvesting berries.

Grapes: Use 20 to 40 pounds for Grape leaf folder, Grape leafhopper and Grape leaf skeletonizer. Apply just before first brood Leaf folder larvae emerge from rolls and as needed for Leafhoppers. Use 40 pounds for Grape berry moth, Japanese beetle, June beetles and Red-banded leaf roller.

Strawberries: Use 20 to 40 pounds for Meadow spittlebug, Strawberry leaf roller and Strawberry weevil. Do not apply later than 1 day before harvesting berries.

VEGETABLES

Asparagus: Use 10 to 20 pounds for Asparagus beetle on seedlings or spears. <u>Do not apply</u> later than 1 day before harvesting asparagus. Use 20 to 40 pounds for Asparagus beetle and Apache cicada on ferns or brush growth in the post-harvest period. Do not apply more often than once every 3 days.

Beans (green beans, lima beans, snap beans, cowpeas and black-eyed peas): Use 10 pounds for Mexican bean beetle, 20 pounds for Bean leaf beetle, Cucumber beetles, Flea beetles, Japanese beetle, Leafhoppers, Velvetbean caterpillar and Westernbean cutworm.

Use 20 to 30 pounds for Armyworms, Corn earworm, Stink bugs and Tarnished plant bug.

50/6

Com: Use 20 to 40 pounds for Corn earworm, Corn rootworm adults, European corn borer, Fall armyworm, Flea beetles, Japanese beetle, Sap beetles and Leafhoppers.

For larvae in whorl (Budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary.

For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control.

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

Cucumber, Melons, Pumpkin and Squash: Use 10 to 20 pounds for Pickleworm and Melonworm; 20 pounds for Cucumber beetles, Flea beetles, Leafhoppers and Squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. RIVERSIDE 5% SEVIN DUST injures watermelons in Florida.

Garden Peas: Use 20 pounds for Leafhoppers. Use 20 pounds for Colorado potato beetle. For Colorado potato beetle control, direct application to weeds in fields of garden peas. This application is intended to help prevent insect contamination in processed peas. Use 20 to 30 pounds for Armyworms and Grasshoppers.

Potato, Tomato, Eggplant and Pepper: Use 10 to 20 pounds for Colorado potato beetle, Flea beetles and Leafhoppers. Use 20 to 30 pounds for European corn borer, Fall armyworm, Lace bugs, Tomato fruitworm, Tomato homworm, Tarnished plant bugs and Stink bugs.

Okra: Use 20 to 40 pounds for Corn earworm and Stink bugs. Repeat at 5 to 7 day intervals or as necessary.

Carrots, Chinese Cabbage, Collards, Dandelion, Endive (Escarole), Garden Beets, Horseradish, Kale, Leaf Lettuce, Mustard Greens, Parsley, Parsnips, Radishes, Rutabagas, Salsify, Spinach, Swiss Chard and Turnips: Use 10 to 20 pounds for Flea beetles, Harlequin bug and Leafhoppers. To control Six spotted leafhoppers (aster yellows vector), apply 20 to 30 pounds in a 5 to 7 day schedule. Use 20 to 40 pounds for Armyworms, Imported cabbageworm, Corn earworm, Tarnished plant bug and Stink bugs. For optimum control of Corn earworm after lettuce heads begin to form, use 40 pounds in a 5 to 7 day schedule.

Where Cabbage looper is the principal problem, use an alternative pesticide that is specifically recommended. For control of small Cabbage loopers after edible portions begin to form, use 20 to 40 pounds in a 5 to 7 day schedule.

Do not apply to Chinese cabbage, Collards, Dandelion, Endive (Escarole), Garden Beets, Kale, Leaf Lettuce, Mustard Greens, Parsley, Salsify, Spinach, Swiss Chard or Turnips later than 14 days before harvest; or to Horseradish, Parsnips, Radishes, and Rutabagas later than 3 days before harvest.

Cabbage, Broccoli, Brussels Sprouts, Cauliflower and Kohlrabi; Use 10 to 20 pounds for Flea beetles and Harlequin bug.

Use 20 to 40 pounds for Armyworms, Imported cabbageworm and Corn earworm.

Where Cabbage looper is the principal problem, use an alternative pesticide that is specifically recommended. For control of small Cabbage loopers after edible portions begin to form, use in a 5 to 7 day schedule. Do not apply to these crops later than 3 days before harvest.

Tobacco: In plant beds (for Tobacco flea beetle), apply 1 to 2 pounds per 100 square yards when insects appear. Avoid excessive application; plant injury may result. Repeat as needed at 10 to 14 day intervals and just before transplanting.

In fields (broadcast applications for Hornworms and Tobacco flea beetles), apply 15 to 25 pounds per acre depending on size of the plants. Good coverage is essential. Use the highest dosage only on large plants and repeat as needed. Heavier applications may cause plant injury.

If late applications are necessary, allow 3 days before priming or cutting.

Treated fields may be entered immediately after dust has settled. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

FORAGE AND FIELD CROPS

Alfalfa, Clovers, Cowpeas, Peanuts and Soybeans: Use 10 to 20 pounds for Blister beetles and Mexican been beetle.

Use 20 pounds for Alfalfa caterpillar, Bean leaf beetle, Cucumber beetles, Green cloverworm, Japanese beetle, Leafhoppers, Three-cornered alfalfa hopper, Thrips and Velvetbean caterpillar.

Use 20 to 30 pounds for Armyworms, Corn earworm, Stink bugs, and Webworms.

To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.

6046

Forage Grasses and Pasture: Use 20 to 30 pounds for Armyworms and Thrips. For optimum Thrips control in grasses grown for seed, direct dust into boot.

Rice (in the Mississippi Delta and Texas): Use 20 to 30 pounds for Armyworms and Stink bugs. Warning! Do not apply before heading if propanil (such as "Stam" F-34) has been or will be applied; plant injury may result. Do not apply later than 15 days before harvesting rice.

Sorghums (including Milo and Grain Sorghum): Use 20 to 40 pounds for Armyworms, Corn earworm, Stink bugs and Webworms. Use 30 pounds for Sorghum midge. For optimum insect control on grain sorghum, direct dust into the forming heads. Treatment for Sorghum midge control should be made 3 to 4 days after heads have emerged from boot. Do not apply later than 21 days before harvesting sorghums.

GRASSHOPPERS: Apply 10 to 30 pounds as often as necessary to control grasshoppers on the above crops. A 10 to 20 pound rate is suggested for nymphs on small plants or sparse vegetation in wasteland, rangeland, ditchbanks, and borders. A 20 to 30 pound dosage should be used when grasshoppers are mature or material is applied to crops requiring greater coverage. Do not apply more than 20 pounds per acre to cucumbers, melons, pumpkins, or squash. Follow the appropriate time limitations as given under the appropriate crop.

POULTRY (On Chickens, Turkeys, Ducks, Geese, Game Birds and Pigeons

Litter Treatment: Apply 1 pound per 40 sq. ft. to control Northern fowl mite, Chicken mite, Lice, Fleas and Bed bugs. Use a scoop and scatter by hand, or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs.

Hand Application to Birds: Use 1 pound to treat 100 birds by means of a shaker can, squeeze bottle, or hand duster to control Northern fowl mite, Chicken mite, Lice and Fleas. Direct dust to vent and fluff areas. Repeat in 4 weeks if necessary. Hand application to birds for Chicken mite and Fleas is a supplement to litter treatment for control of these pests.

Dust-Bath-Boxes: Use 2½ pounds per box for each 50 birds to control Northern fowl mite, Chicken mite, Lice and Fleas. This is a supplement to litter treatment for Chicken mite and Flea control. Con ranges, two natural wallows may be substituted for each dust-bath-box. Do not apply within 7 days of slaughter. Poultry should not be housed in poultry houses within 7 days of slaughter.

PETS (On Dogs and Cats): For Ticks and Fleas ddust liberally over the entire animal, avoiding the eyes, and rub theroughly into the skin. Gomb out dead ticks a few hours ater treatment. Repeat at weekly intervals if needed. Do not treat kitten or nursing puppies:

In or Around Dog and Cat-Quarters: For Ticks-and Fleas dust liberally in and around sleeping quarters, kennels and dog houses by means-of-a shaker-can, hand or power duster. Force dust into cracks and crevices. Repeat as necessary.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.