

**AN EMULSIFIABLE CONCENTRATE CONTAINING 3 POUNDS OF HEPTACHLOR PER GALLON.  
USEFUL FOR THE CONTROL OF CERTAIN INSECT PESTS OF CERTAIN BERRIES, CORN (FIELD), CERTAIN FRUITS, PINEAPPLES, CERTAIN SMALL GRAINS, SOYBEANS, CERTAIN VEGETABLES, ORNAMENTAL TURF, LAWNS, GOLF COURSES, CEMETERIES, AND NURSERIES.**

<b>ACTIVE INGREDIENTS:</b>	
Heptachlor*	32.3%
Related Compounds	12.6%
Aromatic Petroleum Solvent	51.1%
<b>INERT INGREDIENTS:</b>	
TOTAL	100.0%
*Heptachlorotetrahydro-4, 7-methanoindene	
Product contains 3 pounds of Heptachlor per gallon.	

**WARNING: KEEP OUT OF REACH OF CHILDREN**

May be fatal if swallowed! Do not breathe vapor or spray mist. Do not get on skin. In case of contact, wash immediately with soap and water. Do not contaminate feed or foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Keep children and pets off treated areas until material is washed into soil or grass is dry.

**DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.**

**READ ENTIRE LABEL BEFORE USING THIS PRODUCT:** Use of this product in a manner not in accordance with the information on this label may injure or kill humans, domestic animals, wildlife, aquatic life, desirable plants, create excessive and unlawful chemical residues, or produce other undesirable results.

**ANTIDOTE**

**INTERNAL**—Emetic of 1 tablespoonful mustard in tumbler of water. Call a physician.

**EXTERNAL**—In case of spillage on skin, wash promptly with soap and water.

**DISPOSAL OF CONTAINER AND WASTES**

Destroy all containers when empty by punching holes in them and burying with wastes. Do not burn. Do not reuse containers for other purposes. Rinse equipment and containers and dispose of wastes by burying in noncrop lands away from water supplies.

**NOTICE**

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instruction, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

**DIRECTIONS FOR USE**

Dilute RIVERSIDE HEPTACHLOR 3 with sufficient water to give adequate coverage.

Row treatment for corn should be applied in front of planter packing wheels.

Preplant broadcast applications to corn, pineapple, soybeans, small grains, snap beans and lettuce should be worked into the soil immediately after treatment.

Do not apply to berries, fruit, and tomatoes after the edible parts start to form.

On fruit, apply as a soil treatment in late spring or early summer.


Repeat applications to turf as necessary to maintain effective control. Keep children and pets off treated areas until the RIVERSIDE HEPTACHLOR 3 is washed into the soil and the grass is dry.

For control of Cabbage Maggots use ¾ pint of RIVERSIDE HEPTACHLOR 3 per acre and spray in furrow at planting time. RIVERSIDE HEPTACHLOR 3 may be used as a drench at planting time by using ½ pint in 50 gallons of water. Treatment is limited to planting or transplanting only.


For control of Plum Curculio and Conspire Stink Bug, apply uniformly over the soil under the spread of the tree and work into top few inches immediately. Make applications no later than petal fall.

On peaches and pears apply to ground cover in spring before petal fall. On pineapple do not apply foliar application within 60 days of harvest.

Apply as a broadcast treatment only to fruit and vegetables.



EPA REG. NO. 9779-80



MANUFACTURED BY  
**RIVERSIDE CHEMICAL CO.**

A SUBSIDIARY OF COOK INDUSTRIES, INC., MEMPHIS, TN. 38116

Riverside serves agriculture. Agriculture serves everyone.

CROP (OR USE)	INSECT	DOSAGE PER ACRE	
		RIVERSIDE HEPTACHLOR 3	SOIL TREATMENT ONLY
<b>FIELD CROPS</b>		<b>ROW (10"-14")</b>	<b>Broadcast</b>
Corn (field)	Seed Corn Beetle	2 pts.	2 qts.
	Corn Billbug Corn Rootworms, Seed Corn Maggot, Rough Handed Corn Stalk Borer	1½-3 pts.	1½-3 qts.
	Ants, Cutworms, Wireworms, Green June Beetle larvae, Japanese Beetle larvae, White Grubs, White-fringed Beetle larvae, Grape Colaspis	2½-4 pts.	2½-4 qts.
Small Grains (barley, oats, rye, wheat)	White Grubs, Wireworms	Broadcast Soil Treatment Only 3-4 qts.	
Soybeans	Grape Colaspis, Japanese Beetle larvae, Seed Corn Maggot, White Grubs, Wireworms	2½-4 qts.	
<b>VEGETABLES</b>			
beans (snap), lettuce, peppers, tomatoes	Wireworms	3 qts.	
rutabagas	Wireworms	4 qts.	
cabbage	Root Maggots	4-7 qts. Preplant	
	Cabbage Maggot	½ pt.	
<b>BERRIES</b>			
blackberries, blueberries, boysenberries, dewberries, raspberries	Strawberry Root Weevil, Strawberry Root Maggot, Cranberry Root Grub, White Grubs	5½ qts.	
<b>FRUIT</b>			
Pineapple	Ants Anomala Beetle	1¼-2½ qts. Foliar	4 qts. Soil
<b>ORNAMENTAL TURF, LAWNS, GOLF COURSES, CEMETERIES, NURSERIES</b>	Ants, Asiatic Garden Beetle, European Chafer larvae, Imported Fire Ants*, Japanese Beetle larvae, White-fringed Beetle larvae, White Grubs, Wireworms Black Vine Weevil Chiggers, Earwigs, Fleas, Ticks	2 oz. 5 oz. 1 oz.	3 qts. 7 qts. 1½ qts.

\*For mound treatment consult your County Agent or your Cooperative Agricultural Extension Service for specific recommendations in your area.