

SUPPLEMENT TO:

Riverside 7 1/2% SEVIN  
File No. 9779-82  
Dated Feb. 20, 1968

January 16, 1969

DIRECTIONS

BEANS: Apply thoroughly to upper and lower surfaces at first sign of insect infestation before feeding occurs: FOR CONTROL OF MEXICAN BEAN BEETLES, apply 7 1/2 lb. per acre. FOR LEAFHOPPERS, apply 15 lbs. per acre. FOR LYGUS<sup>(b)</sup> AND STINK BUGS, apply 28 lb. per acre. If needed, repeat applications every 7 to 10 days up to and including the day of harvest.

CORN: FOR CONTROL OF CORN EARWORMS, apply 22 1/2 to 25 lbs. per acre. Repeat as necessary for insects attacking silks and ears - apply at 2 to 3 day intervals until silk begins to dry. May be applied up to and including day of harvest on sweet corn for human consumption. Do not apply within 7 days before harvest on corn for use as fodder and forage for dairy and meat animals. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

CUCUMBER, SUMMER SQUASH, CANTALOUPE, WATERMELON: FOR CONTROL OF SQUASH BUGS, CUCUMBER BEETLES, FLEA BEETLES, PICKLEWORM and MELON WORM, apply 13 lbs. per acre. Apply when insects or their damage appear, repeat at 7 to 10 day intervals or as necessary. May be applied up to and including day of harvest.

TOMATO, EGGPLANT, PEPPER: FOR CONTROL OF TOMATO FRUITWORM, FLEA BEETLES, apply at the rate of 22 1/2 to 30 lbs. per acre. FOR CONTROL OF TOMATO HORNWORM and STINK BUGS APPLY at the rate of 30 - 45 lbs. per acre. Apply when insects or their damage appear. Repeat at 7 to 10 day intervals or as necessary, up to and including day of harvest.

POTATOES: Apply on appearance of insects - FOR CONTROL OF COLORADO POTATO BEETLE, apply 7 1/2 to 11 lbs. per acre. FOR LEAFHOPPERS AND FLEA BEETLES, apply 15 to 22 1/2 lbs. per acre. FOR LYGUS BUGS, apply 22 1/2 to 25 lbs. per acre. Repeat if needed every 7 to 10 days up to and including day of harvest.

DOGS AND CATS: FOR CONTROL OF FLEAS AND BROWN DOG TICKS, dust liberally over the animal and rub thoroughly into skin. Apply also to the feet, between toes and on the legs. Dust in and around sleeping quarters and other areas where ticks and fleas occur. Apply at weekly intervals if needed. AVOID TREATMENT OF KITTENS LESS THAN 4 WEEKS OF AGE. DO NOT APPLY TO NURSING PUPPIES.

PET SLEEPING QUARTERS

FLEAS, BROWN DOG TICKS

ACCEPTED