

R I V E R S I D E
75 Peanut Fungicide
COPPER - SULPHUR - PEANUT DUST
Composition

ACTIVE INGREDIENTS:

Copper (From Basic Copper Sulphate) Expressed as Metallic	3.4%
Sulphur (Sulphur 93% Passed 325 Mesh)	75.0%

INERT INGREDIENTS:

21.6%

TOTAL 100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN. SEE RIGHT PANEL FOR ADDITIONAL CAUTIONS.

Effective for the control of Leaf Spot (Cerosporo Arachidicola).

WHEN TO BEGIN

When first spots appear on basal leaves.

NUMBER OF APPLICATIONS


Usually 4. Applied at about 10-day intervals, the first dusting should be about the time of last cultivation.

AMOUNT OF MATERIAL

From 20 to 30 pounds of dust per acre for each dusting depending on size of plants.

CAUTION

Harmful if taken internally. ANTIDOTE: Emetic of Mustard - a cupful, followed with Olive Oil or white of eggs, mucilaginous. AVOID BREATHING DUST. AVOID CONTAMINATION OF FOODSTUFFS. AVOID CONTACT WITH SKIN AND EYES. In case of skin contact, wash with plenty of soap and water. For eyes, flush with plenty of water.



NOT-WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.

Manufactured
MANUFACTURED BY: RIVERSIDE CHEMICAL COMPANY
USDA REG. NO. 9779-

MARKS, MISSISSIPPI

ACCEPTED
May 29, 1968
UNDER FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
EPA REG. NO. 9779-100 (100)