


RIVERSIDE
BORON-COPPER-SULPHUR PEANUT DUST



ACTIVE INGREDIENTS:

Copper (From Basic Copper Sulphate) Expressed as Metallic	-----	3.4%
Sulphur (Sulphur 93% Passed 325 Mesh)	-----	63.0%

INERT INGREDIENTS ----- 33.6%

TOTAL ----- 100.0%

Contains 1.6% Boron

CAUTION: KEEP OUT OF REACH OF CHILDREN. SEE OTHER CAUTIONS ON RIGHT PANEL.

NON-WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Effective for the Control of Leaf Spot (*Cerospora Archidicola*), and to reduce hollow heart in peanuts due to boron deficiencies.

WHEN TO BEGIN

Make a single application of Boron-Copper-Sulphur instead of a fungicidal dust at early bloom stage to dry foliage at 20-30 pounds per acre. Multiple applications or excessive rates may cause crop injury or reduced yields.

NUMBER OF APPLICATIONS

Apply only 1 application of Boron-Copper-Sulphur per year to supply the boron needed to reduce hollow heart in peanuts. Use instead of one of the usual four fungicidal applications. Other fungicidal dusts applied at about 10 day intervals, the first dusting should be about the time of last cultivation.

AMOUNT OF MATERIAL

From 20 to 30 pounds of dust per acre depending on size of plants.

CAUTION

Harmful if taken internally. If taken internally, induce vomiting by giving mustard and water and tickling the throat. Give large doses of magnesia or lime water, followed with doses of Epsom Salts.

AVOID BREATHING DUST. AVOID CONTAMINATION OF FOODSTUFFS. AVOID CONTACT WITH SKIN AND EYES. In case of skin contact, wash with plenty of soap and water. For eyes, flush with plenty of water.

CAUTION

Do not apply more than 1 application of fertilizer or insecticide containing boron per year as crop injury may result.

Keep out of lakes, streams, and ponds. Apply this product only as specified on this label. Do not apply when weather conditions favor drift of spray from areas treated. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

MANUFACTURED BY: RIVERSIDE CHEMICAL COMPANY MARKS, MISSISSIPPI
USDA REG. NO. 9779-GG

