

RIVERSIDE TOXON 63

AN EMULSIFIABLE CONCENTRATE CONTAINING 6 POUNDS OF TOXAPHENE AND 3 POUNDS OF METHYL PARATHION PER GALLON. USEFUL FOR THE CONTROL OF CERTAIN INSECT PESTS OF COTTON, RICE, SMALL GRAINS (Barley, Oats, Rye, Wheat), AND SOYBEANS.

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING

(1) Wear water-proof pants, coat, hat, rubber boots or rubber overshoes. (2) Wear safety goggles. (3) Wear mask or respirator approved by the U.S. Bureau of Mines for methyl parathion protection. (4) Wear heavy duty, natural rubber gloves.

WORK SAFETY RULES

Keep all unprotected persons and children away from treated area or where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to methyl parathion-toxaphene combination and describe his condition. After first aid is given (see First Aid Treatment Section) and if a doctor cannot come, take victim to clinic or hospital.

IMPORTANT! Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smoking, eating or drinking.

AFTER WORK, take off all work clothes and shoes. Shower, using soap and water. Wear only clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirator should be cleaned and filter replaced according to instructions included with respirator.

POISON SIGNS (Symptoms)

Methyl parathion-toxaphene combination is a very dangerous poison. It rapidly enters the body on contact with all skin surfaces and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment or they may die.

Some of the signs and symptoms of poisoning are: Headache, nausea, vomiting, cramps, weakness, blurred vision, pin-point pupils, tightness in chest, labored breathing, nervousness, sweating, watering of eyes, drooping or twitching of mouth and nose, muscle spasms and coma.

FIRST AID TREATMENT

Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to methyl parathion-toxaphene combination and describe his condition.

IF BREATHING HAS STOPPED, start artificial respiration immediately and maintain until doctor sees victim.

IF SWALLOWED and victim is awake (conscious) make him vomit quickly. Induce vomiting by sticking finger down throat or by giving scapy or strong salty water to drink. Repeat until vomit is clear. Never give anything by mouth to an unconscious person. Have victim lie down and keep quiet. See doctor immediately.

IN CASE OF CONTACT, immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. See doctor immediately.

NOTE TO PHYSICIAN

In this product mixture, methyl parathion is the dominant toxic ingredient exerting muscular-like effects, that is, stimulation of the parasympathetic nervous system. Poisoning victims should be treated promptly with atropine to the limit of the victims' tolerance to relieve symptoms and to improve chances of survival. 2-PAM may be administered in conjunction with atropine. The administration of oxygen is helpful. Morphine should not be given to persons suspected of methyl parathion poisoning. Toxaphene produces central nervous system excitation which ordinarily responds to the barbituric acid derivatives, but this drug should be considered only if complete atropinization fails to relieve symptoms.

POST TREATED AREA

Consult your State Agricultural Extension Service or Experiment Station regarding posting treated areas.

ACTIVE INGREDIENTS:

Toxaphene	52.8%
O,O-dimethyl O-p-nitrophenyl phosphorothioate**	26.4%
Aromatic Petroleum Solvent	15.8%
INERT INGREDIENTS:	5.0%
Total	100.0%

*Technical Chlorinated Camphene, Chlorine Content 67-69%
**Methyl Parathion
Contains 6 pounds of Toxaphene and 3 pounds of Methyl Parathion per gallon


STOP -- READ LABEL BEFORE USING!

POISON
SEE SIDE
PANELS FOR
ANTIDOTE AND
PRECAUTIONS




DANGER
KEEP OUT OF
THE REACH OF
CHILDREN

NOT FOR HOME USE




CAN KILL IF SWALLOWED

This product can kill you if swallowed even in small amounts. Spray mist may be fatal if swallowed.



CAN KILL BY SKIN CONTACT

This product can kill you if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes).



CAN KILL IF BREATHED

This product can kill you if vapors or spray mist are breathed.

POISONOUS TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur.

HIGHLY TOXIC TO BEES

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

GENERAL INFORMATION

RIVERSIDE TOXON 63 is useful for the control of certain insect pests of rice, certain small grains (barley, oats, rye, wheat), and soybeans.

RIVERSIDE TOXON 63 may be applied by ground sprayers or by adding the RIVERSIDE TOXON 63 in water to form an emulsion spray. Make the spray solution half fill the spray tank with water, add proper amount of RIVERSIDE TOXON 63 and then fill the spray tank while still agitating. Continue to agitate during use.

When applying with ground sprayers calibrate so as to apply the recommended amount of RIVERSIDE TOXON 63 in not less than 3 gallons of water. Airplanes should apply the recommended amount of RIVERSIDE TOXON 63 in not less than 1 gallon of water per acre. Always use sufficient water to insure adequate coverage of the crop foliage.

Where a dosage range is given use the lower rate on small or young plants and the higher rate on large or mature plants. Use the higher rate on large or mature plants with heavy insect infestations.

The hazards and precautions for handling RIVERSIDE TOXON 63 in tankers are equally applicable to it after dilution with water for spray.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR FLAME.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT: Use in a manner not in accordance with the information on this label may kill humans, domestic animals, wildlife, aquatic life, desirable plants, and produce other undesirable effects and unprofitable chemical residues, or produce other undesirable effects.

DIRECTIONS FOR USE

COTTON — For control of Aphids, Fleahoppers, Cotton Leafworm, Cotton Bollworm, Garden Webworm, Grasshoppers, Lygus Bugs and other Cotton Pests apply 1 1/2 to 2 1/2 pints of RIVERSIDE TOXON 63 per acre per application. For control of Bollworms and Spiny Buds apply 1 pint per acre per application. For control of Beet Armyworm, Tobacco Budworm, Green Leafhopper, Cotton Leaf Perforator, Cotton Bollworm, Saltmarsh Caterpillar, and Yellow-striped Armyworm apply 1 to 6 pints per acre per application. For control of Cutworms and Flea Beetles apply 1 to 6 pints per acre per application. Applications should be made at intervals until control is obtained.

Do not apply RIVERSIDE TOXON 63 within 7 days of handpicking. Workers entering cotton fields within 24 hours after application of RIVERSIDE TOXON 63 should wear protective clothing.

RICE — For control of Stink Bug, use 1 1/2 pints per acre. Do not apply after harvest. Do not feed treated forage to dairy animals or animals intended for slaughter.

SMALL GRAINS — (Barley, Oats, Rye, Wheat) — For control of Armyworm apply 1/2 to 1 1/4 pints per acre per application, depending upon the infestation. For control of Armyworms (up to 3rd instar) and Lygus apply 1 1/2 pints of RIVERSIDE TOXON 63 per acre per application.

Do not apply RIVERSIDE TOXON 63 to small grains later than 15 days before harvest.

59%
SHOOT AT FOR REFLECTIONS REPRODUCE IN 3 COLORS: GREEN (191 LINE NO. GF/3)
WARM RED
BLACK

ACCEPTED

DEC 27 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

9779-134

IN PER GALLON.
, AND SOYBEANS.

FORMATION

ION 63 is useful for the control of certain insect pests of cotton, all grains (barley, oats, rye, wheat), and soybeans.

ION 63 may be applied by ground sprayers or by airplanes, mixed with water to form an emulsion spray solution. To use, fill the spray tank with water, agitate, add the recommended amount of RIVERSIDE TOXON 63 in gallon of water per acre. Always use sufficient water for adequate coverage of the crop foliage.

For ground sprayers calibrate so as to apply the recommended amount of RIVERSIDE TOXON 63 in not less than 3 gallons of water per acre. Do not apply the recommended amount of RIVERSIDE TOXON 63 in less than 3 gallons of water per acre. Always use sufficient water for adequate coverage of the crop foliage.

A range is given use the lower rate on small or young plants or light infestations; use the higher rate on large or mature plants or heavy infestations.

Precaution for handling RIVERSIDE TOXON 63 in this container: applicable to it and dilution with water for spray application.

E. POUR, SPILL, OR STORE NEAR HEAT OR OPEN

LABEL BEFORE USING THIS PRODUCT: Use of this product in accordance with the information on this label may injure or kill domestic animals, wildlife, aquatic life, desirable plants, create undesirable chemical residues, or produce other undesirable results.

S FOR USE

For control of Aphids, Fleahoppers, Cotton Leafworm, Fall Armyworm, Webworm, Grasshoppers, Lygus Bugs and other Midges. Red and Blue apply 1 1/2 to 1 1/4 pints of RIVERSIDE TOXON 63 per acre per application. For control of Bean Beetles and Stink Bugs apply 1 1/2 to 3 pints per application. For control of Beet Armyworm, Bill Worm or Worm, Caterpillar, Loopers, Cotton Leaf Perforator, Cutworms, Feathered Caterpillar, and Yellow-striped Armyworm apply 2 1/2 to 3 pints per application. For control of Cutworms and Pink Beetles apply 1 1/2 to 3 pints per application. Applications should be made at 3 to 5 days until control is obtained.

RIVERSIDE TOXON 63 within 7 days of handpicking or harvest. For control of Stink Bug, use 1 1/2 pint per acre. Do not apply within 15 days of harvest. Do not feed treated forage to dairy animals or animals being slaughtered.

For control of Stink Bug, use 1 1/2 pint per acre. Do not apply within 15 days of harvest. Do not feed treated forage to dairy animals or animals being slaughtered.

15 — (Barley, Oats, Rye, Wheat) — For control of Aphids (Green and Black), Armyworms (up to 3rd instar) and Leafhoppers apply 1 1/2 to 3 pints of RIVERSIDE TOXON 63 per acre per application.

RIVERSIDE TOXON 63 to small grains later than 15 days prior to harvest.

SOYBEANS — For control of Corn Earworm, Fall Armyworm, Green Cloverworm, Mexican Bean Beetle, Stink Bug, Three-Cornered Alfalfa Hopper, and Violet Bean Caterpillar apply 1 quart of RIVERSIDE TOXON 63 per acre.

Do not apply RIVERSIDE TOXON 63 to soybeans later than 21 days prior to harvest. Do not apply more than two applications per growing season. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

DISPOSAL OF CONTAINER AND WASTES

This container should not be reused for ANY purpose. When empty, it should be rendered useless by punching large holes in the top, bottom, and sides with an instrument such as an axe. It should then be mashed as flat as possible and buried underground as deep as practical, with at least 18 inches of soil covering. This should be done in an isolated area, away from habitation, and in a location which would not drain into or cause pollution of lakes, streams, or ponds. Any spray wastes or spillage resulting from the application of this material or the destruction of the container should be neutralized with a caustic solution and buried in a pit with the disposed container.

NOTICE

RIVERSIDE CHEMICAL COMPANY warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to RIVERSIDE CHEMICAL COMPANY, and buyer assumes the risk of any such use.

EPA REG. NO. 9779-134

NET CONTENTS



MANUFACTURED BY
RIVERSIDE CHEMICAL CO.
A SUBSIDIARY OF ECOLAB INDUSTRIES, INC., MEMPHIS, TN 38114
Riverside serves agriculture. Agriculture serves everyone.



COLORS: GREEN (1PI LME No. CF/3y9/05) - SEE TISSUE OVERLAY FOR COLOR DISTRIBUTION.
WARM RED
BLACK
/ 11/11 / 11/11 / 11/11 /