RIVERSIDE 5% SEVINO GRANULAR INSECTICIDE

SEVIN is the registered trade mark of Union Carbide Corporation for carbaryl insecticide.

CAUTION: KEEP OUT OF REACH OF CHILDREN AND ANIMALS

HARMFUL IF INHALED OR SWALLOWED. Avoid Prolonged or Repeated Breathing of Dust. Do Not Take Internally. Avoid Prolonged or Repeated Contact with Skin.

Wear regular long-sleeved work clothing. Change to clean clothing daily.

Wash hands and face before eating.

Wash thoroughly after handling.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS, AND PONDS.

ANTIDOTE: If taken internally induce vomiting and call a physician at once.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT: Use of this product in a manner not in accordance with the information on this label may injure or kill humans, domestic animals, wildlife, aquatic life, desirable plants, create excessive and unlawful chemical residues, or produce other undesirable results.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION:

Tolerances established under the Federal Food and Cosmetic Act permit the sale of crops bearing probable SEVIN residues when SEVIN is used in accordance with label instructions.

If SEVIN Insecticide is used in accordance with label directions, these crops, including corn forage and rice straw, may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat.

DIRECTIONS FOR USE

For use on corn and rice: Recommended dosages refer to pounds of Riverside 5% GRANULAR SEVIN INSECTICIDE per acre. Use suitable ground or air equipment to insure good coverage. Apply when pests appear in damaging numbers and repeat 7 to 14 days later if necessary.

CORN:

Use 30 to 40 pounds for first-generation European corn borer larvae in whorls. No post-treatment time limitations on applications to sweet corn and corn forage and fodder.

RICE:

In California, use 40 pounds for tadpole shrimp. For optimum control, apply to water when pest first appears. Do not apply within 14 days of harvest.

WARNING: To avoid plant injury, do not apply within 15 days of propanil (such as Riverside's "Propanilo", or "Stam") use.

LAWNS:

For control of chinch bugs and certain other insects cut the lawn close, remove clippings and water before treatment to bring chinch bugs to the surface.

Treat the entire lawn evenly and thoroughly at the rate of 4 pounds of RIVER-SIDE 5% SEVIN GRANULAR INSECTICIDE per 1000 square feet. Apply with a spreader for uniform coverage. Work material in with a coarse water spray after treatment. RIVERSIDE 5% SEVIN GRANULAR INSECTICIDE will aid in the control of (all armyworms, sod webworms or lawn moths, ants and millipedes.

CONTAINER DISPOSAL:

Do not reuse empty container. Burn immediately when empty. Stay out of smoke.

NOTICE:

Riverside Chemical Company warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably forseeable to Riverside Chemical Company, and buyer assumes the risk of any such use.

EPA REG. NO. 9779-156

NET CONTENTS____



MANUFACTURED BY

miverside chemical co.

DIVISION OF COOK INDUSTRIES, INC., MEMPHIS, TENN. 38116

Plants at Marks and Canton, Miss., Blytheville and Pine Biuff, Ark:, Montgomery and Decatur, Ala., Rayville, La. and Jackson, Tenn.

75

ACCEPTED

JUL 3 1 1972
UNDER THE HIDERAL INSECTICIDE FUNDICIDIL AND MODERN CIDIE ACT FOR ECONOMIC POSSON FEB. ED ONDER NO LITELES SUBJECT TO ATTACHED COMMENTS.

921-666

The Lo