

## MIXING DIRECTIONS

Always have at least half the required amount of water in the tank before adding the spray concentrate. Have pump overflow operating to assist in mixing.

## DIRECTIONS FOR USE

Nozzles should be set to spray so they are at least 8 to 10 inches above the highest growth when spraying.

**SMALL GRAINS:** For control of fall armyworms, yellow striped armyworms, grasshoppers, chinch bugs and climbing and surface feeding cutworms on rice, wheat, oats, barley and rye, apply Toxaphene EC at the rate of 2 to 2 2/3 pints per acre in sufficient water to give good coverage. Use the higher dosage for chinch bug and for climbing and surface feeding cutworms. Do not apply more than once after heads are formed. Do not apply within 7 days of harvest on wheat, rye, and oats. Do not apply within 14 days of harvest on rice and barley. Do not feed treated forage to dairy animals or animals being finished for slaughter.

**COTTON INSECTS:** For early season control make two or three applications at seven-day intervals starting at the four-leaf stage, earlier if cutworm or thrips damage is apparent. Use One Pint per acre for the first application. For cutworm control apply one quart per acre as soon as cutworm damage appears or as soon as cotton is up to a stand where damage is expected.

MAY 9 1968  
UNDER FEDERAL INSECTICIDE  
FUNCTION AND REGISTRATION ACT  
FOR ECONOMIC POISON REGISTERED  
FD UNDER NO. 3677-52 SUBJECT  
TO ATTACHED COMMENTS.



# Toxaphene 6-EC

A Toxaphene Spray Concentrate designated for low gallonage spraying for control of Armyworms, Thrips, Plant Bugs, Cotton Fleahoppers, Grasshoppers and certain other insects.

ACTIVE INGREDIENTS:	
Toxaphene*	60.0%
Aromatic Petroleum Distillates	35.0%
INERT INGREDIENTS:	
TOTAL	
*Technical Chlorinated Camphene, 67-69% Chlorine	
SOLD FOR INSECTICIDAL PURPOSES	

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
See Additional Caution On Side Panel

NON-WARRANTY: Atkins Phelps, Inc. makes no warranty of any kind, expressed or implied, on the use of this material and assumes no responsibility whatsoever for any personal injury, property damage, or other type of loss resulting from use and handling of this material. User assumes all risk for use and handling whether in accordance with directions or not. Buyers accept and use this product under these conditions.

Net Contents                      Gallons  
Manufactured By

Atkins Phelps, Inc.

CHEMICAL DIVISION  
NORTH LITTLE ROCK, ARKANSAS

GRA  
treat  
fenc  
bree  
Toxa  
  
May  
get  
tami  
tic  
anir  
app  
fish  
this  
tain  
crop  
  
INTI  
wat  
mec  
  
EXT

UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON REGISTERED  
 UNDER NO. 36775 SUBJECT  
 TO ATTACHED COMMENTS.

9779-159



# Toxaphene 6-EC

A Toxaphene Spray Concentrate designated for low gallonage spraying for control of Armyworms, Thrips, Plant Bugs, Cotton Fleahoppers, Grasshoppers and certain other insects.

<b>ACTIVE INGREDIENTS:</b>	
Toxaphene*	60.0%
Aromatic Petroleum Distillates	35.0%
<b>INERT INGREDIENTS:</b>	5.0%
<b>TOTAL</b>	<b>100.0%</b>

\* Technical Chlorinated Camphene, 67-69% Chlorine  
 SOLD FOR INSECTICIDAL PURPOSES

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
 See Additional Caution On Side Panel

**NON WARRANTY:** Adkins Phelps, Inc. makes no warranty of any kind, expressed or implied, on the use of this material and assumes no responsibility whatsoever for any personal injury, property damage, or other type of loss resulting from use and handling of this material. User assumes all risk for use and handling whether in accordance with directions or not. Buyers accept and use this product under these conditions.

Net Contents      Gallons  
 Manufactured By

*Adkins Phelps, Inc.*

CHEMICAL DIVISION  
 NORTH LITTLE ROCK, ARKANSAS

## DIRECTIONS FOR USE (Continued)

**GRASSHOPPERS:** The best control of grasshoppers is obtained by treating when they are small. Spray along ditch banks, turn row fences, weed patches and other similar areas where grasshoppers breed. When young grasshoppers appear in numbers, use 3 Part Toxaphene 6-EC diluted with sufficient water per acre.

## WARNING

May be fatal if swallowed. DO NOT breathe spray mist. Do Not get on skin, in eyes or on clothing. Wash thoroughly. Do Not contaminate feed or foodstuffs. Keep away from children and domestic animals (particularly dogs). Do Not pasture dairy animals or animals being fattened for slaughter in treated fields. Do Not apply nor allow to drift where livestock are grazing. To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material. Do Not store or handle near open flame. Keep container tightly closed when not in use. Use only on recommended crops and at recommended dosages.

## ANTIDOTE

**INTERNAL:** Give One to Two Tablespoons of salt in glass of warm water, and repeat until vomit fluid is clear. Call a physician immediately.

**EXTERNAL:** Wash with Soap and water.

Registration No. 36775